

Labs

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC with platelet and differential STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Comprehensive metabolic panel STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> Sedimentation rate STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd
<input type="checkbox"/> C-reactive protein STAT, Status: Future, Expires: S+366, Clinic Collect	PRN	S	Until discont'd

Nursing Orders

	Interval	Defer Until	Duration
<input type="checkbox"/> Insert Peripheral IV or Access Portacath Details	Once	S	1 treatment
<input type="checkbox"/> Vital signs - T/P/R/BP Per Unit Protocol At start of infusion: obtain vital signs every 15 minutes x 4, then every 30 minutes.	Once	S	1 treatment
<input type="checkbox"/> Nursing communication: Observe patient for 30 minutes Details	Once	S	1 treatment
<input type="checkbox"/> Nursing communication: Monitor patient for any signs of MINOR INFUSION REACTION Monitor patient for any signs of infusion reaction. For MINOR INFUSION REACTIONS (fever, chills, pruritis, urticaria, chest pain, mild hypotension, hypertension, or mild dyspnea) stop or slow infusion rate by 50%. If symptoms do not resolve within 5-10 minutes give Benadryl as ordered. Notify MD if symptoms do not resolve within 5-10 minutes. With MD approval, give Solu-medrol as ordered. When symptoms resolve, restart medication at 25% of the original rate. After 15-30 minutes gently increase rate and monitor patient closely until completion.	Once	S	1 treatment
<input type="checkbox"/> Nursing communication: Monitor patient for signs of SEVERE INFUSION REACTION Monitor patient for signs of SEVERE INFUSION REACTION: severe dyspnea, wheezing, stridor, angioedema, or marked hypotension. For severe infusion reaction take the following steps; STOP Infusion and start Normal Saline solution, Notify physician and administer Benadryl as ordered, Monitor vital signs, Administer Epinephrine as ordered. Consult MD for further orders.	Once	S	1 treatment
<input type="checkbox"/> sodium chloride 0.9% flush 10 mL 10 mL, intravenous, PRN, line care, Starting S, For 10 Doses	Once	S	1 treatment
<input type="checkbox"/> HEParin, porcine (PF) 100 unit/mL injection 500 Units 500 Units, intra-catheter, once, Starting S, For 1 Doses ONLY IF PATIENT HAS PORTACATH: Flush portacath before removing needle from portacath.	Once	S	1 treatment
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Once	S	1 treatment
<input type="checkbox"/> Discontinue IV Details	Once	S	1 treatment

Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 500 mg 500 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Once	S	1 treatment
<input type="checkbox"/> diphenhydramine (BENADRYL) tablet 25 mg 25 mg, oral, once, Starting S, For 1 Doses Give 30 minutes prior to start of infusion.	Once	S	1 treatment

Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> ustekinumab (STELARA) in sodium chloride 0.9% 250 mL IVPB	Once	S	1 treatment

Medications (continued)

	Interval	Defer Until	Duration
intravenous, for 60 Minutes, once, Starting S, For 1 Doses Use an in-line, low protein-binding filter.			
< 55 kg: 260 mg as single dose			
> 55 kg to 85 kg: 390 mg as single dose			
> 85 kg: 520 mg as single dose			

Hypersensitivity

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, oral, every 6 hours PRN, mild pain (score 1-3), headaches, fever, Starting S	PRN	S	Until discontinuation
<input type="checkbox"/> diphenhydramine (BENADRYL) injection 25 mg 25 mg, intravenous, once PRN, itching, for mild to moderate infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> diphenhydramine (BENADRYL) injection 50 mg 50 mg, intravenous, once PRN, itching, for severe infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> epinephrine (ADRENALIN) 1 mg/1 mL injection 0.5 mg 0.5 mL, subcutaneous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection 100 mg 100 mg, intravenous, once PRN, anaphylaxis, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> methylprednisolone sodium succinate (Solu-MEDROL) injection 40 mg 40 mg, intravenous, once PRN, for infusion reaction, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> ondansetron (ZOFRAN) injection 4 mg 4 mg, intravenous, once PRN, nausea, vomiting, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> promethazine (PHENERGAN) tablet 25 mg 25 mg, oral, every 6 hours PRN, nausea, vomiting, Starting S	PRN	S	Until discontinuation
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL 250 mL, intravenous, once PRN, Hydration, Starting S, For 1 Doses	PRN	S	Until discontinuation
<input type="checkbox"/> sodium chloride 0.9% infusion 500 mL 500 mL, intravenous, once PRN, Hydration, Starting S, For 1 Doses	PRN	S	Until discontinuation