

Nursing Orders

	Interval	Defer Until	Duration
☐ Insert Peripheral IV or Access Portacath Details	Once	S	1 treatment
☐ sodium chloride 0.9 % flush 10 mL 10 mL, intravenous, PRN, line care, Starting S	Once	S	1 treatment
☐ sodium chloride 0.9 % infusion 250 mL 250 mL, intravenous, at 30 mL/hr, PRN, Flush bag, Starting S	Once	S	1 treatment
☐ Discontinue IV Details	Once	S	1 treatment

Pre-Medications

	Interval	Defer Until	Duration
☐ methylPREDNISolone sodium succinate (Solu-MEDROL) injection 100 mg 100 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Once	S	1 treatment
☐ metoclopramide (REGLAN) 5 mg/5 mL solution 20 mg 20 mg, oral, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Once	S	1 treatment
☐ diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, intravenous, once, Starting S, For 1 Doses Administer 30 minutes before infusion.	Once	S	1 treatment

Medications

	Interval	Defer Until	Duration
☐ valproate (DEPACON) 1,000 mg in sodium chloride 0.9 % 100 mL IVPB 1,000 mg, intravenous, for 60 Minutes, once, Starting H+30 Minutes, For 1 Doses	Once	S	1 treatment