Alteplase (tPA) Reversal Management - Symptomatic ICH Management within 24 Hours of alteplase (tPA) Administration [4112] PROVIDER: DISCONTINUE ALTEPLASE (TPA) INFUSION IMMEDIATELY

Nursing	
Vital Signs and HM Stroke Change Scale (HMSCS) (Single Response)	
) Vital Signs and HM Stroke Change Scale (HMSCS) Every 15 Minutes	
[] Vital signs - T/P/R/BP	STAT, Per unit protocol
[] HM Stroke Change Scale (HMSCS)	STAT, Every 15 min,
	Perform NIH Stroke Change Scale if/when an increase of 2 or more points in the HM Stroke Change Scale and notify provider.
) Vital Signs and HM Stroke Change Scale (HMSCS) Every 30 Minutes	
[] Vital signs - T/P/R/BP	STAT, Every 30 min
[] HM Stroke Change Scale (HMSCS)	STAT, Every 30 min,
	Perform NIH Stroke Change Scale if/when an increase of 2 or more points in the HM Stroke Change Scale and notify provider.
) Continue with Post-Alteplase Vital Signs and HM Stroke Change Scale (HMSCS)	
[] Continue with Post-Alteplase Vital Signs and HM Stroke Chage Scale (HMSCS)	Routine, Until discontinued, Starting S For Until specified
Notify	
X] Notify Physician for vitals:	Routine, Until discontinued, Starting S Temperature greater than: Temperature less than: Systolic BP greater than: 180 Systolic BP less than: Diastolic BP greater than: 105 Diastolic BP less than: MAP less than: Heart rate greater than (BPM): Heart rate less than (BPM): Respiratory rate greater than: Respiratory rate less than: SpO2 less than:
Medications	
Medications (Single Response)	
) tranexamic acid (CYCLOKAPRON) IVPB	1,000 mg, intravenous, for 10 Minutes, once, Starting S, For 1 Doses
) Aminocaproic Acid - Bolus and Infusion	
[] aminocaproic acid (AMICAR) sodium chloride 0.9 % (loading dose)	5 g, intravenous, for 60 Minutes, once, For 1 Doses
[] aminocaproic acid (AMICAR) infusion	1 g/hr, intravenous, continuous, Starting H+60 Minutes, For 8 Hours
_abs	
_aboratory - STAT	
X] Type and screen	Once

IVI Drothrombia time with IND	CTAT For 4 Occurrences
[X] Prothrombin time with INR	STAT For 1 Occurrences STAT For 1 Occurrences
[X] Partial thromboplastin time, activated[X] Fibrinogen	STAT For 1 Occurrences
[] Glucose level	STAT For 1 Occurrences
	CTATT OF T COCUMENOUS
Blood Products	
[] Red Blood Cells	
[] Prepare RBC	Routine
	Transfusion Indications:
	Transfusion date:
[] Transfuse RBC	Routine
[1 Plotolet phorosis	Transfusion duration per unit (hrs):
[] Platelet pheresis [] Prepare platelet pheresis	Routine
[] Frepare placelet priefesis	Transfusion Indications:
	Transfusion date:
[] Transfuse platelet pheresis	Routine
	Transfusion duration per unit (hrs):
[] Fresh Frozen Plasma	
[] Prepare fresh frozen plasma	Routine
	Transfusion Indications:
I 1 Transfuse fresh frezen plasma	Transfusion date: Routine
[] Transfuse fresh frozen plasma	Transfusion duration per unit (hrs):
[X] Cryoprecipitate	Transitusion duration per unit (mo).
[X] Prepare cryoprecipitate	Routine
	Transfusion Indications: Other (Specify)
	Other indication: Symptomatic alteplase ICH management
	Transfusion date:
[X] Transfuse cryoprecipitate	Routine
	Transfusion duration per unit (hrs):
IV Fluids	
[] sodium chloride 0.9 % infusion	250 mL, intravenous, at 30 mL/hr, continuous
[]	Administer with blood
Consults	
Consults - STAT	
[X] Consult Hematology	Reason for Consult?
	Patient/Clinical information communicated?
IVI Consult Nourceurgery	Patient/clinical information communicated?
[X] Consult Neurosurgery	Reason for Consult? Patient/Clinical information communicated?
	Patient/Clinical information communicated? Patient/clinical information communicated?
	r attempolitical information communicated: