

General

Medications

Pharmacy Consult (Selection Required)

Pharmacy consult to monitor ibutilide (CORVERT) therapy STAT, Until discontinued, Starting S

ibutilide (CORVERT): patient weight greater than or equal to 60 kg

ibutilide (CORVERT) bolus 1 mg, intravenous, for 10 Minutes, once, For 1 Doses RESTRICTED to Cardiology specialists. Are you a Cardiology specialist or ordering on behalf of one?

ibutilide (CORVERT): patient weight less than 60 kg

ibutilide (CORVERT) bolus 0.01 mg/kg, intravenous, for 10 Minutes, once, For 1 Doses RESTRICTED to Cardiology specialists. Are you a Cardiology specialist or ordering on behalf of one?

Electrolyte Replacement

if magnesium level less than 3 mEq/L 2 g, intravenous, at 50 mL/hr, for 2 Hours, once, For 1 Doses

if potassium level less than or equal to 4 mEq/L (CENTRAL LINE) 20 mEq, intravenous, for 60 Minutes, once, For 1 Doses For central line only

potassium chloride IVPB (PERIPHERAL LINE) 10 mEq, intravenous, for 60 Minutes, every 1 hour, For 2 Doses

Nursing

Vital signs

Vital signs Routine, Once For 1 Occurrences Obtain baseline prior to procedure.

Vital signs Routine, Every 15 min Every 15 minutes x 4 occurrences, followed by every 30 minutes x 2 hours, followed by every hour x 2 hours.

Telemetry **"And" Linked Panel**

Telemetry monitoring Routine, Continuous Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box) Reason for telemetry: Can be off of Telemetry for tests and baths? Yes

Telemetry Additional Setup Information Routine, Continuous High Heart Rate (BPM): 120 Low Heart Rate(BPM): 50 High PVC's (per minute): 10 High SBP(mmHg): 175 Low SBP(mmHg): 100 High DBP(mmHg): 95 Low DBP(mmHg): 40 Low Mean BP: 60 High Mean BP: 120 Low SPO2(%): 94

Nursing care

Ibutilide Protocol: Cardiologist, Cardiologist Fellow, or Cardiologist Midlevel must be present during ibutilide administration Routine, Until discontinued, Starting S

Ibutilide Protocol: Notify Cardiology Service before and after administration of Ibutilide Routine, Until discontinued, Starting S

[X] Ibutilide Protocol: Notify central telemetry monitoring before and after administration of Ibutilide	Routine, Until discontinued, Starting S
[X] Ibutilide protocol: Apply electrode Zoll pads for NSVT. If patient develops unstable ventricular arrhythmias, then shock per ACLS protocol.	Routine, Until discontinued, Starting S
[X] Ibutilide Protocol: Continuous ECG monitoring during and at least 4 hours following infusion or until QTc has returned to baseline. Print out ECG strips for QTc monitoring every 15 minutes for 2 hours post infusion.	Routine, Until discontinued, Starting S

IV Fluids

VTE

Labs

Labs (STAT)

[X] Magnesium	STAT For 1 Occurrences Draw baseline if not already completed within 24 hours
[X] Potassium	STAT For 1 Occurrences Draw baseline if not already completed within 24 hours

Cardiology

ECG

[X] ECG 12 lead	Routine, Once For 1 Occurrences Clinical Indications: Cardiac Arrhythmia Interpreting Physician: Baseline ECG
[X] ECG 12 lead	Routine, As needed For 1 Occurrences Clinical Indications: Cardiac Arrhythmia Interpreting Physician: Obtain 4 hours post ibutilide infusion

Imaging

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Additional Orders