

## General

### Common Present on Admission Diagnosis

<input type="checkbox"/> Acidosis	Details
<input type="checkbox"/> Acute Post-Hemorrhagic Anemia	Details
<input type="checkbox"/> Acute Renal Failure	Details
<input type="checkbox"/> Acute Respiratory Failure	Details
<input type="checkbox"/> Acute Thromboembolism of Deep Veins of Lower Extremities	Details
<input type="checkbox"/> Anemia	Details
<input type="checkbox"/> Bacteremia	Details
<input type="checkbox"/> Bipolar disorder, unspecified	Details
<input type="checkbox"/> Cardiac Arrest	Details
<input type="checkbox"/> Cardiac Dysrhythmia	Details
<input type="checkbox"/> Cardiogenic Shock	Details
<input type="checkbox"/> Decubitus Ulcer	Details
<input type="checkbox"/> Dementia in Conditions Classified Elsewhere	Details
<input type="checkbox"/> Disorder of Liver	Details
<input type="checkbox"/> Electrolyte and Fluid Disorder	Details
<input type="checkbox"/> Intestinal Infection due to Clostridium Difficile	Details
<input type="checkbox"/> Methicillin Resistant Staphylococcus Aureus Infection	Details
<input type="checkbox"/> Obstructive Chronic Bronchitis with Exacerbation	Details
<input type="checkbox"/> Other Alteration of Consciousness	Details
<input type="checkbox"/> Other and Unspecified Coagulation Defects	Details
<input type="checkbox"/> Other Pulmonary Embolism and Infarction	Details
<input type="checkbox"/> Phlebitis and Thrombophlebitis	Details
<input type="checkbox"/> Protein-calorie Malnutrition	Details
<input type="checkbox"/> Psychosis, unspecified psychosis type	Details
<input type="checkbox"/> Schizophrenia Disorder	Details
<input type="checkbox"/> Sepsis	Details
<input type="checkbox"/> Septic Shock	Details
<input type="checkbox"/> Septicemia	Details
<input type="checkbox"/> Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Details
<input type="checkbox"/> Urinary Tract Infection, Site Not Specified	Details

### Admission or Observation (Single Response) (Selection Required)

<input type="radio"/> Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
<input type="radio"/> Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
<input type="radio"/> Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

### Transfer

<input type="checkbox"/> Transfer patient	Level of Care: Bed request comments:
---	---

## Code Status

<input type="checkbox"/> Full code	Code Status decision reached by:
<input type="checkbox"/> DNR (Selection Required)	
<input type="checkbox"/> DNR (Do Not Resuscitate)	Does patient have decision-making capacity?
<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult:
<input type="checkbox"/> Modified Code	Does patient have decision-making capacity? Modified Code restrictions:
<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions:

## Isolation

<input type="checkbox"/> Airborne isolation status	
<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics.	Once, Sputum
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

## Precautions

<input type="checkbox"/> Aspiration precautions	Details
<input type="checkbox"/> Fall precautions	Increased observation level needed:
<input type="checkbox"/> Latex precautions	Details
<input type="checkbox"/> Seizure precautions	Increased observation level needed:

## Nursing

### Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 15 min Every 15 minutes x 4, then every 30 minutes x 4, then every 1 hour x 4, then every shift.
---	--

### Activity

<input type="checkbox"/> Bedrest	Routine, Until discontinued, Starting S With *** leg straight after sheath removal for *** hours.
<input type="checkbox"/> Head of bed 30 degrees	Routine, Until discontinued, Starting S Head of bed: 30 degrees No greater than a maximum of 30 degrees for *** hours after sheath removal.
<input type="checkbox"/> Activity as tolerated	Routine, Until discontinued, Starting S Specify: Activity as tolerated

### Nursing

<input type="checkbox"/> National Institute of Health Stroke Score - on admission (done by certified personnel)	Routine, Once For 1 Occurrences
<input type="checkbox"/> Neurological assessment	Routine, Every 15 min Assessment to Perform: Every 15 min times 4, then every 30 min times 2, then every 1 hour times 4, then every 4 hours.
<input type="checkbox"/> Assess cath site	Routine, Every 15 min Every 15 minutes times 4, then every 30 minutes times 4, then every 1 hour times 4, then every shift.

[ ] Pulse checks	Routine, Every 15 min Pulses to assess: Side: Every 15 minutes times 4, then every 30 minutes times 4, then every 1 hour times 4, then every shift.
[ ] Perform ACT	Routine, Until discontinued, Starting S Check ACT 2 hours post procedure.
[ ] Remove sheath	Routine, Once If ACT less than 170 seconds
[ ] Post procedure site assessment	Routine, Every 30 min Procedure Site: sheath site Observe for bleeding/hematoma at sheath site every 30 min times 2, then every 1 hour times 8 hours after sheath removal.
[ ] Foley catheter care	Routine, Until discontinued, Starting S Orders: to gravity
[ ] Apply condom catheter	Routine, Once
[ ] Foley catheter - discontinue	Routine, Once When patient is ambulatory
[ ] Tobacco cessation education	Routine, Once If applicable
[ ] Hold all outpatient antihypertensive therapy	Routine, Until discontinued, Starting S For 24 Hours For 24 hours post procedure

#### Diet

[ ] NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options:
[ ] NPO - except meds	Diet effective now, Starting S NPO: Except meds Pre-Operative fasting options:
[ ] Diet Clear Liquids	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Yes Target Diet: Advance target diet criteria: Liquid Consistency: Fluid Restriction: Foods to Avoid:
[ ] Diet Low Fat, Low Cholesterol	Diet effective now, Starting S Diet(s): Low Fat, Low Cholesterol Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
[ ] Diet 1800 Carb Control Diabetic	Diet effective now, Starting S Diet(s): Other Diabetic/Cal Diabetic/Calorie: 1800 Kcal/202 gm Carbohydrate Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
[ ] Diet Heart Healthy, No added salt	Diet effective now, Starting S Diet(s): Heart Healthy Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: No added salt

<input type="checkbox"/> Diet	Diet effective now, Starting S Diet(s): Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
-------------------------------	--

## Notify

<input type="checkbox"/> Notify Consulting Neurologist for changes in neurologic status	Routine, Until discontinued, Starting S
<input type="checkbox"/> Notify Attending physician for sustained heart rate (BPM) less than 55	Routine, Until discontinued, Starting S
<input type="checkbox"/> Notify Attending physician for sustained systolic blood pressure (mmHg) less than 100	Routine, Until discontinued, Starting S
<input type="checkbox"/> Notify Attending physician for absent or diminished pulse(s) in affected extremity	Routine, Until discontinued, Starting S

## IV Fluids

### IV Fluids (Single Response)

<input type="radio"/> sodium chloride 0.9 % infusion	intravenous, continuous
<input type="radio"/> sodium chloride 0.9 % with potassium chloride 20 mEq/L infusion	intravenous, continuous
<input type="radio"/> lactated Ringer's infusion	intravenous, continuous
<input type="radio"/> sodium chloride with femoral sheath	intravenous, continuous Give with femoral sheath(s)

## Medications

### Light Sedation - Sheath Removal

<input type="checkbox"/> morPHINE injection	intravenous, once PRN, severe pain (score 7-10), prior to sheath removal
<input type="checkbox"/> MIDAZolam (VERSED) injection	intravenous, once PRN, prior to sheath removal Indication(s):

### Scheduled Medications

<input type="checkbox"/> aspirin chewable tablet	81 mg, oral, daily
<input type="checkbox"/> aspirin tablet	325 mg, oral, daily
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	325 mg, oral, daily
<input type="checkbox"/> clopidogrel (PLAVIX) tablet	75 mg, oral, daily
<input type="checkbox"/> ticagrelor (BRILINTA) tablet	90 mg, oral, 2 times daily
<input type="checkbox"/> atorvastatin (LIPITOR) tablet	10 mg, oral, nightly
<input type="checkbox"/> simvastatin (ZOCOR) tablet	40 mg, oral, nightly
<input type="checkbox"/> ezetimibe (ZETIA) tablet	10 mg, oral, nightly

### Medications - IV Infusion

<input type="checkbox"/> DOPamine (INTROPIN) infusion	2-10 mcg/kg/min, intravenous, continuous May titrate to a maximum of 10 microgram/kg/min as needed for systolic blood pressure LESS than 100 mmHg. May titrate as tolerated.
<input type="checkbox"/> vasopressin (PITRESSIN) infusion	0.1-0.4 Units/min, intravenous, continuous May titrate to maximum 0.04 units/min as needed for systolic blood pressure less than 100 mmHg.
<input type="checkbox"/> norepinephrine (LEVOPHED) infusion	1-20 mcg/min, intravenous, continuous May titrate to a maximum of 20 microgram/min as needed for systolic blood pressure LESS than 100 mmHg.

### Medications PRN

<input type="checkbox"/> docusate sodium (COLACE) capsule	100 mg, oral, 2 times daily PRN, constipation
---	---

<input type="checkbox"/> atropine injection	1 mg, intravenous, every 1 hour prn, sustained heart rate LESS than 55 beats/minute. may repeat if necessary
<input type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	<b>"Or" Linked Panel</b>
<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting Give if patient is able to tolerate oral medication.
<input type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.

## Pain Medications

<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, every 6 hours PRN, mild pain (score 1-3)
<input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6)
<input type="checkbox"/> morPHINE injection	4 mg, intravenous, every 3 hours PRN, severe pain (score 7-10)

## VTE

### DVT Risk and Prophylaxis Tool (Single Response) (Selection Required)

URL: "appt1.pdf"

<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> LOW Risk of DVT (Selection Required)	

#### Low Risk Definition

Age less than 60 years and NO other VTE risk factors

<input type="checkbox"/> Low Risk (Single Response) (Selection Required)	
<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation

### ( ) MODERATE Risk of DVT - Surgical (Selection Required)

#### Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

<input type="checkbox"/> Moderate Risk (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device	<b>"And" Linked Panel</b>
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous

( ) Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis		"And" Linked Panel
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis		Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis		Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)		
<input type="checkbox"/> enoxaparin (LOVENOX) syringe		40 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min		30 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min		30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min		40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection		2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection		5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)		5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet		oral, daily at 1700 (TIME CRITICAL), Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)		STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)		
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis		Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous		Routine, Continuous
<input type="checkbox"/> MODERATE Risk of DVT - Non-Surgical (Selection Required)		
Moderate Risk Definition		
Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.		
One or more of the following medical conditions:		
CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome		
Age 60 and above		
Central line		
History of DVT or family history of VTE		
Anticipated length of stay GREATER than 48 hours		
Less than fully and independently ambulatory		
Estrogen therapy		
Moderate or major surgery (not for cancer)		
Major surgery within 3 months of admission		
<input type="checkbox"/> Moderate Risk (Selection Required)		
<input type="checkbox"/> Moderate risk of VTE		Routine, Once

[ ] Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	<b>"And" Linked Panel</b>
[ ] Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	<b>"And" Linked Panel</b>
[ ] Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
[ ] Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (TIME CRITICAL), Starting S
( ) patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (TIME CRITICAL), Starting S For Patients with CrCl LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response) (Selection Required)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) HIGH Risk of DVT - Surgical (Selection Required)	

## High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1
( ) patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1 For Patients with CrCl LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) HIGH Risk of DVT - Non-Surgical (Selection Required)	

## High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (TIME CRITICAL), Starting S
( ) patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (TIME CRITICAL), Starting S For Patients with CrCl LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	

## High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
( ) aspirin chewable tablet	162 mg, oral, daily, Starting S+1
( ) aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1
( ) enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1 For Patients with CrCl LESS than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL), Starting S+1 To be Given on Post Op Day 1. Indications:
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):

( ) Place/Maintain sequential compression device continuous      Routine, Continuous

**DVT Risk and Prophylaxis Tool (Single Response) (Selection Required)**

URL: "\appt1.pdf"

( ) Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis      Routine, Once  
No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.  
Therapy for the following:

**( ) LOW Risk of DVT (Selection Required)**

Low Risk Definition

Age less than 60 years and NO other VTE risk factors

**[ ] Low Risk (Single Response) (Selection Required)**

( ) Low risk of VTE      Routine, Once  
Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation

**( ) MODERATE Risk of DVT - Surgical (Selection Required)**

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

**[ ] Moderate Risk (Selection Required)**

[ ] Moderate risk of VTE      Routine, Once

**[ ] Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)**

( ) Contraindications exist for pharmacologic prophylaxis      **"And" Linked Panel**  
BUT order Sequential compression device

[ ] Contraindications exist for pharmacologic prophylaxis      Routine, Once  
No pharmacologic VTE prophylaxis due to the following contraindication(s):

[ ] Place/Maintain sequential compression device continuous      Routine, Continuous

( ) Contraindications exist for pharmacologic prophylaxis      **"And" Linked Panel**  
AND mechanical prophylaxis

[ ] Contraindications exist for pharmacologic prophylaxis      Routine, Once  
No pharmacologic VTE prophylaxis due to the following contraindication(s):

[ ] Contraindications exist for mechanical prophylaxis      Routine, Once  
No mechanical VTE prophylaxis due to the following contraindication(s):

( ) enoxaparin (LOVENOX) injection (Single Response)  
(Selection Required)

( ) enoxaparin (LOVENOX) syringe      40 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1

( ) patients with CrCL LESS than 30 mL/min      30 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1  
For Patients with CrCL LESS than 30 mL/min

( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input checked="" type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) MODERATE Risk of DVT - Non-Surgical (Selection Required)	
Address pharmacologic prophylaxis by selecting one of the following. Mechanical prophylaxis is optional unless pharmacologic prophylaxis is contraindicated.	
<input checked="" type="checkbox"/> Moderate Risk (Selection Required)	
<input checked="" type="checkbox"/> Moderate risk of VTE	Routine, Once
<input checked="" type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	<b>"And" Linked Panel</b>
<input checked="" type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input checked="" type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	<b>"And" Linked Panel</b>
<input checked="" type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input checked="" type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (TIME CRITICAL), Starting S+1
( ) patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (TIME CRITICAL), Starting S+1 For Patients with CrCL LESS than 30 mL/min

( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

( ) HIGH Risk of DVT - Surgical (Selection Required)

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1
( ) patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

( ) HIGH Risk of DVT - Non-Surgical (Selection Required)

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily, Starting S+1
( ) patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily, Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, every 12 hours at 0900, 2100 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, every 12 hours at 0900, 2100 (TIME CRITICAL) For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
( ) HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
( ) aspirin chewable tablet	162 mg, oral, daily, Starting S+1
( ) aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1
( ) enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1 For Patients with CrCL LESS than 30 mL/min.

( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL), Starting S+1 To be Given on Post Op Day 1. Indications:
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

#### DVT Risk and Prophylaxis Tool (Single Response)

URL: "\appt1.pdf"

( ) Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
( ) <u>LOW Risk of DVT (Selection Required)</u> Low Risk Definition Age less than 60 years and NO other VTE risk factors	

[ ] Low Risk (Single Response) (Selection Required)

( ) Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation
---------------------	---

( ) MODERATE Risk of DVT - Surgical (Selection Required)

Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission
---

[ ] Moderate Risk (Selection Required)

[ ] Moderate risk of VTE	Routine, Once
--------------------------	---------------

[ ] Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device	<b>"And" Linked Panel</b>
[ ] Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	<b>"And" Linked Panel</b>
[ ] Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
[ ] Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1
( ) patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1 For Patients with CrCl LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response) (Selection Required)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) MODERATE Risk of DVT - Non-Surgical (Selection Required)	

## Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

### Moderate Risk (Selection Required)

<input type="checkbox"/> Moderate risk of VTE	Routine, Once
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	<b>"And" Linked Panel</b>
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous
( ) Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	<b>"And" Linked Panel</b>
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (TIME CRITICAL), Starting S
( ) patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (TIME CRITICAL), Starting S For Patients with CrCl LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

[ ] Mechanical Prophylaxis (Single Response) (Selection Required)

( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous

( ) HIGH Risk of DVT - Surgical (Selection Required)

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

[ ] High Risk (Selection Required)

[ ] High risk of VTE	Routine, Once
----------------------	---------------

[ ] High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)

( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1
( ) patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1 For Patients with CrCl LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response) (Selection Required)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
( ) HIGH Risk of DVT - Non-Surgical (Selection Required)	

## High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700 (TIME CRITICAL), Starting S
( ) patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700 (TIME CRITICAL), Starting S For Patients with CrCl LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL) Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):
( ) Place/Maintain sequential compression device continuous	Routine, Continuous
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	

## High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
( ) aspirin chewable tablet	162 mg, oral, daily, Starting S+1
( ) aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1
( ) enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600 (TIME CRITICAL), Starting S+1 For Patients with CrCl LESS than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL), Starting S+1 To be Given on Post Op Day 1. Indications:
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 (TIME CRITICAL), Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s):

( ) Place/Maintain sequential compression device continuous	Routine, Continuous
---	---------------------

## Labs

### Labs

[ ] CBC with differential	Once, Starting S
[ ] Basic metabolic panel	Once, Starting S
[ ] Lipid panel	Once If not done on admission

## Cardiology

### Diagnostic Imaging

#### Diagnostics - CT

[ ] CT Head W Contrast	Routine, 1 time imaging For 1
[ ] CT Head Wo Contrast	Routine, 1 time imaging For 1

## Other Diagnostic Studies

## Respiratory

### Respiratory

[ ] Oxygen therapy	Routine, As needed Device 1: Nasal Cannula Rate in liters per minute: Rate in tenths of a liter per minute: O2 %: Titrate to keep O2 Sat Above: 92% Indications for O2 therapy:
[ ] Oxygen therapy	Routine, Continuous Device 1: Non-rebreather mask Rate in liters per minute: Titrate to keep O2 Sat Above: 92% Indications for O2 therapy:

## Rehab

## Consults

## Additional Orders

## Discharge

### Discharge Order (Single Response)

( ) Discharge patient when criteria met	Routine, Once Discharge Criteria: Scheduling/ADT
---	--

### Discontinue tubes/drains

[ ] Discontinue Foley catheter	Routine, Once, Scheduling/ADT
[ ] Discharge home with Foley catheter	Routine, Once, Scheduling/ADT
[ ] Discontinue IV	Routine, Once For 1 Occurrences, Scheduling/ADT
[ ] Deaccess port	
[ ] Deaccess Port-a-cath	Routine, Once, Scheduling/ADT
[ ] heparin, porcine (PF) 100 unit/mL injection	intra-catheter, once, Scheduling/ADT

### Discharge Activity

[ ] Activity as tolerated	Routine, Normal, Scheduling/ADT
[ ] Ambulate with assistance or assistive device	Routine, Normal, Scheduling/ADT

<input type="checkbox"/> Lifting restrictions	Routine, Normal, Scheduling/ADT, No lifting over 10 pounds.
<input type="checkbox"/> Weight bearing restrictions (specify)	Routine, Normal, Scheduling/ADT Weight Bearing Status: Extremity: ***
<input type="checkbox"/> Moderate bedrest with complete pelvic rest (no tampons, douching, sex)	Routine, Normal, Scheduling/ADT
<input type="checkbox"/> Complete pelvic rest (no tampons, douching, sex)	Routine, Normal, Scheduling/ADT
<input type="checkbox"/> No driving for 2 weeks	Routine, Normal, Scheduling/ADT
<input type="checkbox"/> Shower instructions:	Routine, Normal, Scheduling/ADT, ***
<input type="checkbox"/> Discharge activity	Routine, Normal, Scheduling/ADT
<input type="checkbox"/> Other restrictions (specify):	Routine, Normal, Scheduling/ADT, ***

#### Wound/Incision Care

<input type="checkbox"/> Discharge wound care	Routine, Normal, Scheduling/ADT, ***
<input type="checkbox"/> Discharge incision care	Routine, Normal, Scheduling/ADT, ***
<input type="checkbox"/> Discharge dressing	Routine, Normal, Scheduling/ADT, ***

#### Discharge Diet - REQUIRED (Single Response)

<input type="radio"/> Discharge Diet	Routine, Normal, Scheduling/ADT Discharge Diet:
<input type="radio"/> Discharge Diet- Regular	Routine, Normal, Scheduling/ADT Discharge Diet: Regular

#### Patient to notify physician

<input type="checkbox"/> Call physician for:	Routine, Normal, Scheduling/ADT, Temperature greater than 100.5
<input type="checkbox"/> Call physician for: Persistent nausea or vomiting	Routine, Normal, Scheduling/ADT
<input type="checkbox"/> Call physician for: severe uncontrolled pain	Routine, Normal, Scheduling/ADT
<input type="checkbox"/> Call physician for: redness, tenderness, or signs of infection (pain, swelling, redness, odor or green/yellow discharge from affected area)	Routine, Normal, Scheduling/ADT
<input type="checkbox"/> Call physician for difficulty breathing, chest pain, persistent dizziness or light-headedness	Routine, Normal, Scheduling/ADT
<input type="checkbox"/> Call physician for:	Routine, Normal, Scheduling/ADT, ***

#### Discharge Education

<input type="checkbox"/> Nurse to provide discharge education	Routine, Once Patient/Family: Both Education for: Other (specify) Specify: Nurse to provide patient education Scheduling/ADT
<input type="checkbox"/> Nurse to provide tobacco cessation education	Routine, Once Patient/Family: Both Education for: Other (specify) Specify: Nurse to provide tobacco cessation education Scheduling/ADT

#### Discharge Instructions

<input type="checkbox"/> Additional discharge instructions for Patient	Routine, Normal, Scheduling/ADT, ***
<input type="checkbox"/> Discharge instructions for Nursing- Will not show on AVS	Routine, Once ***, Scheduling/ADT

#### Place Follow-Up Order

<input type="checkbox"/> Follow-up with me	Follow up with me: Clinic Contact: Follow up in: On date: Appointment Time:
<input type="checkbox"/> Follow-up with primary care physician	Routine, Normal, Scheduling/ADT

[ ] Follow-up with physician	Follow up on: Appointment Time: Follow up in: Instructions for Follow Up:
[ ] Follow-up with physician	Follow up on: Appointment Time: Follow up in: Instructions for Follow Up:
[ ] Follow-up with department	Details