

General

Admission or Observation (Single Response) (Selection Required)

<input type="checkbox"/> Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
<input type="checkbox"/> Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
<input type="checkbox"/> Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

Nursing

Nursing

<input checked="" type="checkbox"/> Peripheral vascular assessment	Routine, Once DVT or Arterial-Catheter Directed Thrombolysis Admission, Post-op
<input checked="" type="checkbox"/> Intake and output	Routine, Every hour, Post-op
<input checked="" type="checkbox"/> Foley catheter care	Routine, Until discontinued, Starting S Orders: Maintain To bedside drainage, Post-op
<input checked="" type="checkbox"/> Daily weights	Routine, Daily, Post-op
<input checked="" type="checkbox"/> Nursing wound care	Routine, Every 12 hours Wound care to be performed by: Location: Site: Irrigate wound? Apply: Dressing Type: Reinforce dressing., Post-op
<input checked="" type="checkbox"/> Nursing wound care	Routine, Every 12 hours Wound care to be performed by: Location: Site: Irrigate wound? Apply: Dressing Type: Wrap affected leg in ACE bandage., Post-op
<input checked="" type="checkbox"/> Site care	Routine, Once Site: Place 10 pound sandbag over insertion site, if bleeding. (For groin insertions only), Post-op
<input checked="" type="checkbox"/> Please hold all Anticoagulants while therapy in progress.	Routine, Until discontinued, Starting S, Post-op
<input checked="" type="checkbox"/> Assess IV site	Routine, Every hour Assess catheter access site for bleeding/hematoma every 1 hour., Post-op
<input checked="" type="checkbox"/> No injections	Routine, Until discontinued, Starting S Type of injection: Post-op

Notify

<input checked="" type="checkbox"/> Notify Physician for fibrinogen less than 250	Routine, Until discontinued, Starting S, Post-op
<input checked="" type="checkbox"/> Notify Physician for all changes in TPA and for any bleeding	Routine, Until discontinued, Starting S, Post-op

Activity

<input checked="" type="checkbox"/> Strict bed rest	Routine, Until discontinued, Starting S, Post-op
---	--

Diet

<input checked="" type="checkbox"/> NPO (After Midnight)	Diet effective midnight, Starting S+1 at 12:01 AM NPO: Pre-Operative fasting options: Post-op
<input type="checkbox"/> Diet - Clear Liquids	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op

IV Fluids

IV Fluids

<input type="checkbox"/> sodium chloride 0.9 % bolus	500 mL, intravenous, once, Post-op For systolic BP less than 100 and/or increase in heart rate of 20 BPM or decrease in SBP of 20 mm Hg.
<input type="checkbox"/> sodium chloride 0.45 % infusion	100 mL/hr, intravenous, for 10 Hours, continuous, Post-op
<input checked="" type="checkbox"/> sodium chloride 0.9 % infusion	100 mL/hr, intravenous, for 10 Hours, continuous, Post-op
<input type="checkbox"/> dextrose 5%-0.45% sodium chloride infusion	100 mL/hr, intravenous, for 10 Hours, continuous, Post-op
<input type="checkbox"/> dextrose 5%-0.9% sodium chloride infusion	100 mL/hr, intravenous, for 10 Hours, continuous, Post-op

Medications

Medications - HMWB Only

<input type="checkbox"/> RIGHT (Single Response)	
<input type="checkbox"/> RIGHT - Alteplase 1 mg/hr + Heparin + Sodium Chloride	"And" Linked Panel
<input type="checkbox"/> RIGHT - alteplase (CATHFLO) 8 mg in sodium chloride 0.9 % 240 mL infusion	1 mg/hr, intra-catheter, continuous, Starting H Infuse via drug ports on EKOS x4 hours Fibrinogen >250 - no change in rate of tPa Fibrinogen 200-250 decrease tPa to half of the current rate and notify physician Fibrinogen <200 stop tPa and notify physician (**if tPa is stopped run NaCl through the line at 15 cc/hour for the first 4 hours), recheck fibrinogen after 4 hours
<input type="checkbox"/> RIGHT - HEParin (porcine) 10,000 Units in sodium chloride 0.9 % 1,000 mL infusion	300 Units/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/> sodium chloride 0.9 % infusion	35 mL/hr, intravenous, continuous, Starting H
<input type="checkbox"/> RIGHT - Alteplase 0.5 mg/hr + Heparin + Sodium Chloride	"And" Linked Panel
<input type="checkbox"/> RIGHT - alteplase (CATHFLO) 8 mg in sodium chloride 0.9 % 240 mL infusion	0.5 mg/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/> RIGHT - HEParin (porcine) 10,000 Units in sodium chloride 0.9 % 1,000 mL infusion	300 Units/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/> sodium chloride 0.9 % infusion	35 mL/hr, intravenous, continuous, Starting H
<input type="checkbox"/> LEFT (Single Response)	
<input type="checkbox"/> LEFT - Alteplase 1 mg/hr + Heparin + Sodium Chloride	"And" Linked Panel
<input type="checkbox"/> LEFT - alteplase (CATHFLO) 8 mg in sodium chloride 0.9 % 240 mL infusion	1 mg/hr, intra-catheter, continuous, Starting H Infuse via drug ports on EKOS x4 hours Fibrinogen >250 - no change in rate of tPa Fibrinogen 200-250 decrease tPa to half of the current rate and notify physician Fibrinogen <200 stop tPa and notify physician (**if tPa is stopped run NaCl through the line at 15 cc/hour for the first 4 hours), recheck fibrinogen after 4 hours

<input type="checkbox"/>	LEFT - HEParin (porcine) 10,000 Units in sodium chloride 0.9 % 1,000 mL infusion	300 Units/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/>	sodium chloride 0.9 % infusion	35 mL/hr, intravenous, continuous, Starting H
<input type="checkbox"/>	LEFT - Alteplase 0.5 mg/hr + Heparin + Sodium Chloride	"And" Linked Panel
<input type="checkbox"/>	LEFT - alteplase (CATHFLO) 8 mg in sodium chloride 0.9 % 240 mL infusion	0.5 mg/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/>	LEFT - HEParin (porcine) 10,000 Units in sodium chloride 0.9 % 1,000 mL infusion	300 Units/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/>	sodium chloride 0.9 % infusion	35 mL/hr, intravenous, continuous, Starting H

Medications - HMCL, HMSL, and HMTW Only

<input type="checkbox"/>	RIGHT (Single Response)	
<input type="checkbox"/>	RIGHT - Alteplase 1 mg/hr + Heparin + Sodium Chloride	"And" Linked Panel
<input type="checkbox"/>	alteplase (CATHFLO) 8 mg in sodium chloride 0.9 % 240 mL infusion	1 mg/hr, intra-catheter, continuous, Starting H Infuse via drug ports on EKOS x4 hours Fibrinogen >250 - no change in rate of tPa Fibrinogen 200-250 decrease tPa to half of the current rate and notify physician Fibrinogen <200 stop tPa and notify physician (**if tPa is stopped run NaCl through the line at 15 cc/hour for the first 4 hours), recheck fibrinogen after 4 hours
<input type="checkbox"/>	HEParin (porcine) 10,000 Units in sodium chloride 0.9 % 1,000 mL infusion	300 Units/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/>	sodium chloride 0.9 % infusion	35 mL/hr, intravenous, continuous, Starting H
<input type="checkbox"/>	RIGHT - Alteplase 0.5 mg/hr + Heparin + Sodium Chloride	"And" Linked Panel
<input type="checkbox"/>	alteplase (CATHFLO) 8 mg in sodium chloride 0.9 % 240 mL infusion	0.5 mg/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/>	HEParin (porcine) 10,000 Units in sodium chloride 0.9 % 1,000 mL infusion	300 Units/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/>	sodium chloride 0.9 % infusion	35 mL/hr, intravenous, continuous, Starting H
<input type="checkbox"/>	LEFT (Single Response)	
<input type="checkbox"/>	LEFT - Alteplase 1 mg/hr + Heparin + Sodium Chloride	"And" Linked Panel
<input type="checkbox"/>	alteplase (CATHFLO) 8 mg in sodium chloride 0.9 % 240 mL infusion	1 mg/hr, intra-catheter, continuous, Starting H Infuse via drug ports on EKOS x4 hours Fibrinogen >250 - no change in rate of tPa Fibrinogen 200-250 decrease tPa to half of the current rate and notify physician Fibrinogen <200 stop tPa and notify physician (**if tPa is stopped run NaCl through the line at 15 cc/hour for the first 4 hours), recheck fibrinogen after 4 hours
<input type="checkbox"/>	HEParin (porcine) 10,000 Units in sodium chloride 0.9 % 1,000 mL infusion	300 Units/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/>	sodium chloride 0.9 % infusion	35 mL/hr, intravenous, continuous, Starting H
<input type="checkbox"/>	LEFT - Alteplase 0.5 mg/hr + Heparin + Sodium Chloride	"And" Linked Panel
<input type="checkbox"/>	alteplase (CATHFLO) 8 mg in sodium chloride 0.9 % 240 mL infusion	0.5 mg/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/>	HEParin (porcine) 10,000 Units in sodium chloride 0.9 % 1,000 mL infusion	300 Units/hr, intra-catheter, continuous, Starting H
<input type="checkbox"/>	sodium chloride 0.9 % infusion	35 mL/hr, intravenous, continuous, Starting H

VTE

Labs

<input type="checkbox"/>	Basic metabolic panel	Daily While TPA therapy in progress, Post-op
<input type="checkbox"/>	CBC with platelet and differential	Every 8 hours While TPA therapy in progress, Post-op
<input type="checkbox"/>	Fibrinogen	Every 4 hours While TPA therapy in progress, Post-op

Cardiology

Imaging

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Additional Orders