

**General**

**Code Status**

- Full code Code Status decision reached by: Patient by means of Oral Directive

---

- DNR (Do Not Resuscitate) (Selection Required)
  - DNR (Do Not Resuscitate) Does patient have decision-making capacity?  
L&D Pre-Delivery
  - Consult to Palliative Care Service Priority:  
Reason for Consult?  
Order?  
Name of referring provider:  
Enter call back number:
  - Consult to Social Work Reason for Consult:  
L&D Pre-Delivery

---

- Modified Code Does patient have decision-making capacity?  
Modified Code restrictions:  
L&D Pre-Delivery

---

- Treatment Restrictions Treatment Restriction decision reached by:  
Specify Treatment Restrictions:  
L&D Pre-Delivery

**Isolation**

- Airborne isolation status
  - Airborne isolation status Details
  - Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics. Once, Sputum, L&D Pre-Delivery

---

- Contact isolation status Details

---

- Droplet isolation status Details

---

- Enteric isolation status Details

**Precautions**

- Aspiration precautions L&D Pre-Delivery

---

- Fall precautions Increased observation level needed:  
On Admission and every 8 hours, L&D Pre-Delivery

---

- Latex precautions L&D Pre-Delivery

---

- Seizure precautions Increased observation level needed:  
L&D Pre-Delivery

**Common Present on Admission Diagnosis**

- Acidosis L&D Pre-Delivery

---

- Acute Post-Hemorrhagic Anemia L&D Pre-Delivery

---

- Acute Renal Failure L&D Pre-Delivery

---

- Acute Respiratory Failure L&D Pre-Delivery

---

- Acute Thromboembolism of Deep Veins of Lower Extremities L&D Pre-Delivery

---

- Anemia L&D Pre-Delivery

---

- Bacteremia L&D Pre-Delivery

---

- Bipolar disorder, unspecified L&D Pre-Delivery

---

- Cardiac Arrest L&D Pre-Delivery

---

- Cardiac Dysrhythmia L&D Pre-Delivery

---

- Cardiogenic Shock L&D Pre-Delivery

---

- Decubitus Ulcer L&D Pre-Delivery

---

- Dementia in Conditions Classified Elsewhere L&D Pre-Delivery

---

- Disorder of Liver L&D Pre-Delivery

---

- Electrolyte and Fluid Disorder L&D Pre-Delivery

<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	L&D Pre-Delivery
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	L&D Pre-Delivery
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	L&D Pre-Delivery
<input type="checkbox"/>	Other Alteration of Consciousness	L&D Pre-Delivery
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	L&D Pre-Delivery
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	L&D Pre-Delivery
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	L&D Pre-Delivery
<input type="checkbox"/>	Protein-calorie Malnutrition	L&D Pre-Delivery
<input type="checkbox"/>	Psychosis, unspecified psychosis type	L&D Pre-Delivery
<input type="checkbox"/>	Schizophrenia Disorder	L&D Pre-Delivery
<input type="checkbox"/>	Sepsis	L&D Pre-Delivery
<input type="checkbox"/>	Septic Shock	L&D Pre-Delivery
<input type="checkbox"/>	Septicemia	L&D Pre-Delivery
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	L&D Pre-Delivery
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	L&D Pre-Delivery
<input type="checkbox"/>	Present on Admission-History of preterm premature rupture of membranes	L&D Pre-Delivery

## Nursing

### Activity

<input type="checkbox"/>	Strict bed rest	Routine, Until discontinued, Starting S Flat, L&D Pre-Delivery
<input type="checkbox"/>	Bed rest with bathroom privileges	Routine, Until discontinued, Starting S Bathroom Privileges: with bathroom privileges L&D Pre-Delivery
<input type="checkbox"/>	Ambulate with assistance	Routine, 3 times daily Specify: with assistance L&D Pre-Delivery

### Vital Signs

<input checked="" type="checkbox"/>	Vital signs - T/P/R/BP - Per unit protocol	Routine, Per unit protocol, L&D Pre-Delivery
<input type="checkbox"/>	Pulse oximetry	Routine, Continuous Current FIO2 or Room Air: L&D Pre-Delivery
<input type="checkbox"/>	Pulse oximetry	Routine, Once For 1 Occurrences Current FIO2 or Room Air: L&D Pre-Delivery
<input type="checkbox"/>	Pain Assessment	Routine, Per unit protocol Assess: Pain L&D Pre-Delivery

### Nursing Care

<input type="checkbox"/>	Apply external fetal monitor	
<input type="checkbox"/>	Monitor fetal heart tones continuous	Routine, Continuous Type: Continuous L&D Pre-Delivery
<input type="checkbox"/>	Monitor fetal heart tones intermittent	Routine, Continuous Type: Intermittent L&D Pre-Delivery
<input type="checkbox"/>	Fetal nonstress test	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/>	Tocometry	Routine, Until discontinued, Starting S Type: Continuous L&D Pre-Delivery
<input type="checkbox"/>	Apply internal fetal monitor (FSE)	
<input type="checkbox"/>	Fetal scalp monitor	Routine, Continuous Indication: May place Fetal Scalp Electrode if unable to adequately monitor fetal heart tones and membranes are ruptured. May place if GBS positive: Yes L&D Pre-Delivery

<input type="checkbox"/>	POC nitrazine	Once Rule out ruptured membranes, L&D Pre-Delivery
<input type="checkbox"/>	Sterile vaginal exam	Routine, Once Perform sterile vaginal exam to monitor progression or if clinically indicated, L&D Pre-Delivery
<input type="checkbox"/>	No sterile vaginal exam	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/>	Sterile speculum exam	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/>	Encourage fluids	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/>	Discharge instructions for Nursing	Routine, Once Send patient home with precautions, L&D Pre-Delivery

### Nursing Care

<input type="checkbox"/>	Apply external fetal monitor	
<input type="checkbox"/>	Monitor fetal heart tones continuous	Routine, Continuous Type: Continuous L&D Pre-Delivery
<input type="checkbox"/>	Monitor fetal heart tones intermittent	Routine, Continuous Type: Intermittent L&D Pre-Delivery
<input type="checkbox"/>	Fetal nonstress test	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/>	Tocometry	Routine, Until discontinued, Starting S Type: Continuous L&D Pre-Delivery
<input type="checkbox"/>	Apply internal fetal monitor (FSE)	
<input type="checkbox"/>	Fetal scalp monitor	Routine, Continuous Indication: May place Fetal Scalp Electrode if unable to adequately monitor fetal heart tones and membranes are ruptured. May place if GBS positive: Yes L&D Pre-Delivery
<input type="checkbox"/>	POC AmnioTest	Once Rule out ruptured membranes, L&D Pre-Delivery
<input type="checkbox"/>	Sterile vaginal exam	Routine, Once Perform sterile vaginal exam to monitor progression or if clinically indicated, L&D Pre-Delivery
<input type="checkbox"/>	No sterile vaginal exam	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/>	Sterile speculum exam	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/>	Encourage fluids	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/>	Discharge instructions for Nursing	Routine, Once Send patient home with precautions, L&D Pre-Delivery

### Nursing Care

<input type="checkbox"/>	Apply external fetal monitor	
<input type="checkbox"/>	Monitor fetal heart tones continuous	Routine, Continuous Type: Continuous L&D Pre-Delivery
<input type="checkbox"/>	Monitor fetal heart tones intermittent	Routine, Continuous Type: Intermittent L&D Pre-Delivery
<input type="checkbox"/>	Fetal nonstress test	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/>	Tocometry	Routine, Until discontinued, Starting S Type: Continuous L&D Pre-Delivery
<input type="checkbox"/>	Apply internal fetal monitor (FSE)	
<input type="checkbox"/>	Fetal scalp monitor	Routine, Continuous Indication: May place Fetal Scalp Electrode if unable to adequately monitor fetal heart tones and membranes are ruptured. May place if GBS positive: Yes L&D Pre-Delivery
<input type="checkbox"/>	POC Amnisure	Once Rule out ruptured membranes, L&D Pre-Delivery

<input type="checkbox"/> Sterile vaginal exam	Routine, Once Perform sterile vaginal exam to monitor progression or if clinically indicated, L&D Pre-Delivery
<input type="checkbox"/> No sterile vaginal exam	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/> Sterile speculum exam	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Encourage fluids	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/> Discharge instructions for Nursing	Routine, Once Send patient home with precautions, L&D Pre-Delivery

### Nursing Care

<input type="checkbox"/> Apply external fetal monitor	
<input type="checkbox"/> Monitor fetal heart tones continuous	Routine, Continuous Type: Continuous L&D Pre-Delivery
<input type="checkbox"/> Monitor fetal heart tones intermittent	Routine, Continuous Type: Intermittent L&D Pre-Delivery
<input type="checkbox"/> Fetal nonstress test	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Tocometry	Routine, Until discontinued, Starting S Type: Continuous L&D Pre-Delivery
<input type="checkbox"/> Apply internal fetal monitor (FSE)	
<input type="checkbox"/> Fetal scalp monitor	Routine, Continuous Indication: May place Fetal Scalp Electrode if unable to adequately monitor fetal heart tones and membranes are ruptured. May place if GBS positive: Yes L&D Pre-Delivery
<input type="checkbox"/> Amnisure	Once, Amniotic fluid Rule out ruptured membranes, L&D Pre-Delivery
<input type="checkbox"/> Sterile vaginal exam	Routine, Once Perform sterile vaginal exam to monitor progression or if clinically indicated, L&D Pre-Delivery
<input type="checkbox"/> No sterile vaginal exam	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/> Sterile speculum exam	Routine, Once, L&D Pre-Delivery
<input type="checkbox"/> Encourage fluids	Routine, Until discontinued, Starting S, L&D Pre-Delivery
<input type="checkbox"/> Discharge instructions for Nursing	Routine, Once Send patient home with precautions, L&D Pre-Delivery

### Notify

<input checked="" type="checkbox"/> Notify Physician when initial assessment is complete OR within one hour.	Routine, Until discontinued, Starting S, L&D Pre-Delivery
--	---

### Diet

<input checked="" type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options: L&D Pre-Delivery
<input type="checkbox"/> Diet - Clear Liquid	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: L&D Pre-Delivery
<input type="checkbox"/> Diet - Regular	Diet effective now, Starting S Diet(s): Regular Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: L&D Pre-Delivery

<input type="checkbox"/> Diet -	Diet effective now, Starting S Diet(s): Other Options: Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: L&D Pre-Delivery
---------------------------------	--

## IV Fluids

### IV Fluids

<input type="checkbox"/> lactated ringers infusion And bolus	<b>"And" Linked Panel</b>
<input type="checkbox"/> lactated Ringer's infusion	125 mL/hr, intravenous, continuous, L&D Pre-Delivery
<input type="checkbox"/> lactated Ringer's bolus from bag	1,000 mL, intravenous, at 999 mL/hr, PRN, Bolus as needed for Non-reassuring FHR with Tachysystole, L&D Pre-Delivery

### Peripheral IV Access

<input type="checkbox"/> Initiate and maintain IV	
<input type="checkbox"/> Insert peripheral IV	Routine, Once
<input type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 12 hours scheduled
<input type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care

## Medications

### Tocolytics

<input type="checkbox"/> NIFEdipine (PROCARDIA) capsule	10 mg, oral, once, For 1 Doses, L&D Pre-Delivery HOLD parameters for this order: Contact Physician if:
<input type="checkbox"/> terbutaline (BRETHINE) injection	0.25 mg, subcutaneous, every 20 min PRN, For Tocolysis, For 3 Doses, L&D Pre-Delivery May give up to 3 doses as needed. HOLD for pulse GREATER than 120 BPM.

### Mild Pain (Pain Score 1-3) (Single Response)

<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, once PRN, mild pain (score 1-3), L&D Pre-Delivery
---	---

### Moderate Pain (Pain Score 4-6) (Single Response)

<input type="checkbox"/> butorphanol (STADOL) injection	1 mg, intravenous, once PRN, moderate pain (score 4-6), L&D Pre-Delivery
<input type="checkbox"/> nalbuphine (NUBAIN) injection	5 mg, intravenous, once PRN, moderate pain (score 4-6), L&D Pre-Delivery

### Moderate Pain (Pain Score 4-6) (Single Response)

<input type="checkbox"/> butorphanol (STADOL) injection	1 mg, intravenous, once PRN, moderate pain (score 4-6), L&D Pre-Delivery
<input type="checkbox"/> nalbuphine (NUBAIN) injection	5 mg, intravenous, once PRN, moderate pain (score 4-6), L&D Pre-Delivery

### Severe Pain (Pain Score 7-10) (Single Response)

<input type="checkbox"/> butorphanol (STADOL) injection	2 mg, intravenous, once PRN, severe pain (score 7-10), L&D Pre-Delivery
<input type="checkbox"/> nalbuphine (NUBAIN) injection	10 mg, intravenous, once PRN, severe pain (score 7-10), L&D Pre-Delivery

### PRN Antihypertensives

<input type="checkbox"/> For blood pressure GREATER than or EQUAL to 160 mmHg - labetalol (NORMODYNE) tablet	200 mg, oral, 2 times daily PRN, high blood pressure, For Systolic BP GREATER than 160mmHg and Diastolic BP GREATER than 110mmHg., L&D Pre-Delivery DO NOT give incremental dosing as this will compromise fetal circulation. HOLD parameters for this order: Hold Parameters requested HOLD for: HOLD for: Contact Physician if: For Systolic BP GREATER than 160mmHG and Diastolic BP GREATER than 110mmHg
<input type="checkbox"/> hydrALAZINE (APRESOLINE) tablet	5 mg, oral, every 6 hours PRN, high blood pressure, For blood pressure GREATER than 160mmHG DO NOT give incremental dosing as this will compromise fetal circulation. HOLD parameters for this order: Contact Physician if:

**PRN Severe Hypertension**

<input type="checkbox"/> NIFEdipine (PROCARDIA) capsule	10 mg, oral, once PRN, high blood pressure, for severe BP elevations of 15 min or more. Recheck BP in 15 min., For 1 Doses, L&D Pre-Delivery HOLD parameters for this order: Contact Physician if: For Systolic BP GREATER than 160mmHG and Diastolic BP GREATER than 110mmHg
<input type="checkbox"/> labetalol (NORMODYNE, TRANDATE) injection	20 mg, intravenous, once PRN, high blood pressure, for severe blood pressure elevation (systolic BP GREATER than or EQUAL to 110 mm Hg) persisting for 15 minutes or more., For 1 Doses, L&D Pre-Delivery Give IV Push over 2 minutes. Repeat BP measurements in 10 minutes and record results. HOLD parameters for this order: Contact Physician if: For Systolic BP GREATER than 160mmHG and Diastolic BP GREATER than 110mmHg

**Antiemetics - HHM, HMSJ, HMW, HMSTC, HMTW Only**

<input type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required) <b>"Or" Linked Panel</b>	
<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is able to tolerate oral medication.
<input type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal <b>"Or" Linked Panel</b>	
<input type="checkbox"/> promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

**Antiemetics - HMSL, HMWB Only**

<input type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required) <b>"Or" Linked Panel</b>	
<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is able to tolerate oral medication.
<input type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.

<input type="checkbox"/>	promethazine (PHENERGAN) IV or Oral or Rectal	<b>"Or" Linked Panel</b>
<input type="checkbox"/>	promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9 % 0.9 % 20 mL for Alaris pump syringe option	12.5 mg, intravenous, at 60 mL/hr, for 20 Minutes, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/>	promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/>	promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

### Antiemetics - HMSTJ Only

<input type="checkbox"/>	ondansetron (ZOFTRAN) IV or Oral (Selection Required)	<b>"Or" Linked Panel</b>
<input type="checkbox"/>	ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is able to tolerate oral medication.
<input type="checkbox"/>	ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/>	promethazine (PHENERGAN) IVPB or Oral or Rectal	<b>"Or" Linked Panel</b>
<input type="checkbox"/>	promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, intravenous, for 30 Minutes, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/>	promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.
<input type="checkbox"/>	promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, L&D Pre-Delivery Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

## Labs

### COVID-19 Qualitative PCR

<input type="checkbox"/>	COVID-19 qualitative PCR - Nasopharyngeal swab	STAT For 1 Occurrences Specimen Source: Nasopharyngeal Swab Is this for pre-procedure or non-PUI assessment? Yes L&D Pre-Delivery
--------------------------	--	--

### Hematology and Coagulation - NOT HMCL

<input type="checkbox"/>	CBC hemogram	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Fibrinogen	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Kleihauer-Betke	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Prothrombin time with INR	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Partial thromboplastin time	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Type and screen, obstetrical patient	STAT For 1 Occurrences, L&D Pre-Delivery

### Hematology and Coagulation - HMCL ONLY

<input type="checkbox"/>	CBC hemogram	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Fibrinogen	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Kleihauer-Betke	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Prothrombin time with INR	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Partial thromboplastin time	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Type and screen	STAT For 1 Occurrences, L&D Pre-Delivery

### Chemistry

<input type="checkbox"/>	Basic metabolic panel	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	Comprehensive metabolic panel	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/>	D-dimer	Once, L&D Pre-Delivery

<input type="checkbox"/> Magnesium	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Uric acid	STAT For 1 Occurrences, L&D Pre-Delivery

### Chemistry HMWB

<input type="checkbox"/> Basic metabolic panel	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Comprehensive metabolic panel	Once, L&D Pre-Delivery
<input type="checkbox"/> D-dimer	Once, L&D Pre-Delivery
<input type="checkbox"/> Magnesium	STAT For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Uric acid	STAT For 1 Occurrences, L&D Pre-Delivery

### OB Screening Markers

<input type="checkbox"/> Fetal fibronectin	Once, L&D Pre-Delivery
<input type="checkbox"/> POC Amnisure	Once, L&D Pre-Delivery

### OB Screening Markers

<input type="checkbox"/> POC AmnioTest	Once Rule out ruptured Membranes, L&D Pre-Delivery
<input type="checkbox"/> Amnisure	Once, Amniotic fluid, L&D Pre-Delivery
<input type="checkbox"/> Fern	Once, Vaginal fluid, L&D Pre-Delivery
<input type="checkbox"/> Fetal fibronectin	Once, L&D Pre-Delivery
<input type="checkbox"/> POC nitrazine	Once, L&D Pre-Delivery

### OB Screening Markers

<input type="checkbox"/> POC nitrazine	Once, L&D Pre-Delivery
<input type="checkbox"/> POC AmnioTest	Once Rule out ruptured Membranes, L&D Pre-Delivery
<input type="checkbox"/> Fetal fibronectin	Once, L&D Pre-Delivery
<input type="checkbox"/> POC Amnisure	Once, L&D Pre-Delivery

### Pre-Eclamptic Lab Panel

<input type="checkbox"/> Pre-Eclamptic Lab Panel	
<input type="checkbox"/> CBC with differential	Once, L&D Pre-Delivery
<input type="checkbox"/> Comprehensive metabolic panel	Once, L&D Pre-Delivery
<input type="checkbox"/> Prothrombin time with INR	Once, L&D Pre-Delivery
<input type="checkbox"/> Partial thromboplastin time	Once, L&D Pre-Delivery
<input type="checkbox"/> Fibrinogen	Once, L&D Pre-Delivery
<input type="checkbox"/> Uric acid	Once, L&D Pre-Delivery
<input type="checkbox"/> LDH	Once, L&D Pre-Delivery
<input type="checkbox"/> Urine Protein and Creatinine	
<input type="checkbox"/> Creatinine level, urine, random	Once For 1 Occurrences, L&D Pre-Delivery
<input type="checkbox"/> Protein, urine, random	Once For 1 Occurrences, L&D Pre-Delivery

### Magnesium and D-dimer

<input type="checkbox"/> D-dimer	Once, L&D Pre-Delivery
<input type="checkbox"/> Magnesium level	Once, L&D Pre-Delivery

### 24 Hour urine

<input type="checkbox"/> 24 Hour urine	
<input type="checkbox"/> Creatinine clearance, urine, 24 hour	Once, L&D Pre-Delivery
<input type="checkbox"/> Protein, urine, 24 hour	Once, L&D Pre-Delivery

### Microbiology

<input type="checkbox"/> STI Screen	
<input type="checkbox"/> Sexually Transmitted Infections	



<input type="checkbox"/> Chlamydia trachomatis by PCR	Once Urine is the only acceptable source for patients less than or equal to 13 years old. Specimen source: Urine Specimen Source: Specimen Site: L&D Pre-Delivery
<input type="checkbox"/> Neisseria gonorrhoeae by PCR	Once Urine is the only acceptable source for patients less than or equal to 13 years old. Specimen source: Urine Specimen Source: Specimen Site: L&D Pre-Delivery
<input type="checkbox"/> Beta Strep Screen Culture with Lim Broth	Once, Vaginal, L&D Pre-Delivery
<input type="checkbox"/> Urinalysis screen and microscopy, with reflex to culture	Once Specimen Source: Urine Specimen Site: L&D Pre-Delivery
<input type="checkbox"/> Wet prep	Once, L&D Pre-Delivery
<input type="checkbox"/> POC wet mount, vaginal	Once, L&D Pre-Delivery
<input type="checkbox"/> Fern	Once, Vaginal fluid, L&D Pre-Delivery

#### Urine Dipstick HWMB

<input type="checkbox"/> Urinalysis, dipstick only	Once, L&D Pre-Delivery
--	------------------------

### Other Studies

#### Ultrasound

<input type="checkbox"/> US Fetal Biophysical Profile	STAT, 1 time imaging For 1 , L&D Pre-Delivery
<input type="checkbox"/> US Pregnancy Transvaginal	STAT, 1 time imaging For 1 , L&D Pre-Delivery
<input type="checkbox"/> US Pregnancy Single Less Than 14 Weeks	STAT, 1 time imaging For 1 , L&D Pre-Delivery
<input type="checkbox"/> US Pregnancy Greater Than 14 Weeks	STAT, 1 time imaging For 1 , L&D Pre-Delivery
<input type="checkbox"/> US Pregnancy Limited	STAT, 1 time imaging For 1 , L&D Pre-Delivery

### Consults

For Physician Consult orders use sidebar

#### Ancillary consults

<input type="checkbox"/> Consult to PT eval and treat	Reasons for referral to Physical Therapy (mark all applicable): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation( if values are very abnormal): Weight Bearing Status: L&D Pre-Delivery
<input type="checkbox"/> Consult to Social Work	Reason for Consult: L&D Pre-Delivery
<input type="checkbox"/> Consult to Spiritual Care	Reason for consult? L&D Pre-Delivery