

Outpatient Gynecologic Surgery PostOp [1441]

General

Isolation

<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Airborne isolation status	Once, Sputum, Post-op
<input type="checkbox"/> Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics.	
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op

Nursing

Vital Signs

<input checked="" type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol, Post-op
--	-------------------------------------

Activity

<input checked="" type="checkbox"/> Ambulate with assistance	Routine, 3 times daily Specify: with assistance Post-op
--	---

Diet

<input checked="" type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options: NPO until no longer nauseated., Post-op
<input type="checkbox"/> Diet -	Diet effective now, Starting S Diet(s): Regular Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid:
<input type="checkbox"/> Diet clear liquid	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Yes Target Diet: Advance target diet criteria: Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op
<input type="checkbox"/> Diet - Regular	Diet effective now, Starting S Diet(s): Regular Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op

Nursing Care

<input type="checkbox"/> Document Pad Count	Routine, Once For 1 Occurrences Prior to Discharge, Post-op
<input type="checkbox"/> Foley catheter - discontinue	Routine, Once, Post-op
<input type="checkbox"/> Post-op voiding trial	Routine, Once, Post-op
<input type="checkbox"/> Straight cath	Routine, Conditional Frequency For 1 Occurrences Post bladder scan and bedpan, Post-op

Discharge

<input checked="" type="checkbox"/> Discharge instructions	Routine, Until discontinued, Starting S Evaluate for discharge if patient is voiding, ambulating, and has no nausea/vomiting. May discharge with copy of printed post operative discharge instructions and copies of prescriptions, if indicated., Post-op
<input type="checkbox"/> Discharge Patient	Routine, Until discontinued, Starting S Discharge patient at ***. Confirm that medication reconciliation is complete prior to discharge., Post-op

IV Fluids

IV Fluids

<input type="checkbox"/> dextrose 5 % and lactated Ringer's infusion	125 mL/hr, intravenous, continuous, Post-op
<input type="checkbox"/> dextrose 5%-0.9% sodium chloride infusion	125 mL/hr, intravenous, continuous, Post-op

Peripheral IV Access

<input checked="" type="checkbox"/> Initiate and maintain IV	
<input checked="" type="checkbox"/> Insert peripheral IV	Routine, Once
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 12 hours scheduled
<input checked="" type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care

Medications

Moderate Pain (Pain Score 4-6) (Single Response)

<input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	2 tablet, oral, every 3 hours PRN, moderate pain (score 4-6), Post-op
<input type="checkbox"/> ketorolac (TORADOL) tablet - Not recommended in patients with eGFR LESS than 30 mL/min OR in acute kidney injury	10 mg, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Not recommended in patients with eGFR LESS than 30 mL/min OR in acute kidney injury

Severe Pain (Pain Score 7-10)

<input type="checkbox"/> morPHINE injection	intravenous, every 3 hours PRN, severe pain (score 7-10), and Notify MD, Post-op
[] ketorolac (TORADOL) IV (Single Response)	

Do NOT use in patients with eGFR LESS than 30 mL/min AND/OR patients LESS than 17 years of age.
WARNING: Use is contraindicated for treatment of perioperative pain OR in the setting of coronary artery bypass graft (CABG) surgery.

<input type="checkbox"/> For patients ages GREATER than 64 OR weight LESS than 50 kg OR eGFR 30-59 mL/min - ketorolac (TORADOL) injection	15 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6)
<input type="checkbox"/> For patients ages 17-64 AND weight GREATER than or EQUAL to 50 kg AND eGFR at least 60 mL/min - ketorolac (TORADOL) injection	30 mg, intravenous, every 6 hours PRN, moderate pain (score 4-6)

Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	"Or" Linked Panel
<input checked="" type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.

[X] ondansetron (ZOFRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
[X] promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics - HMSL, HMWB Only

[X] ondansetron (ZOFRAN) IV or Oral (Selection Required)	"Or" Linked Panel
[X] ondansetron ODT (ZOFRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
[X] ondansetron (ZOFRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IV or Oral or Rectal	"Or" Linked Panel
[X] promethazine (PHENERGAN) 12.5 mg in sodium chloride 0.9 % 0.9 % 20 mL for Alaris pump syringe option	12.5 mg, intravenous, at 60 mL/hr, for 20 Minutes, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics - HMSTJ Only

[X] ondansetron (ZOFRAN) IV or Oral (Selection Required)	"Or" Linked Panel
[X] ondansetron ODT (ZOFRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
[X] ondansetron (ZOFRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) IVPB or Oral or Rectal	"Or" Linked Panel
[X] promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB	12.5 mg, intravenous, for 30 Minutes, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
[X] promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFRAN) is ineffective and patient is able to tolerate oral medication.
[X] promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFRAN) is ineffective and patient is UNable to tolerate oral medication.

Rh Negative Mother

Nursing

[X] Rhogam Workup: If Mother is Rh Negative, complete Rhogam workup and administer Rh immune globulin 50 mcg (or dose determined by lab antibody results) IM within 72 hours of delivery.	Routine, Until discontinued, Starting S, Post-op
---	--

Labs

<input type="checkbox"/> Fetal Screen	Conditional Frequency For 1 Occurrences Conditional- One activation- If Rh Negative Mom and Rh Positive infant, Post-op
<input type="checkbox"/> Rhogam Type and Screen	Once, Post-op
Medication	
<input type="checkbox"/> rho(D) immune globulin (HYPERRHO/RHOGAM) injection	300 mcg, intramuscular, PRN, Rhogam Workup: If Mother is Rh Negative, complete Rhogam workup and administer Rh immune globulin 50 mcg (or dose determined by lab antibody results) IM within 72 hours of delivery., For 1 Doses, Post-op