

Newborn Circumcision [1447]

Nursing

Consent

<input checked="" type="checkbox"/> Complete consent for Circumcision	Routine, Once Procedure: Circumcision Diagnosis/Condition: Physician: Ensure circumcision consent signed by parent/guardian
---	---

PreProcedure

<input checked="" type="checkbox"/> Please confirm: initial physical exam by physician must be documented in medical record prior to circumcision	Routine, Once For 1 Occurrences
---	---------------------------------

PostProcedure

<input checked="" type="checkbox"/> Circumcision care	Routine, Every 15 min For Until specified Check site for bleeding every 15 minutes x4, then until stable
<input type="checkbox"/> Vaseline gauze to bedside	Routine, Until discontinued, Starting S Apply to circumcision site following circumcision with Mogen clamp or Gomco; follow with Vaseline applied to gauze with each diaper change
<input checked="" type="checkbox"/> Infant must be observed for 2 hours post circumcision prior to discharge	Routine, Until discontinued, Starting S
<input checked="" type="checkbox"/> Patient education: Post Circumcision Care	Routine, Once Patient/Family: Family Education for: Other (specify) Specify: post-circumcision care
<input checked="" type="checkbox"/> Notify Physician immediately if excessive bleeding noted	Routine, Until discontinued, Starting S, immediately if excessive bleeding noted

Medications

Medications - HHM, HMWB, HMTW Only

<input checked="" type="checkbox"/> povidone-iodine (BETADINE) 10 % external solution	1 application, Topical, once, For 1 Doses Apply to penis pre-procedure for site cleaning
<input type="checkbox"/> lidocaine-prilocaine (EMLA) cream	1 application, Topical, once, For 1 Doses Apply to Penis Pre-Procedure
<input checked="" type="checkbox"/> lidocaine PF (XYLOCAINE) 10 mg/mL (1%) injection	1 mL, infiltration, once, For 1 Doses
<input checked="" type="checkbox"/> sucrose 24 % oral solution	0.1 mL, oral, every 5 min PRN, mild pain (score 1-3), pain control during procedure, For 3 Doses
<input type="checkbox"/> microfibrillar collagen (AVITENE) powder	Topical, PRN, bleeding
<input type="checkbox"/> white petrolatum ointment	1 application, Topical, PRN, Diaper Changes Apply to Penis with Diaper Changes
<input type="checkbox"/> silver nitrate applicators	1 application, Topical, PRN

Medications - HMB Only

<input checked="" type="checkbox"/> povidone-iodine (BETADINE) 10 % external solution	1 application, Topical, once, For 1 Doses Apply to penis pre-procedure for site cleaning
<input checked="" type="checkbox"/> lidocaine-prilocaine (EMLA) cream	1 application, Topical, PRN, mild pain (score 1-3)
<input checked="" type="checkbox"/> lidocaine PF (XYLOCAINE) 10 mg/mL (1%) injection	1 mL, infiltration, once, For 1 Doses
<input checked="" type="checkbox"/> sucrose 24 % oral solution	0.1 mL, oral, every 5 min PRN, mild pain (score 1-3), pain control during procedure, For 3 Doses
<input type="checkbox"/> microfibrillar collagen (AVITENE) powder	Topical, PRN, bleeding
<input type="checkbox"/> white petrolatum ointment	1 application, Topical, PRN, Diaper Changes Apply to Penis with Diaper Changes
<input type="checkbox"/> silver nitrate applicators	1 application, Topical, PRN

Medications - HMSTJ Only

<input type="checkbox"/> lidocaine-prilocaine (EMLA) cream	1 application, Topical, once, For 1 Doses Apply to Penis Pre-Procedure
<input checked="" type="checkbox"/> lidocaine PF (XYLOCAINE) 10 mg/mL (1%) injection	1 mL, infiltration, once, For 1 Doses
<input checked="" type="checkbox"/> sucrose 24 % oral solution	0.1 mL, oral, every 5 min PRN, mild pain (score 1-3), pain control during procedure, For 3 Doses
<input type="checkbox"/> microfibrillar collagen (AVITENE) powder	Topical, PRN, bleeding
<input type="checkbox"/> white petrolatum ointment	1 application, Topical, PRN, Diaper Changes Apply to Penis with Diaper Changes
<input type="checkbox"/> silver nitrate applicators	1 application, Topical, PRN

Medications - HMSL Only

<input checked="" type="checkbox"/> povidone-iodine (BETADINE) 10 % external solution	1 application, Topical, once, For 1 Doses Apply to penis pre-procedure for site cleaning
<input type="checkbox"/> lidocaine-prilocaine (EMLA) cream	1 application, Topical, once, For 1 Doses Apply to Penis Pre-Procedure
<input checked="" type="checkbox"/> lidocaine PF (XYLOCAINE) 10 mg/mL (1%) injection	1 mL, infiltration, once, For 1 Doses
<input checked="" type="checkbox"/> sucrose 24 % oral solution	0.1 mL, oral, every 5 min PRN, mild pain (score 1-3), pain control during procedure, For 3 Doses
<input type="checkbox"/> microfibrillar collagen (HEMOSTAT, AVITENE) pad	1 application, Topical, PRN, bleeding
<input type="checkbox"/> white petrolatum ointment	1 application, Topical, PRN, Diaper Changes Apply to Penis with Diaper Changes
<input type="checkbox"/> silver nitrate applicators	1 application, Topical, PRN

Medications - HMW Only

<input checked="" type="checkbox"/> povidone-iodine (BETADINE) 10 % external solution	1 application, Topical, once, For 1 Doses Apply to penis pre-procedure for site cleaning
<input type="checkbox"/> lidocaine-prilocaine (EMLA) cream	1 application, Topical, once, For 1 Doses Apply to Penis Pre-Procedure
<input checked="" type="checkbox"/> lidocaine PF (XYLOCAINE) 10 mg/mL (1%) injection	1 mL, infiltration, once, For 1 Doses
<input checked="" type="checkbox"/> sucrose 24 % oral solution	0.1 mL, oral, every 5 min PRN, mild pain (score 1-3), pain control during procedure, For 3 Doses
<input type="checkbox"/> microfibrillar collagen (HEMOSTAT, AVITENE) pad	1 application, Topical, PRN, bleeding
<input type="checkbox"/> white petrolatum ointment	1 application, Topical, PRN, Diaper Changes Apply to Penis with Diaper Changes
<input type="checkbox"/> silver nitrate applicators	1 application, Topical, PRN