

Newborn Hyperbilirubinemia [1457]

Nursing

Nursing

<input checked="" type="checkbox"/> Phototherapy	Routine, Continuous Type of Phototherapy: Cover infant's eyes during phototherapy
<input checked="" type="checkbox"/> Measure and maintain irradiance level greater than 30 microwatts/squared centimeters/nanometers	Routine, Until discontinued, Starting S
<input checked="" type="checkbox"/> Measure and maintain a distance of light 12-18 inches from baby	Routine, Until discontinued, Starting S
<input checked="" type="checkbox"/> Plot the total serum bilirubin on the nomogram for phototherapy and exchange transfusions provided by the guidelines for babies greater than 35 weeks gestation, OR enter total serum bilirubin and baby's hours of age on BILITOOL to determine risk and inform physician about the levels	Routine, Until discontinued, Starting S

Notify

<input checked="" type="checkbox"/> Notify Physician	Routine, Until discontinued, Starting S, -Immediately, if bilirubin level plots at High risk on Nomogram or Bilitool -During morning rounds or no later than 9am, if bilirubin level plots at High Interim risk on Nomogram or Bilitool
--	--

Labs

Laboratory

<input checked="" type="checkbox"/> Bilirubin, neonatal	Daily For 3 Occurrences
<input type="checkbox"/> Total bilirubin	Once
<input type="checkbox"/> Hemoglobin and hematocrit	Once
<input type="checkbox"/> CBC with differential	Once
<input type="checkbox"/> Reticulocyte count	Once