

General**Notify Physician (Single Response) (Selection Required)**

Notify provider if amIODarone therapy held for heart rate less than 60 or pauses greater than 2.5 seconds Routine, Until discontinued, Starting S, Notify provider if amIODarone therapy held for heart rate less than 60 or pauses greater than 2.5 seconds

Nursing**Vital Signs**

| [X] Telemetry | "And" Linked Panel |
|--|--|
| [X] Telemetry monitoring | Routine, Continuous Order: Place in Centralized Telemetry Monitor: EKG Monitoring Only (Telemetry Box) Reason for telemetry: Can be off of Telemetry for tests and baths? Yes |
| [X] Telemetry Additional Setup Information | Routine, Continuous High Heart Rate (BPM): 120 Low Heart Rate(BPM): 50 High PVC's (per minute): 10 High SBP(mmHg): 175 Low SBP(mmHg): 100 High DBP(mmHg): 95 Low DBP(mmHg): 40 Low Mean BP: 60 High Mean BP: 120 Low SPO2(%): 94 |

IV Fluids**Infusions - Select for CENTRAL or Peripheral Line Administration (Single Response) (Selection Required)**

() CENTRAL Line Administration (Single Response)

() Loading Dose and Maintenance Infusion (Single Response)

Select Standard or Double concentration

() Standard

[] CENTRAL Line Administration: amIODarone (CORDArone) 150 mg LOADING Dose followed by STANDARD concentration 24-hour Infusion for Atrial Fibrillation- NOT HMWB

"Followed by" Linked Panel

[] amIODarone (CORDArone) 150 mg BOLUS

150 mg, intravenous, once, Starting S, For 1 Doses
Patients should be monitored for QTc prolongation.

[] amIODarone 1.8 mg/mL (STANDARD concentration) infusion

1 mg/min, intravenous, continuous, Starting H+10 Minutes
Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours.
HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours.

[] REDUCE rate for amIODarone (CORDArone) 450 mg/ 250 mL NS

0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses
HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. Do not take down 1st infusion until entire content of bag is infused.

[] amIODarone 1.8 mg/mL (STANDARD concentration) infusion - 2nd bag

0.5 mg/min, intravenous, continuous, Starting H+8 Hours
HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours.

| | |
|--|---|
| <input type="checkbox"/> CENTRAL Line Administration: amiODarone (CORDArone) 150 mg LOADING Dose followed by STANDARD concentration 24-hour Infusion for Atrial Fibrillation-HMWB ONLY | "Followed by" Linked Panel |
| <input type="checkbox"/> amiODarone (CORDArone) 150 mg BOLUS | 150 mg, intravenous, once, Starting S, For 1 Doses Patients should be monitored for QTc prolongation. |
| <input type="checkbox"/> amiODarone 1.8 mg/mL (STANDARD concentration) infusion | 1 mg/min, intravenous, continuous, Starting H+10 Minutes Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. |
| <input type="checkbox"/> REDUCE rate for amiODarone (CORDArone) infusion | 0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. Do not take down 1st infusion until entire content of bag is infused. |
| <input type="checkbox"/> amiODarone 1.8 mg/mL (STANDARD concentration) infusion - 2nd bag | 0.5 mg/min, intravenous, continuous, Starting H+8 Hours HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours. |
| () Double | "Followed by" Linked Panel |
| <input type="checkbox"/> CENTRAL Line Administration: amiODarone (CORDArone) 150 mg LOADING Dose followed by DOUBLE concentration 24-hour Infusion for Atrial Fibrillation | 150 mg, intravenous, once, Starting S, For 1 Doses Patients should be monitored for QTc prolongation. |
| <input type="checkbox"/> amiODarone (CORDArone) 150 mg BOLUS | 1 mg/min, intravenous, continuous, Starting H+10 Minutes Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line. |
| <input type="checkbox"/> amiODarone (CORDArone) 900 mg/ 250 mL NS | 0.5 mg/min, intravenous, continuous, Starting H+6 Hours Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line. Do not take down 1st infusion until entire content of bag is infused. |
| () Maintenance Infusion (Single Response) | |
| Select Standard or Double Concentration | |
| () Standard | "Followed by" Linked Panel |
| <input type="checkbox"/> NO LOADING DOSE - Central Line Administration: amiODarone (CORDArone) STANDARD concentration 24-hour Infusion for Atrial Fibrillation - NOT HMWB | 1 mg/min, intravenous, continuous Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line. |
| <input type="checkbox"/> amiODarone 1.8 mg/mL (STANDARD concentration) infusion | 0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. Do not take down 1st infusion until entire content of bag is infused. |
| <input type="checkbox"/> REDUCE rate for amiODarone (CORDArone) 450 mg/ 250 mL NS | 0.5 mg/min, intravenous, continuous, Starting H+8 Hours HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. |
| <input type="checkbox"/> amiODarone 1.8 mg/mL (STANDARD concentration) infusion - 2nd bag | 0.5 mg/min, intravenous, continuous, Starting H+10 Minutes Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line. |

| | |
|---|---|
| <input type="checkbox"/> NO LOADING DOSE - Central Line Administration: amIODarone (CORDArone) STANDARD concentration 24-hour Infusion for Atrial Fibrillation - HMWB Only | "Followed by" Linked Panel |
| <input type="checkbox"/> amIODarone 1.8 mg/mL (STANDARD concentration) infusion | 1 mg/min, intravenous, continuous Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. |
| <input type="checkbox"/> REDUCE rate for amIODarone (CORDArone) 360 mg/ 200 mL NS | 0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. Do not take down 1st infusion until entire content of bag is infused. |
| <input type="checkbox"/> amIODarone 1.8 mg/mL (STANDARD concentration) infusion - 2nd bag | 0.5 mg/min, intravenous, continuous, Starting H+8 Hours HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. |
| () Double (Single Response) | |
| <input type="checkbox"/> NO LOADING DOSE - Central Line Administration: amIODarone (CORDArone) Double Concentration 24-hour Infusion for Atrial Fibrillation | "Followed by" Linked Panel |
| <input type="checkbox"/> amIODarone (CORDArone) 900 mg/ 250 mL NS | 1 mg/min, intravenous, continuous Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line. |
| <input type="checkbox"/> REDUCE rate for amIODarone (CORDArone) 900 mg/ 250 mL NS | 0.5 mg/min, intravenous, continuous, Starting H+6 Hours Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line. Do not take down 1st infusion until entire content of bag is infused. |
| () Peripheral Line Administration (Single Response) | |
| () Loading Dose and Maintenance Infusion | |
| <input type="checkbox"/> PERIPHERAL Line Administration: amIODarone (CORDArone) 150 mg LOADING Dose followed by 24-hour Infusion for Atrial Fibrillation -NOT HMWB (Selection Required) | "Followed by" Linked Panel |
| <input type="checkbox"/> amIODarone (CORDArone) 150 mg LOADING Dose | 150 mg, intravenous, once, Starting S, For 1 Doses Patients should be monitored for QTc prolongation. |
| <input type="checkbox"/> amIODarone (CORDArone) 450 mg/ 250 mL infusion - 1st bag | 1 mg/min, intravenous, continuous, Starting H+10 Minutes Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. |
| <input type="checkbox"/> REDUCE rate for amIODarone (CORDArone) 450 mg/ 250 mL infusion | 0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. Do not take down 1st infusion until entire content of bag is infused. |
| <input type="checkbox"/> amIODarone (CORDArone) 450 mg/ 250 mL infusion - 2nd bag | 0.5 mg/min, intravenous, continuous, Starting H+8 Hours HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours. |
| <input type="checkbox"/> PERIPHERAL Line Administration: amIODarone (CORDArone) 150 mg LOADING Dose followed by 24-hour Infusion for Atrial Fibrillation -HMWB ONLY (Selection Required) | "Followed by" Linked Panel |

| | |
|--|---|
| [] amiODarone (CORDArone) 150 mg LOADING Dose | 150 mg, intravenous, once, Starting S, For 1 Doses Patients should be monitored for QTc prolongation. |
| [] amiODarone (CORDArone) 360 mg/ 200 mL infusion - 1st bag | 1 mg/min, intravenous, continuous, Starting H+10 Minutes Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. |
| [] REDUCE rate for amiODarone (CORDArone) infusion | 0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. Do not take down 1st infusion until entire content of bag is infused. |
| [] amiODarone (CORDArone) 360 mg/ 200 mL infusion - 2nd bag | 0.5 mg/min, intravenous, continuous, Starting H+8 Hours HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours. |
| () Maintenance Infusion | |
| [] NO LOADING DOSE - Peripheral Line Administration: amiODarone (CORDArone) STANDARD concentration 24-hour Infusion for Atrial Fibrillation | "Followed by" Linked Panel |
| [] amiODarone (CORDArone) 450 mg/ 250 mL infusion - 1st bag | 1 mg/min, intravenous, continuous, Starting H Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. |
| [] REDUCE rate for amiODarone (CORDArone) 450 mg/ 250 mL infusion | 0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours. Do not take down 1st infusion until entire content of bag is infused. |
| [] amiODarone (CORDArone) infusion solution -2nd bag | 0.5 mg/min, intravenous, continuous, Starting H+8 Hours HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours |
| [] NO LOADING DOSE - Peripheral Line Administration: amiODarone (CORDArone) STANDARD concentration 24-hour Infusion for Atrial Fibrillation - HMWB Only | "Followed by" Linked Panel |
| [] amiODarone (CORDArone) 360 mg/ 200 mL infusion - 1st bag | 1 mg/min, intravenous, continuous, Starting H Start with 1 mg/min for 6 hours. Decrease to 0.5 mg/min for 18 hours. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a CENTRAL line or PICC line if infusion duration is GREATER than 24 hours. |
| [] REDUCE rate for amiODarone (CORDArone) infusion | 0.5 mg/min, intravenous, once, Starting H+6 Hours, For 1 Doses Patients should be monitored for QTc prolongation. Use in-line filter to help prevent Phlebitis. HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours. Do not take down 1st infusion until entire content of bag is infused. |
| [] amiODarone (CORDArone) infusion solution -2nd bag | 0.5 mg/min, intravenous, continuous, Starting H+8 Hours HOLD Infusion for heart rate LESS THAN 60 or pauses GREATER THAN 2.5 seconds. MUST be infused via a central line or PICC line if infusion duration is GREATER THAN 24 hours |

amiODarone (PACERONE) tablet (Selection Required)

You MUST be sure the oral tablet order is set to start TOMORROW with the start time set to 24 hours AFTER the start time of the INITIAL infusion order above.

[X] amIODarone (PACERONE) tablet **** You MUST CHANGE the START DATE to TOMORROW and set the Start TIME to be 24 hours after the Start Time of the Infusion

200 mg, oral, every 24 hours scheduled, Starting H+24 Hours amiodarone (Pacerone) tablets must start 24 hours after the start of the infusion order.

Labs

Labs (Baseline checks required if not already completed)

[X] Hepatic function panel

Once

Draw for baseline if not already completed within 24 hours

[X] TSH

Once

Draw for baseline if not already completed within 24 hours

Cardiology

ECG

[X] ECG 12 lead

STAT, Once For 1 Occurrences

Clinical Indications: Cardiac Arrhythmia

Interpreting Physician:

Baseline ECG if not already completed

Imaging

X-ray

[X] Chest 2 Vw

Routine, 1 time imaging For 1

Other Studies

Respiratory

Consults

For Physician Consult orders use sidebar

Additional Orders