

## General

### Common Present on Admission Diagnosis

[ ] Acidosis	Details
[ ] Acute Post-Hemorrhagic Anemia	Details
[ ] Acute Renal Failure	Details
[ ] Acute Respiratory Failure	Details
[ ] Acute Thromboembolism of Deep Veins of Lower Extremities	Details
[ ] Anemia	Details
[ ] Bacteremia	Details
[ ] Bipolar disorder, unspecified	Details
[ ] Cardiac Arrest	Details
[ ] Cardiac Dysrhythmia	Details
[ ] Cardiogenic Shock	Details
[ ] Decubitus Ulcer	Details
[ ] Dementia in Conditions Classified Elsewhere	Details
[ ] Disorder of Liver	Details
[ ] Electrolyte and Fluid Disorder	Details
[ ] Intestinal Infection due to Clostridium Difficile	Details
[ ] Methicillin Resistant Staphylococcus Aureus Infection	Details
[ ] Obstructive Chronic Bronchitis with Exacerbation	Details
[ ] Other Alteration of Consciousness	Details
[ ] Other and Unspecified Coagulation Defects	Details
[ ] Other Pulmonary Embolism and Infarction	Details
[ ] Phlebitis and Thrombophlebitis	Details
[ ] Protein-calorie Malnutrition	Details
[ ] Psychosis, unspecified psychosis type	Details
[ ] Schizophrenia Disorder	Details
[ ] Sepsis	Details
[ ] Septic Shock	Details
[ ] Septicemia	Details
[ ] Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Details
[ ] Urinary Tract Infection, Site Not Specified	Details

### Admission or Observation (Single Response) (Selection Required)

( ) Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
( ) Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
( ) Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

### Admission or Observation (Single Response)

Patient has active status order on file

( ) Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
( ) Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments:
( ) Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments:

#### Admission (Single Response)

Patient has active status order on file.

( ) Admit to inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
------------------------	--

#### Code Status

[ ] Full code	Code Status decision reached by: Post-op
[ ] DNR (Do Not Resuscitate) (Selection Required)	
[ ] DNR (Do Not Resuscitate)	Does patient have decision-making capacity? Post-op
[ ] Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
[ ] Consult to Social Work	Reason for Consult: Post-op
[ ] Modified Code	Does patient have decision-making capacity? Modified Code restrictions: Post-op
[ ] Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

#### Isolation

[ ] Airborne isolation status	Details
[ ] Airborne isolation status	Once, Sputum, Post-op
[ ] Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics.	
[ ] Contact isolation status	Details
[ ] Droplet isolation status	Details
[ ] Enteric isolation status	Details

#### Precautions

<input type="checkbox"/> Aspiration precautions	PACU & Post-op
<input checked="" type="checkbox"/> Fall precautions	Increased observation level needed: PACU & Post-op
<input type="checkbox"/> Latex precautions	PACU & Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: PACU & Post-op
<input type="checkbox"/> Spinal precautions	PACU & Post-op

## Nursing

### Vital Signs (Single Response)

(X) Vital signs - T/P/R/BP	Routine, Per unit protocol, PACU & Post-op
----------------------------	--

### Activity

<input type="checkbox"/> Strict bed rest	Routine, Until discontinued, Starting S, PACU & Post-op
<input type="checkbox"/> Up with assistance	Routine, Until discontinued, Starting S Specify: Up with assistance PACU & Post-op
<input type="checkbox"/> Activity as tolerated	Routine, Until discontinued, Starting S Specify: Activity as tolerated PACU & Post-op
<input type="checkbox"/> All meals out of bed	Routine, Until discontinued, Starting S All meals out of bed, PACU & Post-op
<input type="checkbox"/> Head of bed 30 degrees	Routine, Until discontinued, Starting S Head of bed: 30 degrees PACU & Post-op
<input type="checkbox"/> Head of bed flat	Routine, Until discontinued, Starting S Head of bed: flat PACU & Post-op

### Nursing

<input checked="" type="checkbox"/> Assess operative site	Routine, Every 8 hours For bleeding, reinforce dressing, notify physician., PACU & Post-op
<input checked="" type="checkbox"/> Assess for pain	Routine, Every 4 hours Assess: for Pain PACU & Post-op
<input type="checkbox"/> Neurological assessment	Routine, Once Assessment to Perform: PACU & Post-op
<input type="checkbox"/> Straight cath	Routine, Every 6 hours If unable to void after second attempt, insert Foley and call physician., PACU & Post-op
<input type="checkbox"/> <u>Insert/Maintain Foley and Notify</u>	
<input type="checkbox"/> Insert Foley catheter	Routine, Once Type: Size: Urinometer needed: PACU & Post-op
<input type="checkbox"/> Foley catheter care	Routine, Until discontinued, Starting S Orders: Maintain PACU & Post-op
<input type="checkbox"/> Notify Physician if unable to void after second attempt at straight cath and Foley inserted	Routine, Until discontinued, Starting S, PACU & Post-op

[ ] Surgical/incision site care	Routine, Once Location: Site: Apply: Dressing Type: Open to air? PACU & Post-op
[ ] Reinforce dressing	Routine, As needed Reinforce with: If saturated., PACU & Post-op
[ ] Cervical collar - soft	Routine, Once Type of Collar to Apply: Soft cervical collar Special Instructions: obtain from central supply PACU & Post-op
[ ] Place antiembolic stockings	Routine, Once, PACU & Post-op
[ ] No anticoagulants INcluding UNfractionated heparin	Routine, Until discontinued, Starting S Reason for "No" order: PACU & Post-op
[ ] No anti-platelet agents INcluding aspirin	Routine, Until discontinued, Starting S Reason for "No" order: PACU & Post-op

#### Notify

[X] Notify Physician if acute change in neurological status	Routine, Until discontinued, Starting S, PACU & Post-op
[X] Notify Physician for itching	Routine, Until discontinued, Starting S, PACU & Post-op
[X] Notify Physician bleeding at site	Routine, Until discontinued, Starting S, PACU & Post-op
[X] Notify Physician of No Bowel Movement for more than 72 hours	Routine, Until discontinued, Starting S, PACU & Post-op

#### Diet

[ ] Diet - Clear liquids (advance as tolerated to Regular)	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Yes Target Diet: Regular Advance target diet criteria: Please assess bowel sounds between progressions. Liquid Consistency: Fluid Restriction: Foods to Avoid: PACU & Post-op
[ ] Diet - Full liquids	Diet effective now, Starting S Diet(s): Full Liquids Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: PACU & Post-op
[ ] Diet - Regular	Diet effective now, Starting S Diet(s): Regular Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: PACU & Post-op
[ ] Diet - Heart healthy	Diet effective now, Starting S Diet(s): Heart Healthy Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: PACU & Post-op

<input type="checkbox"/> Diet - 2000 Kcal/225 gm Carb	Diet effective now, Starting S Diet(s): 2000 Kcal/225 gm Carbohydrate Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: PACU & Post-op
<input type="checkbox"/> Diet	Diet effective now, Starting S Diet(s): Other Options: Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: PACU & Post-op

## IV Fluids

### IV Fluids (Single Response)

<input type="checkbox"/> lactated Ringer's infusion	intravenous, continuous, Post-op
<input type="checkbox"/> sodium chloride 0.9 % infusion	intravenous, continuous, Post-op
<input type="checkbox"/> sodium chloride 0.9 % with potassium chloride 20 mEq/L infusion	intravenous, continuous, Post-op
<input type="checkbox"/> dextrose 5 % and sodium chloride 0.45 % with potassium chloride 20 mEq/L infusion - for NPO Patients	intravenous, continuous, Post-op

## Medications

### Muscle Relaxants (Single Response)

<input type="checkbox"/> methocarbamol (ROBAXIN) 500 mg in sodium chloride 0.9 % 100 mL IVPB	500 mg, intravenous, for 60 Minutes, every 8 hours PRN, muscle spasms, Post-op
<input type="checkbox"/> methocarbamol (ROBAXIN) tablet	500 mg, oral, every 8 hours PRN, muscle spasms, Post-op
<input type="checkbox"/> cyclobenzaprine (FLEXERIL) tablet	5 mg, oral, every 8 hours PRN, muscle spasms, Post-op

### Muscle Relaxants - Refractory Treatments (Single Response)

<input type="checkbox"/> diazepam (VALIUM) injection	2.5 mg, intravenous, every 8 hours PRN, muscle spasms, inadequate muscle spasm relief following administration of other agents, Post-op Indication(s): Other Specify: Muscle Relaxant
<input type="checkbox"/> diazepam (VALIUM) tablet	2.5 mg, oral, every 8 hours PRN, muscle spasms, inadequate muscle spasm relief following administration of other agents, Post-op Indication(s): Other Specify: Muscle Relaxant

### Antiemetics

<input checked="" type="checkbox"/> ondansetron (ZOFTRAN) IV or Oral (Selection Required)	<b>"Or" Linked Panel</b>
<input type="checkbox"/> ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet	4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.
<input type="checkbox"/> ondansetron (ZOFTRAN) 4 mg/2 mL injection	4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) IV or Oral or Rectal	<b>"Or" Linked Panel</b>
<input type="checkbox"/> promethazine (PHENERGAN) 12.5 mg IV	12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.
<input type="checkbox"/> promethazine (PHENERGAN) tablet	12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.

<input type="checkbox"/> promethazine (PHENERGAN) suppository	12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.
<input type="checkbox"/> scopolamine (TRANSDERM-SCOP) 1.5 mg (1 mg over 3 days) - For Patients LESS than 65 years old	1 patch, transdermal, for 72 Hours, every 72 hours, Post-op

#### PRN Medications - Pain - Pain Score (1-3) (Single Response)

<input type="checkbox"/> traMADol (ULTRAM) tablet	25 mg, oral, every 4 hours PRN, mild pain (score 1-3), Post-op Maximum Daily Dose: 200 mg/day
---	--

#### PRN Medications - Pain - Pain Score (4-6) (Single Response)

<input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	1 tablet, oral, every 4 hours PRN, moderate pain (score 4-6), Post-op Do not exceed 3000 mg of acetaminophen daily from all sources.
<input type="checkbox"/> traMADol (ULTRAM) tablet	50 mg, oral, every 4 hours PRN, moderate pain (score 4-6), Post-op Maximum Daily Dose: 200 mg/day

#### PRN Medications - Pain - Pain Score (7-10) (Single Response)

<input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	2 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Do not exceed 3000 mg of acetaminophen daily from all sources.
--	--

#### Breakthrough Pain (Single Response)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	25 mcg, intravenous, every 2 hour PRN, other, pain (score: 4-10) - inadequate pain relief following administration of oral agents, Post-op Administer after pain re-assessment for inadequate pain relief.
<input type="checkbox"/> morphine 2 mg/mL injection	2 mg, intravenous, every 3 hours PRN, other, pain (score: 4-10) - inadequate pain relief following administration of oral agents, Post-op Administer after pain re-assessment for inadequate pain relief.
<input type="checkbox"/> HYDROmorphine (DILAUDID) injection	0.5 mg, intravenous, every 3 hours PRN, other, pain (score: 4-10) - inadequate pain relief following administration of oral agents, Post-op Administer after pain re-assessment for inadequate pain relief.

## VTE

#### DVT Risk and Prophylaxis Tool (Single Response) (Selection Required)

URL: "\appt1.pdf"

<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> LOW Risk of DVT (Selection Required)	

Low Risk Definition

Age less than 60 years and NO other VTE risk factors

#### Low Risk (Single Response) (Selection Required)

<input type="checkbox"/> Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation PACU & Post-op
--	---

#### MODERATE Risk of DVT - Surgical (Selection Required)



( ) warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
( ) MODERATE Risk of DVT - Non-Surgical (Selection Required)	
Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.	
One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome	
Age 60 and above	
Central line	
History of DVT or family history of VTE	
Anticipated length of stay GREATER than 48 hours	
Less than fully and independently ambulatory	
Estrogen therapy	
Moderate or major surgery (not for cancer)	
Major surgery within 3 months of admission	

[ ] Moderate Risk (Selection Required)	
[ ] Moderate risk of VTE	Routine, Once, PACU & Post-op
[ ] Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	<b>"And" Linked Panel</b>
[ ] Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
[ ] Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
( ) Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	<b>"And" Linked Panel</b>
[ ] Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
[ ] Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S
( ) patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCl LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700, PACU & Post-op Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

( ) HIGH Risk of DVT - Surgical (Selection Required)

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

[ ] High Risk (Selection Required)

[ ] High risk of VTE	Routine, Once, PACU & Post-op
[ ] High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
( ) patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response) (Selection Required)	

( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
( ) Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
( ) HIGH Risk of DVT - Non-Surgical (Selection Required)	
High Risk Definition	
Both pharmacologic AND mechanical prophylaxis must be addressed.	
One or more of the following medical conditions:	
Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)	
Severe fracture of hip, pelvis or leg	
Acute spinal cord injury with paresis	
Multiple major traumas	
Abdominal or pelvic surgery for CANCER	
Acute ischemic stroke	
History of PE	
[ ] High Risk (Selection Required)	
[ ] High risk of VTE	Routine, Once, PACU & Post-op
[ ] High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S
( ) patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCl LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[ ] Mechanical Prophylaxis (Single Response) (Selection Required)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
( ) Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
( ) HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	

## High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
( ) apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
( ) aspirin chewable tablet	162 mg, oral, daily, Starting S+1
( ) aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
( ) enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1
( ) enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCl LESS than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL), Starting S+1 To be Given on Post Op Day 1. Indications:
( ) warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	

( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
( ) Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

### DVT Risk and Prophylaxis Tool (Single Response)

URL: "\appt1.pdf"

( ) Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
--	--

#### ( ) LOW Risk of DVT (Selection Required)

Low Risk Definition

Age less than 60 years and NO other VTE risk factors

##### [ ] Low Risk (Single Response) (Selection Required)

( ) Low risk of VTE	Routine, Once Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation PACU & Post-op
---------------------	---

#### ( ) MODERATE Risk of DVT - Surgical (Selection Required)

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

##### [ ] Moderate Risk (Selection Required)

[ ] Moderate risk of VTE	Routine, Once, PACU & Post-op
--------------------------	-------------------------------

##### [ ] Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)

( ) Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device	<b>"And" Linked Panel</b>
--	---------------------------

[ ] Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
---	---

[ ] Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
---	-------------------------------------

( ) Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	<b>"And" Linked Panel</b>
---	---------------------------

[ ] Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
---	---

[ ] Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
--	--

( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
( ) patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCl LESS than 30 mL/min
( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

## Moderate Risk of DVT - Non-Surgical (Selection Required)

## Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

### Central line

#### History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

## Estrogen therapy

### Moderate or major surgery (not for cancer)

### Major surgery within 3 months of admission

---

Digitized by srujanika@gmail.com

### ]] Moderate Risk (Selection Required)

<input type="checkbox"/> Moderate Risk (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device	<b>"And" Linked Panel</b>
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis	<b>"And" Linked Panel</b>

<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<b>( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)</b>	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

**( ) HIGH Risk of DVT - Surgical (Selection Required)**

**High Risk Definition**

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

**[ ] High Risk (Selection Required)**

<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
---	-------------------------------

**[ ] High Risk Pharmacological Prophylaxis - Surgical Patient  
(Single Response) (Selection Required)**

Contraindications exist for pharmacologic prophylaxis

Routine, Once  
No pharmacologic VTE prophylaxis due to the following contraindication(s):  
PACU & Post-op

enoxaparin (LOVENOX) injection (Single Response)  
(Selection Required)

40 mg, subcutaneous, daily at 0600, Starting S+1

enoxaparin (LOVENOX) syringe

30 mg, subcutaneous, daily at 0600, Starting S+1

patients with CrCL LESS than 30 mL/min

For Patients with CrCL LESS than 30 mL/min

( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

Mechanical Prophylaxis (Single Response) (Selection Required)

( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
( ) Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

( ) HIGH Risk of DVT - Non-Surgical (Selection Required)

High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

High Risk (Selection Required)

<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
---	-------------------------------

High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)

( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
---	---

( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

40 mg, subcutaneous, daily at 1700, Starting S

( ) patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCL LESS than 30 mL/min
--	--

( ) patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min

30 mg, subcutaneous, 2 times daily, Starting S  
For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

( ) patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:

[ ] Mechanical Prophylaxis (Single Response) (Selection Required)

( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
( ) Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

( ) HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)

#### High Risk Definition

Both pharmacologic AND mechanical prophylaxis must be addressed.

One or more of the following medical conditions:

Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)

Severe fracture of hip, pelvis or leg

Acute spinal cord injury with paresis

Multiple major traumas

Abdominal or pelvic surgery for CANCER

Acute ischemic stroke

History of PE

[ ] High Risk (Selection Required)

[ ] High risk of VTE Routine, Once, PACU & Post-op

[ ] High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)

( ) Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
---	---

( ) apixaban (ELIQUIS) tablet	2.5 mg, oral, every 12 hours, Starting S+1 Indications:
-------------------------------	--

( ) aspirin chewable tablet	162 mg, oral, daily, Starting S+1
-----------------------------	-----------------------------------

( ) aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
---	-----------------------------------

( ) enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

( ) enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
( ) enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1

( ) enoxaparin (LOVENOX) syringe - For Patients with CrCl LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCl LESS than 30 mL/min.
---	---

( ) enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
( ) enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
( ) fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
( ) heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
( ) heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
( ) rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL), Starting S+1 To be Given on Post Op Day 1. Indications:
( ) warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
( ) Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
( ) Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
( ) Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

## Labs

### Labs

<input type="checkbox"/> Hemoglobin and hematocrit	Once, PACU & Post-op
--	----------------------

### Labs - AM

<input type="checkbox"/> Basic metabolic panel	AM draw, Starting S+1 For 1 Occurrences, PACU & Post-op
<input type="checkbox"/> CBC with platelet and differential	AM draw, Starting S+1 For 1 Occurrences, PACU & Post-op
<input checked="" type="checkbox"/> Partial thromboplastin time	AM draw, Starting S+1 For 1 Occurrences, PACU & Post-op
<input type="checkbox"/> Prothrombin time with INR	AM draw, Starting S+1 For 1 Occurrences, PACU & Post-op

### Labs - AM Daily x 3

<input type="checkbox"/> Hemoglobin	AM draw repeats For 3 Occurrences, PACU & Post-op
-------------------------------------	---

## Imaging

### CT

<input type="checkbox"/> CT Cervical Spine Wo Contrast	Routine, 1 time imaging For 1 , PACU & Post-op
<input type="checkbox"/> CT Thoracic Spine Wo Contrast	Routine, 1 time imaging For 1 , PACU & Post-op
<input type="checkbox"/> CT Lumbar Spine Wo Contrast	Routine, 1 time imaging For 1 , PACU & Post-op

### X-ray

<input type="checkbox"/> Chest 1 Vw Portable	Routine, 1 time imaging For 1 , PACU & Post-op
<input type="checkbox"/> Chest 1 Vw Portable in AM	Routine, 1 time imaging, Starting S+1 For 1 , PACU & Post-op

[ ] XR Spine Scoliosos 2-3 Views	Routine, 1 time imaging For 1 Please add 32 millimeter image calibration necklace to the field of view., PACU & Post-op
[ ] Cervical Spine 1 Vw	Routine, 1 time imaging For 1 , PACU & Post-op
[ ] Cervical Spine 2 Or 3 Vw	Routine, 1 time imaging For 1 , PACU & Post-op
[ ] Thoracic Spine 1 Vw	Routine, 1 time imaging For 1 , PACU & Post-op
[ ] Lumbar Spine 1 Vw	Routine, 1 time imaging For 1 , PACU & Post-op
[ ] Thoracolumbar Spine 2 Vw	Routine, 1 time imaging For 1 , PACU & Post-op

## Respiratory

### Respiratory

[ ] Oxygen therapy - Simple face mask	Routine, Continuous Device: Simple Face Mask Rate in liters per minute: 6 lpm Rate in tenths of a liter per minute: O2 %: Titrate to keep O2 Sat Above: 92% Indications for O2 therapy: Wean prn., PACU & Post-op
[ ] Incentive spirometry	Routine, Every hour While awake., PACU & Post-op

## Consults

For Physician Consult orders use sidebar

### Ancillary Consults

[ ] Consult to Case Management	Consult Reason: PACU & Post-op
[ ] Consult to Social Work	Reason for Consult: PACU & Post-op
[ ] Consult to PT eval and treat	Reasons for referral to Physical Therapy (mark all applicable): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation( if values are very abnormal): Weight Bearing Status: PACU & Post-op
[ ] Consult PT wound care	Special Instructions: Location of Wound? PACU & Post-op
[ ] Consult to OT eval and treat	Reason for referral to Occupational Therapy (mark all that apply): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation( if values are very abnormal): Weight Bearing Status: PACU & Post-op
[ ] Consult to Nutrition Services	Reason For Consult? Purpose/Topic: PACU & Post-op
[ ] Consult to Spiritual Care	Reason for consult? PACU & Post-op
[ ] Consult to Speech Language Pathology	Routine, Once Reason for consult: PACU & Post-op

[ ] Consult to Wound Ostomy Care nurse	Reason for consult: Reason for consult: Reason for consult: Reason for consult: Consult for NPWT: Reason for consult: PACU & Post-op
[ ] Consult to Respiratory Therapy	Reason for Consult? PACU & Post-op

#### Physician Consults

[ ] Consult Intensive Care	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated? PACU & Post-op
[ ] Consult Physical Medicine Rehab	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated? PACU & Post-op