

Amphotericin B LIPOSOMAL (Ambisome) Infusion (RESTRICTED) [2389]

!!! LIPOSOMAL formulation !!!

Amphotericin B Liposomal (AMBISOME) infusion is RESTRICTED to Infectious Diseases (ID), Solid Organ Transplant (SOT), Bone Marrow Transplant (BMT), and Hematology/Oncology (Heme/Onc) specialists.

General

Nursing

IV Fluids

Medications

Hydration

<input checked="" type="checkbox"/> sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours Give one hour PRIOR to amphotericin B infusion
<input checked="" type="checkbox"/> sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours, Starting H+3 Hours Give one hour AFTER amphotericin B infusion

Pre-medications (for fever and chills)

Acute, infusion-related reactions, including fever, chills, and rigors may occur during or shortly after (within 1-2 hours) the administration of Amphotericin B.

acetaminophen (TYLENOL) (Single Response)

<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, every 24 hours, Starting H+30 Minutes
<input type="checkbox"/> acetaminophen (TYLENOL) 325 mg/10.15 mL suspension	650 mg, oral, every 24 hours, Starting H+30 Minutes If patient cannot swallow...
<input type="checkbox"/> acetaminophen (TYLENOL) suppository	650 mg, rectal, every 24 hours, Starting H+30 Minutes

diphenhydramine (BENADRYL) (Single Response)

<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection	25 mg, intravenous, every 24 hours, Starting H+30 Minutes Administered via slow IV push
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) tablet	25 mg, oral, every 24 hours, Starting H+30 Minutes
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) 12.5 mg/5 mL elixir	25 mg, oral, every 24 hours, Starting H+30 Minutes If patient cannot swallow...

hydrocortisone (SOLU-CORTEF) IV (Single Response)

<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection	25 mg, intravenous, every 24 hours, Starting H+30 Minutes Administered via IV push
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection	50 mg, intravenous, every 24 hours, Starting H+30 Minutes Administered via IV push

Dextrose 5% Flush

<input checked="" type="checkbox"/> dextrose 5% flush syringe	10 mL, intravenous, every 24 hours Flush line immediately BEFORE starting amphotericin infusion
<input checked="" type="checkbox"/> dextrose 5% flush syringe	10 mL, intravenous, every 24 hours, Starting H+3 Hours Flush line immediately AFTER amphotericin infusion

AMPHOTERICIN B (AMBISOME) 3-6 mg/kg/day

<input type="checkbox"/> amphotericin B (liposomal) (AmBisome) infusion	3 mg/kg/day, intravenous, for 2 Hours, every 24 hours, Starting H+1 Hours See orders for pre-/post-hydration and/or pre-medications. Infuse over 2 hours. RESTRICTED to Infectious Diseases (ID), Solid Organ Transplant (SOT), Bone Marrow Transplant (BMT), and Hematology/Oncology (Heme/Onc) specialists. Are you an ID, SOT, BMT, or Heme/Onc specialist or ordering on behalf of one? Reason for Therapy:
---	---

Labs

Laboratory

<input type="checkbox"/>	Basic metabolic panel - STAT	STAT For 1 Occurrences
<input type="checkbox"/>	Basic metabolic panel	Once
<input type="checkbox"/>	Magnesium level	Once
<input type="checkbox"/>	Phosphorus level	Once

Cardiology

Imaging

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Additional Orders