

Amphotericin B Conventional (Fungizone) Infusion (RESTRICTED) [2390]

!!! CONVENTIONAL formulation !!!

Amphotericin B conventional (FUNGIZONE) infusion is RESTRICTED to Infectious Diseases (ID), Solid Organ Transplant (SOT), Bone Marrow Transplant (BMT), and Hematology/Oncology (Heme/Onc) specialists.

General

Nursing

IV Fluids

Medications

Hydration

[X] sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours Give one hour PRIOR to amphotericin B infusion
[X] sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours, Starting H+5 Hours Give one hour AFTER amphotericin B infusion

Pre-medications (for fever and chills)

Acute, infusion-related reactions, including fever, chills, and rigors may occur during or shortly after (within 1-2 hours) the administration of Amphotericin B.

[] acetaminophen (TYLENOL) (Single Response)	
() acetaminophen (TYLENOL) tablet	650 mg, oral, every 24 hours, Starting H+30 Minutes
() acetaminophen (TYLENOL) 325 mg/10.15 mL suspension	650 mg, oral, every 24 hours, Starting H+30 Minutes If patient cannot swallow...
() acetaminophen (TYLENOL) suppository	650 mg, rectal, every 24 hours, Starting H+30 Minutes
[] diphenhydramine (BENADRYL) (Single Response)	
() diphenhydramine (BENADRYL) injection	25 mg, intravenous, every 24 hours, Starting H+30 Minutes Administered via slow IV push
() diphenhydramine (BENADRYL) tablet	25 mg, oral, every 24 hours, Starting H+30 Minutes
() diphenhydramine (BENADRYL) 12.5 mg/5 mL elixir	25 mg, oral, every 24 hours, Starting H+30 Minutes If patient cannot swallow...
[] hydrocortisone (SOLU-CORTEF) IV (Single Response)	
() hydrocortisone sodium succinate (Solu-CORTEF) injection	25 mg, intravenous, every 24 hours, Starting H+30 Minutes Administered via IV push
() hydrocortisone sodium succinate (Solu-CORTEF) injection	50 mg, intravenous, every 24 hours, Starting H+30 Minutes Administered via IV push

Dextrose 5% Flush

[X] dextrose 5% flush syringe	10 mL, intravenous, every 24 hours Flush line immediately BEFORE starting amphotericin infusion
[X] dextrose 5% flush syringe	10 mL, intravenous, every 24 hours, Starting H+5 Hours Flush line immediately AFTER amphotericin infusion

amphotericin B Conventional (FUNGIZONE) (Single Response)

() amphotericin B (FUNGIZONE) in dextrose 5% 500 mL IVPB (PERIPHERAL LINE)	intravenous, for 4 Hours, every 24 hours, Starting H+1 Hours See orders for pre-/post-hydration and/or pre-medications RESTRICTED to Infectious Diseases (ID), Solid Organ Transplant (SOT), Bone Marrow Transplant (BMT), and Hematology/Oncology (Heme/Onc) specialists. Are you an ID, SOT, BMT, or Heme/Onc specialist or ordering on behalf of one? Reason for Therapy:
---	---

() amphotericin B (FUNGIZONE) in dextrose 5% 250 mL
IVPB (CENTRAL LINE)

intravenous, for 4 Hours, every 24 hours, Starting H+1 Hours
See orders for pre-/post-hydration and/or pre-medications
RESTRICTED to Infectious Diseases (ID), Solid Organ
Transplant (SOT), Bone Marrow Transplant (BMT), and
Hematology/Oncology (Heme/Onc) specialists. Are you an ID,
SOT, BMT, or Heme/Onc specialist or ordering on behalf of
one?
Reason for Therapy:

Labs

Laboratory

<input type="checkbox"/> Basic metabolic panel - STAT	STAT For 1 Occurrences
<input type="checkbox"/> Basic metabolic panel	Once
<input type="checkbox"/> Magnesium level	Once
<input type="checkbox"/> Phosphorus level	Once

Cardiology

Imaging

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Additional Orders