

General

Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Post-op
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Post-op
<input type="checkbox"/>	Acute Renal Failure	Post-op
<input type="checkbox"/>	Acute Respiratory Failure	Post-op
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Post-op
<input type="checkbox"/>	Anemia	Post-op
<input type="checkbox"/>	Bacteremia	Post-op
<input type="checkbox"/>	Bipolar disorder, unspecified	Post-op
<input type="checkbox"/>	Cardiac Arrest	Post-op
<input type="checkbox"/>	Cardiac Dysrhythmia	Post-op
<input type="checkbox"/>	Cardiogenic Shock	Post-op
<input type="checkbox"/>	Decubitus Ulcer	Post-op
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Post-op
<input type="checkbox"/>	Disorder of Liver	Post-op
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Post-op
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Post-op
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Post-op
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Post-op
<input type="checkbox"/>	Other Alteration of Consciousness	Post-op
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Post-op
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Post-op
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Post-op
<input type="checkbox"/>	Protein-calorie Malnutrition	Post-op
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Post-op
<input type="checkbox"/>	Schizophrenia Disorder	Post-op
<input type="checkbox"/>	Sepsis	Post-op
<input type="checkbox"/>	Septic Shock	Post-op
<input type="checkbox"/>	Septicemia	Post-op
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Post-op
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Post-op

Elective Outpatient, Observation, or Admission (Single Response)

<input type="radio"/>	Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
<input type="radio"/>	Outpatient observation services under general supervision	Diagnosis: Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
<input type="radio"/>	Outpatient in a bed - extended recovery	Diagnosis: Admitting Physician: Bed request comments: PACU & Post-op
<input type="radio"/>	Admit to Inpatient	Diagnosis: Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op

Admission or Observation (Single Response)

Patient has active outpatient status order on file

- | | |
|--|--|
| <input type="checkbox"/> Admit to Inpatient | Diagnosis:
Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Outpatient observation services under general supervision | Diagnosis:
Admitting Physician:
Patient Condition:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Outpatient in a bed - extended recovery | Diagnosis:
Admitting Physician:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Admission (Single Response)

Patient has active status order on file

- | | |
|---|--|
| <input type="checkbox"/> Admit to inpatient | Diagnosis:
Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Transfer (Single Response)

Patient has active inpatient status order on file

- | | |
|---|---|
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Code Status

- | | |
|--|--|
| <input type="checkbox"/> Full Code | Code Status decision reached by:
Post-op |
| <input type="checkbox"/> DNR (Do Not Resuscitate) (Selection Required) | |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | Does patient have decision-making capacity?
Post-op |

<input type="checkbox"/> Consult to Palliative Care Service	Priority: Reason for Consult? Order? Name of referring provider: Enter call back number:
<input type="checkbox"/> Consult to Social Work	Reason for Consult: Post-op
<input type="checkbox"/> Modified Code	Does patient have decision-making capacity? Modified Code restrictions: Post-op
<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

Isolation

<input type="checkbox"/> Airborne isolation status	
<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics.	Once, Sputum, Post-op
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op

Nursing

Notify Physician

<input checked="" type="checkbox"/> Notify Physician for vitals:	Routine, Until discontinued, Starting S Temperature greater than: Temperature less than: Systolic BP greater than: 150 Systolic BP less than: 90 Diastolic BP greater than: Diastolic BP less than: MAP less than: Heart rate greater than (BPM): 100 Heart rate less than (BPM): 60 Respiratory rate greater than: 26 Respiratory rate less than: 12 SpO2 less than: 95
<input checked="" type="checkbox"/> Notify Attending	Routine, Until discontinued, Starting S, Neck swelling or difficulty breathing, Post-op
<input checked="" type="checkbox"/> Notify Attending	Routine, Until discontinued, Starting S, Tingling around mouth or fingers/hands, cramping in hands, Post-op
<input checked="" type="checkbox"/> Notify Attending	Routine, Until discontinued, Starting S, Calcium less than 7, Post-op
<input checked="" type="checkbox"/> Notify Attending	Routine, Until discontinued, Starting S, Nausea/vomiting not controlled, Post-op
<input checked="" type="checkbox"/> Notify Attending	Routine, Until discontinued, Starting S, Inability to keep down oral medications, Post-op

Vitals

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 4 hours For 2 Occurrences, Post-op
---	---

Vital signs - T/P/R/BP Routine, Every 8 hours, Starting H+8 Hours, Post-op

Activity

Out of bed Routine, 3 times daily For Until specified
Specify: Out of bed
Out of bed for 1 hour at a time, or as tolerated, Post-op

Nursing Assessment

Neurological assessment Routine, Every 4 hours
Assessment to Perform:
Post-op

Strict intake and output Routine, Every 8 hours
Record all oral and IV fluid intake and all output including drains, tubes, nasogastric tube and foley catheter, Post-op

Height and weight Routine, Once, Post-op

Nursing Interventions

Head of bed Routine, Until discontinued, Starting S
Head of bed:
Unless contraindicated, Post-op

Patient education Routine, Once
Patient/Family:
Education for:
Post-op

Wound Care

Provide suture tray to patient bedside Routine, Once, Post-op

Diet

NPO Diet effective now, Starting S
NPO:
Pre-Operative fasting options:
Post-op

Diet - Clear Liquids Diet effective now, Starting S
Diet(s): Clear Liquids
Advance Diet as Tolerated?
Liquid Consistency:
Fluid Restriction:
Foods to Avoid:
Post-op

Medications

Medications

calcium carbonate (TUMS) chewable tablet 3,000 mg, oral, 3 times daily, Post-op

calcium gluconate IVPB 1 g, intravenous, for 30 Minutes, every 30 min
Discontinue if ionized calcium GREATER than 1.25 mmol/L

calcitriol (ROCALTRON) capsule 1 mcg, oral, 2 times daily, Post-op

Pain Management

For Mild Pain - acetaminophen (TYLENOL) tablet 500 mg, oral, every 6 hours PRN, mild pain (score 1-3),
Post-op

For Moderate Pain - acetaminophen (TYLENOL) tablet 1,000 mg, oral, every 8 hours PRN, moderate pain (score
4-6), Post-op

For Severe Pain - acetaminophen-codeine (TYLENOL
WITH CODEINE #3) 300-30 mg per tablet 1 tablet, oral, every 6 hours PRN, severe pain (score 7-10),
Post-op

Labs

Labs

[] Parathyroid hormone	Once, Starting H+30 Minutes For 1 Occurrences Check PTH level 30 mins post operation, PACU
[] Calcium level	Every 6 hours For Until specified Check every 6 hours post operation until discontinued, PACU & Post-op
[] Ionized calcium	Every 6 hours For Until specified Check every 6 hours post operation until discontinued, PACU & Post-op

Consults

For Physician Consult orders use sidebar

Consult

[] Consult Nephrology for high calcium HD dialysate	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated?
--	---