

# Non-Emergent Reversal of Warfarin (Coumadin) Anticoagulation (Procedure or Supratherapeutic INR without Bleeding) [4057]

DISCONTINUE WARFARIN (COUMADIN)

\*Vitamin K is not indicated for the reversal of DOACs (apixaban, rivaroxaban, edoxaban or dabigatran)\*

## Nursing

### Nursing

[X] Bleeding precautions Monitor for bleeding. Contact provider for any signs or symptoms of bleeding occur.

## Medications

### Medications - Non-Emergent (Procedure or Supratherapeutic INR) (Single Response)

If the patient has an active, life-threatening bleed use "Emergent Reversal of warfarin (Coumadin) Anticoagulation" order set.

#### ( ) INR GREATER than 10 (Single Response)

##### ( ) Oral: preferred (Single Response)

( ) phytonadione (MEPHYTON) suspension 2.5 mg, oral, once  
Indication:

( ) phytonadione (MEPHYTON) suspension 5 mg, oral, once  
Indication:

##### ( ) Subcutaneous: Variable absorption; alternative if patient is NPO or concern for poor gut absorption (Single Response)

( ) phytonadione \*(AQUA-MEPHYTON) subcutaneous 2.5 mg, subcutaneous, once  
Indication:

( ) phytonadione \*(AQUA-MEPHYTON) subcutaneous 5 mg, subcutaneous, once  
Indication:

##### ( ) Intravenous: Reserved for situations where other routes are not feasible. Anaphylactic reactions have been described with IV use. Monitoring recommended. (Single Response)

( ) phytonadione (AQUA-Mephyton) IV piggyback 5 mg, intravenous, for 30 Minutes, once  
Indication:

( ) phytonadione (AQUA-Mephyton) IV piggyback 10 mg, intravenous, for 30 Minutes, once  
Indication:

#### ( ) INR LESS than 10 (Single Response)

INR less than 10 Vitamin K is not recommended by ACCP guidelines for INR less than 10; optional treatment at provider discretion

##### ( ) Oral: preferred (Single Response)

( ) phytonadione (MEPHYTON) suspension 1 mg, oral, once  
Indication:

( ) phytonadione (MEPHYTON) suspension 2.5 mg, oral, once  
Indication:

##### ( ) Subcutaneous: Variable absorption; alternative if patient is NPO or concern for poor gut absorption (Single Response)

( ) phytonadione \*(AQUA-MEPHYTON) subcutaneous 1 mg, subcutaneous, once  
Indication:

( ) phytonadione \*(AQUA-MEPHYTON) subcutaneous 2.5 mg, subcutaneous, once  
Indication:

## Laboratory

### Laboratory

[X] Prothrombin time with INR AM draw For 1 Occurrences

CBC with platelet and differential

AM draw For 1 Occurrences

CBC with platelet and differential

Every 4 hours, Starting S For 3 Occurrences