

Nursing

Nursing

[X] Weigh patient	STAT, Once For 1 Occurrences Weigh patient if baseline measured weight not in medical record. Notify pharmacist if patient cannot be weighed.
[X] Notify pharmacist (Tandem)	Routine, Until discontinued, Starting S, -Notify pharmacist if a physician writes heparin orders. Heparin dose or infusion changes to be made by pharmacist only. -Notify pharmacist if compatibility with other IV drugs or if IV access is a concern. -Notify pharmacist IMMEDIATELY if the patient transfers or has an off unit procedure. -Notify pharmacist IMMEDIATELY if heparin infusion is stopped for any reason. -Notify pharmacist if patient cannot be weighed.
[X] Notify pharmacist and physician (Tandem)	Routine, Until discontinued, Starting S, -Notify pharmacist and physician if Platelet count less than 150,000 microliter or a 50% drop from baseline or hematoma, bleeding or suspected bleeding occurs.
[X] Heparin instructions (Tandem)	Routine, Until discontinued, Starting S -Document: Tandem Heart infusate rate and heparin concentration hourly. -Do not interrupt heparin infusion unless ordered. -Notify pharmacist regarding compatibility with other IV drugs if access is a concern. -Notify pharmacist immediately if heparin infusion is stopped for any reason.

Medications

Tandem Heart Infusate Heparin (Single Response)

(X) Standard Infusate: Heparin 90,000 units/1000 mL Normal Saline	900 Units/hr, Device, continuous - Document: Hourly infusate flow rates and heparin concentration. - Do not interrupt heparin infusion unless ordered. - Notify pharmacist immediately if heparin infusion is stopped for any reason.
() ALTERNATE Infusate: Heparin 45,000 units/1000 mL Normal Saline	450 Units/hr, Device, continuous - Document: Hourly infusate flow rates and heparin concentration. - Do not interrupt heparin infusion unless ordered. - Notify pharmacist immediately if heparin infusion is stopped for any reason.

Supplemental Systemic Heparin

[] Supplemental systemic heparin HEParin 25,000 unit/500 mL (50 unit/mL)	intravenous, titrated Begin when PTT sub-therapeutic on Tandem Infusate Heparin alone. Indication: Other Therapeutic Monitoring Target: PTT - Other Specify Target: 60 - 80 sec Specify Indication: Tandem Heart
---	---

Labs

Labs-Initiation

<input type="checkbox"/> Partial thromboplastin time	STAT For 1 Occurrences Draw blood for PTT from arm that does not have heparin infusion. If there is no other access than the heparin line, stop the heparin for 10 minutes, flush the line, aspirate 10 mL of blood to waste, obtain sample, and reflush the line after drawing specimen.
<input type="checkbox"/> Prothrombin time with INR	STAT For 1 Occurrences Do not draw blood from the arm that has heparin infusion. Do not draw from heparin flushed lines.
<input type="checkbox"/> CBC hemogram	STAT For 1 Occurrences Do not draw blood from the arm that has heparin infusion. Do not draw from heparin flushed lines.

Labs-Continuing

<input checked="" type="checkbox"/> CBC hemogram	Daily, Starting S+1 For 3 Occurrences Daily while on heparin. Do not draw blood from the arm that has heparin infusion. Do not draw from heparin flushed lines.
<input checked="" type="checkbox"/> Partial thromboplastin time	Every 4 hours, Starting H+4 Hours For 1 Occurrences Draw 4 hours after initiation of heparin infusion. Draw blood for PTT/ Anti Xa UFH from arm that does not have heparin infusion. If there is no other access than the heparin line, stop the heparin for 10 minutes, flush the line, aspirate 10 mL of blood to waste, obtain sample, and reflush the line after drawing specimen.
<input checked="" type="checkbox"/> Occult blood, stool	Conditional Frequency, Starting S For 3 Occurrences, Stool quiac stools when there is a hemoglobin decrease of more than 2 gm/dL in a 24 hour period

Consults

Pharmay Consult

<input checked="" type="checkbox"/> Pharmacy consult for Heparin Tandem Heart	STAT, Until discontinued, Starting S
---	--------------------------------------