

Retinopathy of Prematurity Exam [714]

General

Nursing

Nursing

<input type="checkbox"/> Provide equipment / supplies at bedside- Retinopathy	Routine, Once For 1 Occurrences Supplies: Other (specify) Other: Have the sterile lid speculum and lens loop at the infant's bedside.
<input type="checkbox"/> Apply phototherapy mask to eyes for 4 hours post ROP exam	Routine, Continuous Phototherapy: Double Apply phototherapy mask to eyes 4 hours post dilation.

IV Fluids

Medications

Medications

<input checked="" type="checkbox"/> cyclopentolate-phenylephrine (CYCLOMYDRYL) 0.2-1 % ophthalmic solution	1 drop, Both Eyes, PRN, instill every 5 minutes for 3 doses To be administered 45 minutes PRIOR to eye exam.
<input checked="" type="checkbox"/> sucrose 24 % (TOOT SWEET) oral solution	0.1 mL, oral, once PRN, mild pain (score 1-3), pain
<input checked="" type="checkbox"/> proparacaine (ALCAINE) 0.5 % ophthalmic solution	1 drop, Both Eyes, PRN, for eye exam To be administered 30 seconds PRIOR to eye exam.
<input type="checkbox"/> artificial tears(hypromellose) (SYSTANE/GENTEAL) 0.3 % ophthalmic gel	1 application, Both Eyes, PRN, dry eyes, for eye exam To be administered for each eye exam.

VTE

Labs

Cardiology

Diagnostic Imaging

Other Diagnostic Studies

Respiratory

Rehab

Consults

Additional Orders