

General

Common Present on Admission Diagnosis

<input type="checkbox"/>	Acidosis	Post-op
<input type="checkbox"/>	Acute Post-Hemorrhagic Anemia	Post-op
<input type="checkbox"/>	Acute Renal Failure	Post-op
<input type="checkbox"/>	Acute Respiratory Failure	Post-op
<input type="checkbox"/>	Acute Thromboembolism of Deep Veins of Lower Extremities	Post-op
<input type="checkbox"/>	Anemia	Post-op
<input type="checkbox"/>	Bacteremia	Post-op
<input type="checkbox"/>	Bipolar disorder, unspecified	Post-op
<input type="checkbox"/>	Cardiac Arrest	Post-op
<input type="checkbox"/>	Cardiac Dysrhythmia	Post-op
<input type="checkbox"/>	Cardiogenic Shock	Post-op
<input type="checkbox"/>	Decubitus Ulcer	Post-op
<input type="checkbox"/>	Dementia in Conditions Classified Elsewhere	Post-op
<input type="checkbox"/>	Disorder of Liver	Post-op
<input type="checkbox"/>	Electrolyte and Fluid Disorder	Post-op
<input type="checkbox"/>	Intestinal Infection due to Clostridium Difficile	Post-op
<input type="checkbox"/>	Methicillin Resistant Staphylococcus Aureus Infection	Post-op
<input type="checkbox"/>	Obstructive Chronic Bronchitis with Exacerbation	Post-op
<input type="checkbox"/>	Other Alteration of Consciousness	Post-op
<input type="checkbox"/>	Other and Unspecified Coagulation Defects	Post-op
<input type="checkbox"/>	Other Pulmonary Embolism and Infarction	Post-op
<input type="checkbox"/>	Phlebitis and Thrombophlebitis	Post-op
<input type="checkbox"/>	Protein-calorie Malnutrition	Post-op
<input type="checkbox"/>	Psychosis, unspecified psychosis type	Post-op
<input type="checkbox"/>	Schizophrenia Disorder	Post-op
<input type="checkbox"/>	Sepsis	Post-op
<input type="checkbox"/>	Septic Shock	Post-op
<input type="checkbox"/>	Septicemia	Post-op
<input type="checkbox"/>	Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled	Post-op
<input type="checkbox"/>	Urinary Tract Infection, Site Not Specified	Post-op

Elective Outpatient, Observation, or Admission (Single Response)

<input type="radio"/>	Elective outpatient procedure: Discharge following routine recovery	Routine, Continuous, PACU & Post-op
<input type="radio"/>	Outpatient observation services under general supervision	Admitting Physician: Patient Condition: Bed request comments: PACU & Post-op
<input type="radio"/>	Outpatient in a bed - extended recovery	Admitting Physician: Bed request comments: PACU & Post-op
<input type="radio"/>	Admit to Inpatient	Admitting Physician: Level of Care: Patient Condition: Bed request comments: Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights. PACU & Post-op

Admission or Observation (Single Response)

Patient has active outpatient status order on file

- | | |
|--|--|
| <input type="checkbox"/> Admit to Inpatient | Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Outpatient observation services under general supervision | Admitting Physician:
Patient Condition:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Outpatient in a bed - extended recovery | Admitting Physician:
Bed request comments:
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Admission (Single Response)

Patient has active status order on file

- | | |
|---|--|
| <input type="checkbox"/> Admit to inpatient | Admitting Physician:
Level of Care:
Patient Condition:
Bed request comments:
Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.
PACU & Post-op |
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Transfer (Single Response)

Patient has active inpatient status order on file

- | | |
|---|---|
| <input type="checkbox"/> Transfer patient | Level of Care:
Bed request comments:
Scheduling/ADT |
| <input type="checkbox"/> Return to previous bed | Routine, Until discontinued, Starting S, Scheduling/ADT |

Code Status

- | | |
|--|--|
| <input type="checkbox"/> Full Code | Code Status decision reached by:
Post-op |
| <input type="checkbox"/> DNR (Do Not Resuscitate) (Selection Required) | |
| <input type="checkbox"/> DNR (Do Not Resuscitate) | Does patient have decision-making capacity?
Post-op |
| <input type="checkbox"/> Consult to Palliative Care Service | Priority:
Reason for Consult?
Order?
Name of referring provider:
Enter call back number: |
| <input type="checkbox"/> Consult to Social Work | Reason for Consult:
Post-op |

<input type="checkbox"/> Modified Code	Does patient have decision-making capacity? Modified Code restrictions: Post-op
<input type="checkbox"/> Treatment Restrictions	Treatment Restriction decision reached by: Specify Treatment Restrictions: Post-op

Isolation

<input type="checkbox"/> Airborne isolation status	
<input type="checkbox"/> Airborne isolation status	Details
<input type="checkbox"/> Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics.	Once, Sputum, Post-op
<input type="checkbox"/> Contact isolation status	Details
<input type="checkbox"/> Droplet isolation status	Details
<input type="checkbox"/> Enteric isolation status	Details

Precautions

<input type="checkbox"/> Aspiration precautions	Post-op
<input type="checkbox"/> Fall precautions	Increased observation level needed: Post-op
<input type="checkbox"/> Latex precautions	Post-op
<input type="checkbox"/> Seizure precautions	Increased observation level needed: Post-op

Nursing

Vital Signs

<input checked="" type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 4 hours Pulse check q2 x12 then every 4 hours if patient is stable. Notify MD if absent pulse or doppler signals or change in vascular exam., Post-op
--	--

Activity

<input type="checkbox"/> Up with assistance to chair	Routine, Daily, Starting S+1 Specify: Out of bed, Up in chair, Up with assistance Additional modifier: Out of bed to chair with assistance, Post-op
<input type="checkbox"/> Ambulate	Routine, 4 times daily, Starting S+1 Specify: with assistance Post-op
<input type="checkbox"/> Bed rest - lay flat for 6 hrs	Routine, Until discontinued, Starting S Bathroom Privileges: Patient must lay flat for 6 hours post-op, Post-op
<input type="checkbox"/> Bed rest	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Bed rest - lay flat for 4 hrs	Routine, Until discontinued, Starting S Patient must lay flat for 4 hours post-op, Post-op

Nursing

<input type="checkbox"/> Intake and output	Routine, Every 8 hours, Post-op
<input type="checkbox"/> Foley catheter - discontinue	Routine, Once, Starting S+1, Post-op
<input type="checkbox"/> Tobacco cessation education	Routine, Once, Post-op
<input type="checkbox"/> Nasogastric tube maintenance	Routine, Until discontinued, Starting S Tube Care Orders: To Continuous Suction Irrigate with 30 cubic cm of saline q4, Post-op
<input type="checkbox"/> Neurological assessment	Routine, Every 2 hours For 6 Occurrences Assessment to Perform: Post CEA, Post-op

<input type="checkbox"/> Assess operative site	Routine, Every 8 hours If patient is status post Carotid Endarterectomy, monitor neck incision for increased swelling, hematoma formation, difficulty swallowing or difficulty speaking., Post-op
<input type="checkbox"/> Assess operative site	Routine, Now then every 8 hours If status post angiogram, monitor angiogram access site for hematoma formation, Post-op
<input type="checkbox"/> No peripheral IV or blood draws on side of anticipated dialysis access, place sign over patient's bed	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Measure drainage	Routine, Every 6 hours Type of drain: Jackson Pratt Post-op
<input type="checkbox"/> Have wound dressings available at bedside	Routine, Once Supplies: Have wound dressings available at bedside, Post-op
<input type="checkbox"/> Request for central wound care equipment	Routine, Once Equipment Requested: Equipment Requested: Special Instructions: Post-op
<input type="checkbox"/> Elevate extremity	Routine, Until discontinued, Starting S Position: Additional instructions: elevate extremity Extremity: Post-op
<input type="checkbox"/> Nursing communication: OK to cannulate AV access for dialysis	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Nursing communication: OK to resume heparin post-op	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Nursing communication: Discontinue heparin post-op	Routine, Until discontinued, Starting S, Post-op

Notify

<input checked="" type="checkbox"/> Notify Physician for vitals:	Routine, Until discontinued, Starting S Temperature greater than: 38.5 Temperature less than: Systolic BP greater than: 160 Systolic BP less than: 90 Diastolic BP greater than: 100 Diastolic BP less than: MAP less than: 60 Heart rate greater than (BPM): 100 Heart rate less than (BPM): 50 Respiratory rate greater than: 25 Respiratory rate less than: 8 SpO2 less than: 94
<input type="checkbox"/> Notify Vascular Surgery team for urine output less than 30 milliliters/hour	Routine, Until discontinued, Starting S, Post-op
<input type="checkbox"/> Notify Vascular Surgery team absent pulses or Doppler signals or change in the vascular exam	Routine, Until discontinued, Starting S, Post-op

Diet

<input type="checkbox"/> NPO	Diet effective now, Starting S NPO: Pre-Operative fasting options: Post-op
<input type="checkbox"/> NPO	Diet effective midnight, Starting S+1 at 12:01 AM NPO: Except meds Pre-Operative fasting options: Post-op

<input type="checkbox"/> Diet - clear liquid. Advance as tolerated	Diet effective now, Starting S Diet(s): Clear Liquids Advance Diet as Tolerated? Yes Target Diet: Advance target diet criteria: Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op
<input type="checkbox"/> Diet - heart healthy	Diet effective now, Starting S Diet(s): Heart Healthy Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op
<input type="checkbox"/> Diet - Renal	Diet effective now, Starting S Diet(s): Renal (80GM Pro, 2-3GM Na, 2-3GM K) Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op
<input type="checkbox"/> Diet - diabetic 1800 Carb	Diet effective now, Starting S Diet(s): Other Diabetic/Cal Diabetic/Calorie: 1800 Kcal/202 gm Carbohydrate Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op
<input type="checkbox"/> Diet - Regular	Diet effective now, Starting S Diet(s): Other Options: Advance Diet as Tolerated? Liquid Consistency: Fluid Restriction: Foods to Avoid: Post-op

IV Fluids

Post-Procedure Hydration (Single Response)

<input type="checkbox"/> Inpatient (Single Response)	
<input type="checkbox"/> Patients with EF LESS than 40% or with evidence of fluid overload	0.5 mL/kg/hr, intravenous, continuous Infuse for 6 hours Post-Procedure
<input type="checkbox"/> Patients with EF GREATER than 40% or no evidence of fluid overload	1 mL/kg/hr, intravenous, continuous Infuse for 6 hours Post-Procedure
<input type="checkbox"/> Outpatient (Single Response)	
<input type="checkbox"/> Patients with EF LESS than 40% or with evidence of fluid overload	0.5 mL/kg/hr, intravenous, continuous Infuse for 6 hours Post-Procedure or until discharge, whichever comes first.
<input type="checkbox"/> Patients with EF GREATER than 40% or no evidence of fluid overload	1 mL/kg/hr, intravenous, continuous Infuse for 6 hours Post-Procedure or until discharge, whichever comes first.

IV Fluids (Single Response)

<input type="checkbox"/> sodium chloride 0.9 % infusion	75 mL/hr, intravenous, continuous, Post-op
<input type="checkbox"/> lactated Ringer's infusion	75 mL/hr, intravenous, continuous, Post-op
<input type="checkbox"/> dextrose 5 % and sodium chloride 0.45 % with potassium chloride 20 mEq/L infusion	75 mL/hr, intravenous, continuous, Post-op
<input type="checkbox"/> dextrose 5%-0.45% sodium chloride infusion	75 mL/hr, intravenous, continuous, Post-op

<input type="checkbox"/> sodium chloride 0.45 % infusion	75 mL/hr, intravenous, continuous, Post-op
<input type="checkbox"/> sodium chloride 0.45 % 1,000 mL with sodium bicarbonate 75 mEq/L infusion	75 mL/hr, intravenous, continuous, Post-op

Medications

Respiratory

<input type="checkbox"/> Scheduled - albuterol nebulizer	2.5 mg, nebulization, Respiratory Therapy - every 6 hours, Post-op Aerosol Delivery Device:
<input type="checkbox"/> As needed - albuterol nebulizer	2.5 mg, nebulization, every 6 hours PRN, wheezing, Post-op Aerosol Delivery Device:
<input type="checkbox"/> Scheduled - ipratropium nebulizer	0.5 mg, nebulization, Respiratory Therapy - every 6 hours, Post-op Aerosol Delivery Device:
<input type="checkbox"/> As needed - ipratropium nebulizer	0.5 mg, nebulization, every 6 hours PRN, wheezing, Post-op Aerosol Delivery Device:
<input checked="" type="checkbox"/> Incentive spirometry	Routine, Every 2 hours while awake, Post-op

Anti-hypertensives

<input type="checkbox"/> hydrALAZINE (APRESOLINE) injection	10 mg, intravenous, every 4 hours PRN, high blood pressure, SBP GREATER than 140 mmHg, Post-op Hold if heart rate is GREATER than 100. HOLD parameters for this order: Contact Physician if:
<input type="checkbox"/> labetalol (TRANDATE) injection	10 mg, intravenous, every 4 hours PRN, high blood pressure, SBP GREATER than 140 mmHg, Post-op hold for heart rate LESS than 60.

Anti-platelets

<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	oral, daily, Post-op
<input type="checkbox"/> clopidogrel (PLAVIX) tablet	300 mg, oral, once, For 1 Doses, Post-op
<input type="checkbox"/> clopidogrel (PLAVIX) tablet	75 mg, oral, daily, Starting S+1, Post-op

Statin Therapy (Single Response)

<input type="checkbox"/> simvastatin (ZOCOR) tablet	40 mg, oral, nightly, Post-op
<input type="checkbox"/> simvastatin (ZOCOR) tablet	20 mg, oral, nightly, Post-op
<input type="checkbox"/> atorvastatin (LIPITOR) tablet	40 mg, oral, nightly, Post-op
<input type="checkbox"/> atorvastatin (LIPITOR) tablet	10 mg, oral, nightly, Post-op

Pharmacy Consult: Renal Adjustment

<input checked="" type="checkbox"/> Pharmacy consult to manage dose adjustments for renal function	STAT, Until discontinued, Starting S Adjust dose for:
--	--

Anti-coagulation

<input type="checkbox"/> Pharmacy Consult to Manage Heparin: STANDARD dose protocol (DVT/PE) - with titration boluses	STAT, Until discontinued, Starting S Heparin Indication: Specify: Give initial Bolus Monitoring: Anti-Xa
<input type="checkbox"/> HEParin 25,000 unit/500 mL (50 unit/mL)	500 Units/hr, intravenous, continuous, Post-op Indication: Therapeutic Monitoring Target:
<input type="checkbox"/> Resume heparin (Single Response)	
<input type="checkbox"/> Ok to resume heparin post-op	Routine, Once For 1 Occurrences, Post-op
<input type="checkbox"/> Discontinue heparin post-op	Routine, Once For 1 Occurrences, Post-op
<input type="checkbox"/> Heparin bolus and infusion	
<input type="checkbox"/> HEParin (porcine) injection	80 Units/kg, intravenous, once, Starting S, For 1 Doses

<input type="checkbox"/> HEParin 25,000 unit/500 mL (50 unit/mL)	18 Units/kg/hr, intravenous, titrated, Starting S, Post-op Indication: Peripheral vascular disease Therapeutic Monitoring Target: PTT - 61 - 112 sec Specify Target: None - Non-titrated
<input type="checkbox"/> enoxaparin (LOVENOX)	1 mg/kg, subcutaneous, every 12 hours, Post-op
<input type="checkbox"/> For renal impairment (GFR <30) - enoxaparin (LOVENOX)	1 mg/kg, subcutaneous, daily at 1700, Post-op FOR PATIENTS WITH CRCL OF LESS THAN 30 MILLILITERS PER MINUTE OR PATIENTS WITH ESRD.
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	Routine, Until discontinued, Starting S Indication:

PostOp Antibiotics: For Patients LESS than or EQUAL to 120 kg (Single Response)

<input checked="" type="checkbox"/> ceFAZolin (ANCEF) IV - For Patients LESS than or EQUAL to 120 kg	2 g, intravenous, every 8 hours, For 2 Doses, Post-op Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> If Beta-Lactam Allergic - vancomycin (VANCOGIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Post-op Administer 12 hours after procedure Reason for Therapy: Surgical Prophylaxis

Post-Op Antibiotics: For Patients GREATER than 120 kg (Single Response)

<input checked="" type="checkbox"/> ceFAZolin (ANCEF) IV - For Patients GREATER than 120 kg	3 g, intravenous, every 8 hours, For 2 Doses, Post-op Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> If Beta-Lactam Allergic - vancomycin (VANCOGIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Post-op Administer 12 hours after procedure Reason for Therapy: Surgical Prophylaxis

Multi-modal pain regimen

<input type="checkbox"/> acetaminophen (OFIRMEV) injection and/or lidocaine 5% patch (Single Response)	
<input type="checkbox"/> acetaminophen (OFIRMEV) intravenous solution	1,000 mg, intravenous, for 15 Minutes, every 8 hours, For 2 Doses, Post-op IV acetaminophen (Ofirmev) is restricted to use only in OR, PACU, or ICU areas, and for patients that cannot tolerate oral, per tube, or rectal routes of administration. Do you attest that this restriction has been met?
<input type="checkbox"/> acetaminophen (OFIRMEV) intravenous solution	1,000 mg, intravenous, for 15 Minutes, once PRN, moderate pain (score 4-6), For 1 Doses, Post-op IV acetaminophen (Ofirmev) is restricted to use only in OR, PACU, or ICU areas, and for patients that cannot tolerate oral, per tube, or rectal routes of administration. Do you attest that this restriction has been met?
<input type="checkbox"/> lidocaine (LIDODERM) 5 %	2 patch, transdermal, for 12 Hours, every 24 hours, For 4 Doses, Post-op
<input type="checkbox"/> gabapentin (NEURONTIN) capsule (Single Response)	
<input type="checkbox"/> GFR GREATER than 60 - gabapentin 300 mg PO TID x 5 days	300 mg, oral, 3 times daily, Post-op
<input type="checkbox"/> GFR BETWEEN 30-60 - gabapentin 200 mg PO TID x 5 days	200 mg, oral, 3 times daily, Post-op
<input type="checkbox"/> GFR BETWEEN 15-30 - gabapentin 100 mg PO TID x 5 days	100 mg, oral, 3 times daily, Post-op
<input type="checkbox"/> GFR LESS THAN 15 or on dialysis - gabapentin 100 mg PO TID x 5 days	100 mg, oral, 3 times daily, Post-op
<input type="checkbox"/> pregabalin (LYRICA) capsule (Single Response)	
<input type="checkbox"/> GFR GREATER than 60 - pregabalin 50 mg PO TID x 5 days	50 mg, oral, 3 times daily, Post-op
<input type="checkbox"/> GFR BETWEEN 30-60 - pregabalin 25 mg PO TID x 5 days	25 mg, oral, 3 times daily, Post-op
<input type="checkbox"/> GFR BETWEEN 15-30 - pregabalin 25 mg PO BID x 5 days	25 mg, oral, 2 times daily, Post-op
<input type="checkbox"/> GFR LESS than 15 or on dialysis - pregabalin 25 mg PO BID x 5 days	25 mg, oral, 2 times daily, Post-op
<input type="checkbox"/> tramadol (ULTRAM) tablet (Single Response)	
<input type="checkbox"/> GFR GREATER than 60 - traMADoL 100 mg PO Q8H	100 mg, oral, every 8 hours, Post-op

() GFR BETWEEN 30-60 - traMADoL 50 mg PO Q8H	50 mg, oral, every 8 hours, Post-op
() Elderly Age GREATER than 75 years old - traMADoL 50 mg PO Q8H	50 mg, oral, every 8 hours, Post-op
[] PRN Breakthrough Pain (Single Response)	
() oxyCODone (ROXICODONE) immediate release tablet	10 mg, oral, every 4 hours PRN, severe pain (score 7-10), Post-op
() hydromorPHONE (DILAUDID) injection	0.5 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op
() morPHINE injection	2 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op

PRN Oral for Moderate Pain (Pain Score 4-6): For Patients LESS than 65 years old (Single Response)
(adjust dose for renal/liver function and age)

() acetaminophen-codeine (TYLENOL #3) tablet OR elixir	"Or" Linked Panel Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
[] acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Give if patient is able to tolerate oral medication.
[] acetaminophen-codeine 300 mg-30 mg /12.5 mL solution	12.5 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Use if patient cannot swallow tablet.
() HYDROcodone-acetaminophen 5/325 (NORCO) tablet OR elixir	"Or" Linked Panel Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
[] HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6)
[] HYDROcodone-acetaminophen (HYCET) 2.5-108.3 mg/5 mL solution	10 mL, oral, every 6 hours PRN, moderate pain (score 4-6) If patient cannot swallow tablet.
() HYDROcodone-acetaminophen 7.5/325 (NORCO) tablet OR elixir	"Or" Linked Panel Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
[] HYDROcodone-acetaminophen (NORCO) 7.5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Give if patient is able to tolerate oral medication.
[] HYDROcodone-acetaminophen (HYCET) 7.5-325 mg/15 mL solution	15 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Use if patient cannot swallow tablet.
() HYDROcodone-acetaminophen 10/325 (NORCO) tablet OR elixir	"Or" Linked Panel Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)
[] HYDROcodone-acetaminophen (NORCO) 10-325 mg per tablet	1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Give if patient is able to tolerate oral medication.
[] HYDROcodone-acetaminophen (HYCET) 7.5-325 mg/15 mL solution	20 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op Use if patient can not swallow tablet.
() traMADoL (ULTRAM) tablet - For eGFR LESS than 30 mL/min, change frequency to every 12 hours)	50 mg, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op (Max Daily dose not to exceed 200 mg/day). Give if patient is able to tolerate oral medication

PRN Oral for Moderate Pain (Pain Score 4-6): For Patients GREATER than 65 years old (Single Response)
NOTICE: Before any pain medication is used you MUST NOTIFY MD and get approval.
(adjust dose for renal/liver function and age)

() acetaminophen-codeine (TYLENOL #3) tablet OR elixir	"Or" Linked Panel
---	--------------------------

Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)

- | | |
|--|--|
| <input type="checkbox"/> acetaminophen-codeine (TYLENOL #3) 300-30 mg per tablet | 1 tablet, oral, once PRN, moderate pain (score 4-6), Post-op
Give if patient is able to tolerate oral medication. |
| <input type="checkbox"/> acetaminophen-codeine 300 mg-30 mg /12.5 mL solution | 12.5 mL, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op
Use if patient cannot swallow tablet. |

- HYDROcodone-acetaminophen 5/325 (NORCO) tablet **"Or" Linked Panel**
OR elixir

Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources)

- | | |
|---|--|
| <input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet | 1 tablet, oral, every 6 hours PRN, moderate pain (score 4-6) |
| <input type="checkbox"/> HYDROcodone-acetaminophen (HYCET) 2.5-108.3 mg/5 mL solution | 10 mL, oral, every 6 hours PRN, moderate pain (score 4-6)
If patient cannot swallow tablet. |

- traMADol (ULTRAM) tablet - For eGFR LESS than 30 mL/min, change frequency to every 12 hours) 25 mg, oral, every 6 hours PRN, moderate pain (score 4-6), Post-op
(Max Daily dose not to exceed 200 mg/day). Give if patient can tolerate oral medication.

PRN IV for Moderate Pain (Pain Score 4-6): For Patients LESS than 65 years old (Single Response)

If you select a PCA option you will not be allowed to also order IV PRN pain medications from this section.
(adjust dose for renal/liver function and age)

- | | |
|---|---|
| <input type="checkbox"/> fentaNYL (SUBLIMAZE) injection | 25 mcg, intravenous, every 2 hour PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |
| <input type="checkbox"/> morphine 2 mg/mL injection | 2 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |
| <input type="checkbox"/> HYDROmorphine (DILAUDID) injection | 0.5 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |

PRN IV for Moderate Pain (Pain Score 4-6): For Patients GREATER than 65 years old (Single Response)

If you select a PCA option you will not be allowed to also order IV PRN pain medications from this section.
(adjust dose for renal/liver function and age)

- | | |
|---|--|
| <input type="checkbox"/> fentaNYL (SUBLIMAZE) injection | 12.5 mcg, intravenous, every 2 hour PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |
| <input type="checkbox"/> morphine 2 mg/mL injection | 1 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |
| <input type="checkbox"/> HYDROmorphine (DILAUDID) injection | 0.2 mg, intravenous, every 3 hours PRN, moderate pain (score 4-6), Post-op
Use if patient is unable to swallow or faster onset is needed |

PRN Oral for Severe Pain (Pain Score 7-10): For Patients LESS than 65 years old (Single Response)

(adjust dose for renal/liver function and age)

- | | |
|--|--|
| <input type="checkbox"/> HYDROmorphine (DILAUDID) tablet | 2 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Give if patient is able to tolerate oral medication |
| <input type="checkbox"/> morphine (MSIR) tablet | 15 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Give if patient is able to tolerate oral medication |
| <input type="checkbox"/> oxyCODONE (ROXICODONE) immediate release tablet | 10 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op
Give if patient is able to tolerate oral medication |

PRN Oral for Severe Pain (Pain Score 7-10): For Patients GREATER than 65 years old (Single Response)
(adjust dose for renal/liver function and age)

<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO) 7.5-325 mg per tablet	1 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Give if patient is able to tolerate oral medication
<input type="checkbox"/> HYDROcodone-acetaminophen (NORCO 10-325) 10-325 mg per tablet	1 tablet, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Give if patient is able to tolerate oral medication
<input type="checkbox"/> HYDROmorphine (DILAUDID) tablet	2 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Give if patient is able to tolerate oral medication
<input type="checkbox"/> morphine (MSIR) tablet	15 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Give if patient is able to tolerate oral medication
<input type="checkbox"/> oxyCODONE (ROXICODONE) immediate release tablet	5 mg, oral, every 6 hours PRN, severe pain (score 7-10), Post-op Give if patient is able to tolerate oral medication

PRN IV for Severe Pain (Pain Score 7-10): For Patients LESS than 65 years old (Single Response)

If you select a PCA option you will not be allowed to also order IV PRN pain medications from this section.
(adjust dose for renal/liver function and age)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	50 mcg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed
<input type="checkbox"/> morphine injection	4 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed
<input type="checkbox"/> HYDROmorphine (DILAUDID) injection	0.8 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed

PRN IV for Severe Pain (Pain Score 7-10): For Patients GREATER than 65 years old (Single Response)

If you select a PCA option you will not be allowed to also order IV PRN pain medications from this section.
(adjust dose for renal/liver function and age)

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	25 mcg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed
<input type="checkbox"/> morphine injection	2 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed
<input type="checkbox"/> HYDROmorphine (DILAUDID) injection	0.5 mg, intravenous, every 3 hours PRN, severe pain (score 7-10), Post-op Use if patient is unable to swallow or faster onset is needed

PCA Medications - Opioid Naive (Single Response)

<input type="checkbox"/> morPHINE PCA 30 mg/30 mL	Nurse Loading Dose: Not Ordered PCA Dose: 1 mg Lockout Interval: Not Ordered Basal Rate: 0 mg/hr MAX (Four hour dose limit): 20 mg intravenous, continuous, Post-op Management of breakthrough pain. Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber. For breakthrough pain in patients ages 19-59 years old with normal renal function, may bolus {Bolus Dose:26657::"2"} mg every {Bolus Frequency:26659::"3"} hours as needed. If pain persists, may increase PCA demand dose by {PCA Dose:26660::"0.5"} mg ONCE. Adjust doses for age, renal function or other factors.
---	--

<p>() hydromorPHONE (DILAUDID) 15 mg/30 mL PCA</p>	<p>Nurse Loading Dose: Not Ordered
PCA Dose: 0.2 mg
Lockout: Not Ordered
Basal Rate: 0 mg/hr
MAX (Four hour dose limit): 3 mg intravenous, continuous, Post-op Management of breakthrough pain. Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber. For breakthrough pain in patients ages 19-59 years old with normal renal function, may bolus {Bolus Dose:26662::"0.2"} mg every {Bolus Frequency:26663::"3"} hours as needed. If pain persists, may increase PCA demand dose by {PCA Dose:26664::"0.1"} mg ONCE. Adjust doses for age, renal function or other factors.</p>
<p>() fentaNYL (SUBLIMAZE) 1500 mcg/30 mL PCA</p>	<p>Nurse Loading Dose: Not Ordered
PCA Dose: 10 mcg
Lockout: Not Ordered
Basal Rate: 0 mcg/hr
Four Hour Dose Limit: 150 mcg intravenous, continuous, Post-op Management of breakthrough pain. Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber. For breakthrough pain in patient 19-59 years old, may bolus {Bolus Dose:26656::"25"} mcg every {Bolus Frequency:26655::"2"} hours as needed. If pain persists, may increase PCA demand dose by {PCA Dose:26654::"10"} mcg ONCE. Adjust doses for age, renal function or other factors.</p>

Nausea

<p>[] ondansetron (ZOFTRAN) IV or Oral (Selection Required)</p>	<p>"Or" Linked Panel</p>
<p>[] ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet</p>	<p>4 mg, oral, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is able to tolerate oral medication.</p>
<p>[] ondansetron (ZOFTRAN) 4 mg/2 mL injection</p>	<p>4 mg, intravenous, every 8 hours PRN, nausea, vomiting, Post-op Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.</p>
<p>[] promethazine (PHENERGAN) IV or Oral or Rectal</p>	<p>"Or" Linked Panel</p>
<p>[] promethazine (PHENERGAN) 12.5 mg IV</p>	<p>12.5 mg, intravenous, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.</p>
<p>[] promethazine (PHENERGAN) tablet</p>	<p>12.5 mg, oral, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.</p>
<p>[] promethazine (PHENERGAN) suppository</p>	<p>12.5 mg, rectal, every 6 hours PRN, nausea, vomiting, Post-op Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.</p>

Bowel regimen (Single Response)

<p>() docusate sodium (COLACE) capsule</p>	<p>100 mg, oral, 2 times daily PRN, constipation, Post-op</p>
<p>() bisacodyl (DULCOLAX) suppository</p>	<p>10 mg, rectal, daily PRN, constipation, Post-op</p>
<p>() sennosides-docusate sodium (SENOKOT-S) 8.6-50 mg per tablet</p>	<p>1 tablet, oral, daily PRN, constipation, Post-op</p>

VTE

DVT Risk and Prophylaxis Tool (Single Response) (Selection Required)

URL: "\lappt1.pdf"

- () Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Single Response) (Selection Required)

Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required) **"And" Linked Panel**

Moderate risk of VTE Routine, Once, PACU & Post-op

Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis Routine, Once
No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:
PACU & Post-op

High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required) **"And" Linked Panel**

High risk of VTE Routine, Once, PACU & Post-op

Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis Routine, Once
No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:
PACU & Post-op

LOW Risk of DVT (Selection Required)

Low Risk Definition

Age less than 60 years and NO other VTE risk factors

Low Risk (Single Response) (Selection Required)

Low risk of VTE Routine, Once
Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation
PACU & Post-op

MODERATE Risk of DVT - Surgical (Selection Required)

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Moderate Risk (Selection Required)

Moderate risk of VTE Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis BUT order Sequential compression device **"And" Linked Panel**

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

Place/Maintain sequential compression device continuous Routine, Continuous, PACU & Post-op

Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis **"And" Linked Panel**

Contraindications exist for pharmacologic prophylaxis Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> MODERATE Risk of DVT - Non-Surgical (Selection Required)	
Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission	
<input type="checkbox"/> Moderate Risk (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	

<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection 40 mg	40 mg, subcutaneous, daily at 1700
<input type="checkbox"/> enoxaparin (LOVENOX) injection 30 mg	30 mg, subcutaneous, daily at 1700 if CrCL LESS than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection 2.5 mg	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> HEParin (porcine) injection 5,000 Units	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> HEParin (porcine) injection 5,000 Units	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy to dose warfarin	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Non-Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
[] High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
[] High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
[] High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
[] High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> Apixaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, 2 times daily, Starting S+1 Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min.

<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> Rivaroxaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL) Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

DVT Risk and Prophylaxis Tool (Single Response)

URL: "\appt1.pdf"

<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Single Response) (Selection Required)	
<input type="checkbox"/> Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	"And" Linked Panel
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Selection Required)	"And" Linked Panel
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following: PACU & Post-op
<input type="checkbox"/> LOW Risk of DVT (Selection Required)	

Low Risk Definition
Age less than 60 years and NO other VTE risk factors

Low Risk (Single Response) (Selection Required)

Low risk of VTE
Routine, Once
Low risk: Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation
PACU & Post-op

MODERATE Risk of DVT - Surgical (Selection Required)

Moderate Risk Definition

Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.

One or more of the following medical conditions:

CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome

Age 60 and above

Central line

History of DVT or family history of VTE

Anticipated length of stay GREATER than 48 hours

Less than fully and independently ambulatory

Estrogen therapy

Moderate or major surgery (not for cancer)

Major surgery within 3 months of admission

Moderate Risk (Selection Required)

Moderate risk of VTE
Routine, Once, PACU & Post-op

Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)

Contraindications exist for pharmacologic prophylaxis
BUT order Sequential compression device
"And" Linked Panel

Contraindications exist for pharmacologic prophylaxis
Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

Place/Maintain sequential compression device continuous
Routine, Continuous, PACU & Post-op

Contraindications exist for pharmacologic prophylaxis
AND mechanical prophylaxis
"And" Linked Panel

Contraindications exist for pharmacologic prophylaxis
Routine, Once
No pharmacologic VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

Contraindications exist for mechanical prophylaxis
Routine, Once
No mechanical VTE prophylaxis due to the following contraindication(s):
PACU & Post-op

enoxaparin (LOVENOX) injection (Single Response) (Selection Required)

enoxaparin (LOVENOX) syringe
40 mg, subcutaneous, daily at 0600, Starting S+1

patients with CrCL LESS than 30 mL/min
30 mg, subcutaneous, daily at 0600, Starting S+1
For Patients with CrCL LESS than 30 mL/min

patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min
30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1
For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min

patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min
40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1
For Patient weight of 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
[] Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> MODERATE Risk of DVT - Non-Surgical (Selection Required)	
Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated. One or more of the following medical conditions: CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome Age 60 and above Central line History of DVT or family history of VTE Anticipated length of stay GREATER than 48 hours Less than fully and independently ambulatory Estrogen therapy Moderate or major surgery (not for cancer) Major surgery within 3 months of admission	
[] Moderate Risk (Selection Required)	
<input type="checkbox"/> Moderate risk of VTE	Routine, Once, PACU & Post-op
[] Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Patient is currently receiving therapeutic anticoagulation	Routine, Once No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s):
<input type="checkbox"/> enoxaparin (LOVENOX) injection 40 mg	40 mg, subcutaneous, daily at 1700
<input type="checkbox"/> enoxaparin (LOVENOX) injection 30 mg	30 mg, subcutaneous, daily at 1700 if CrCL LESS than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection 2.5 mg	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> HEParin (porcine) injection 5,000 Units	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> HEParin (porcine) injection 5,000 Units	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy to dose warfarin	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800, Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1, PACU & Post-op If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):

<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM, PACU & Post-op
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM, PACU & Post-op Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1, PACU & Post-op Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Non-Surgical (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 1700, Starting S
<input type="checkbox"/> patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 1700, Starting S For Patients with CrCL LESS than 30 mL/min
<input type="checkbox"/> patients weight between 100-139 kg AND CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily, Starting S For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min
<input type="checkbox"/> patients weight 140 kg or GREATER AND CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily, Starting S For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min
<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min. This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.

<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op
<input type="checkbox"/> HIGH Risk of DVT - Surgical (Hip/Knee) (Selection Required)	
High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed. One or more of the following medical conditions: Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders) Severe fracture of hip, pelvis or leg Acute spinal cord injury with paresis Multiple major traumas Abdominal or pelvic surgery for CANCER Acute ischemic stroke History of PE	
<input type="checkbox"/> High Risk (Selection Required)	
<input type="checkbox"/> High risk of VTE	Routine, Once, PACU & Post-op
<input type="checkbox"/> High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for pharmacologic prophylaxis	Routine, Once No pharmacologic VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> aspirin chewable tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> aspirin (ECOTRIN) enteric coated tablet	162 mg, oral, daily, Starting S+1
<input type="checkbox"/> Apixaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> apixaban (ELIQUIS) tablet	2.5 mg, oral, 2 times daily, Starting S+1 Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor apixaban (ELIQUIS) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> enoxaparin (LOVENOX) injection (Single Response) (Selection Required)	
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	40 mg, subcutaneous, daily at 0600, Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients with CrCL LESS than 30 mL/min	30 mg, subcutaneous, daily at 0600, Starting S+1 For Patients with CrCL LESS than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min	30 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight between 100-139 kg and CrCl GREATER than 30 mL/min.
<input type="checkbox"/> enoxaparin (LOVENOX) syringe - For Patients weight between 140 kg or GREATER and CrCl GREATER than 30 mL/min	40 mg, subcutaneous, 2 times daily at 0600, 1800 (TIME CRITICAL), Starting S+1 For Patients weight 140 kg or GREATER and CrCl GREATER than 30 mL/min

<input type="checkbox"/> fondaparinux (ARIXTRA) injection	2.5 mg, subcutaneous, daily, Starting S+1 If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min This patient has a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT):
<input type="checkbox"/> heparin (porcine) injection	5,000 Units, subcutaneous, every 8 hours, S+1 at 6:00 AM
<input type="checkbox"/> heparin (porcine) injection (Recommended for patients with high risk of bleeding, e.g. weight < 50kg and age > 75yrs)	5,000 Units, subcutaneous, every 12 hours, S+1 at 6:00 AM Recommended for patients with high risk of bleeding, e.g. weight LESS than 50kg and age GREATER than 75yrs.
<input type="checkbox"/> Rivaroxaban and Pharmacy Consult (Selection Required)	
<input type="checkbox"/> rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission	10 mg, oral, daily at 0600 (TIME CRITICAL) Indications: VTE prophylaxis
<input type="checkbox"/> Pharmacy consult to monitor rivaroxaban (XARELTO) therapy	STAT, Until discontinued, Starting S Indications: VTE prophylaxis
<input type="checkbox"/> warfarin (COUMADIN) tablet	oral, daily at 1700, Starting S+1 Indication:
<input type="checkbox"/> Pharmacy consult to manage warfarin (COUMADIN)	STAT, Until discontinued, Starting S Indication:
<input type="checkbox"/> Mechanical Prophylaxis (Single Response) (Selection Required)	
<input type="checkbox"/> Contraindications exist for mechanical prophylaxis	Routine, Once No mechanical VTE prophylaxis due to the following contraindication(s): PACU & Post-op
<input type="checkbox"/> Place/Maintain sequential compression device continuous	Routine, Continuous, PACU & Post-op

Labs

Laboratory Today

<input type="checkbox"/> CBC with platelet and differential	Once, Post-op
<input type="checkbox"/> Basic metabolic panel	Once, Post-op
<input type="checkbox"/> Prothrombin time with INR	Once, Post-op
<input type="checkbox"/> Partial thromboplastin time	Once, Post-op
<input type="checkbox"/> Magnesium level	Once, Post-op

Laboratory Tomorrow

<input type="checkbox"/> CBC with platelet and differential	AM draw For 1 Occurrences, Post-op
<input type="checkbox"/> Basic metabolic panel	AM draw For 1 Occurrences, Post-op
<input type="checkbox"/> Prothrombin time with INR	AM draw For 1 Occurrences, Post-op
<input type="checkbox"/> Partial thromboplastin time	AM draw For 1 Occurrences, Post-op
<input type="checkbox"/> Magnesium level	AM draw For 1 Occurrences, Post-op

Cardiology

Imaging

HM IP XRAY VASCULAR FL

<input type="checkbox"/> Chest 1 Vw Portable	STAT, 1 time imaging For 1 , Post-op
<input type="checkbox"/> Abdomen 1 Vw Portable	STAT, 1 time imaging For 1 , Post-op
<input type="checkbox"/> XR Chest 1 Vw Portable - AM	Routine, 1 time imaging, Starting S+1 at 6:00 AM For 1 , Post-op
<input type="checkbox"/> XR Abdomen 1 Vw Portable - AM	Routine, 1 time imaging, Starting S+1 at 6:00 AM For 1 , Post-op

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Ancillary Consults

<input type="checkbox"/> Consult to Case Management	Reason for Consult? Discharge planning Post-op
<input type="checkbox"/> Consult to Social Work	Reason for Consult? Discharge planning Post-op
<input checked="" type="checkbox"/> Consult PT eval and treat	Reasons for referral to Physical Therapy (mark all applicable): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal): Weight Bearing Status: Post-op
<input type="checkbox"/> Consult PT wound care	Special Instructions: Location of Wound? Post-op
<input checked="" type="checkbox"/> Consult OT eval and treat	Reason for referral to Occupational Therapy (mark all that apply): Are there any restrictions for positioning or mobility? Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal): Weight Bearing Status: Post-op
<input type="checkbox"/> Consult to Nutrition Services	Reason For Consult? Purpose/Topic: Post-op
<input type="checkbox"/> Consult to Wound Ostomy Care nurse	Reason for consult: Reason for consult: Reason for consult: Reason for consult: Consult for NPWT: Reason for consult: Reason for consult: Post-op
<input type="checkbox"/> Consult Intensive Care	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated? Post-op
<input type="checkbox"/> Consult Nephrology	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated? Post-op
<input type="checkbox"/> Consult Neurology	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated? Post-op
<input type="checkbox"/> Consult Diabetes/Endocrinology	Reason for Consult? Patient/Clinical information communicated? Patient/clinical information communicated? Post-op

Additional Orders