

Treprostinil (Remodulin) IV or SQ Infusion Orders [4644]

A calculator is available at formweb.com/houstonmethodist under Drug Information and Clinical Calculators to assist with epoprostenol or treprostinil calculations. To navigate to this site, please open an internet browser and use the web address noted previously.

Nursing

Nursing

<input type="checkbox"/> Treprostinil (REMODULIN) instructions for Nursing:	Routine, Until discontinued, Starting S
	1. Do NOT infuse any other IV fluids or medications through treprostinil (Remodulin) line.
	2. Do NOT draw blood from a lumen adjacent to treprostinil (Remodulin) line.
	3. Do NOT STOP treprostinil (Remodulin) for ANY REASON without first contacting MD who ordered treprostinil (Remodulin)
	4. DO NOT STOP treprostinil (Remodulin) for a CODE BLUE without first contacting MD who ordered treprostinil (Remodulin)
	5. Ensure back-up back-up infusion pump are readily available Document any infusion rate changes at appropriate times in the MAR

Medications

Medications (Single Response) (Selection Required)

<input type="checkbox"/> Continuation of Home Therapy	
<input type="checkbox"/> Vital Signs	"And" Linked Panel
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol Upon initiation of therapy or dose increase, every 15 minutes x 4, then every 30 minutes x2, then every 4 hours.
<input type="checkbox"/> Infusion Options (Single Response)	
<input type="checkbox"/> Intravenous Infusion - CADD-Legacy Pump (mg/100mL)	intravenous, every 48 hours at 1400, S at 2:00 PM Do NOT STOP treprostinil (Remodulin) for ANY REASON without first contacting MD who ordered treprostinil (Remodulin) For CENTAL LINE administration Infusion Pump Rate (mL per 24 hours): Dosing Weight: (in kg): Dose: (in ng/kg/min):
<input type="checkbox"/> SQ/IV Infusion - Crono 5 Syringe (mg/20mL)	every 48 hours at 1400, S at 2:00 PM Do NOT STOP treprostinil (Remodulin) for ANY REASON without first contacting MD who ordered treprostinil (Remodulin) For CENTAL LINE subcutaneous administration Infusion Pump Rate (mcl per hour): Dosing Weight: (in kg): Dose: (in ng/kg/min):
<input type="checkbox"/> SQ Infusion - CADD MS3 pump (3mL)	3 mL, subcutaneous, every 72 hours at 1400, S at 2:00 PM Do NOT STOP treprostinil (Remodulin) for ANY REASON without first contacting MD who ordered treprostinil (Remodulin) For CENTAL LINE subcutaneous administration Infusion Pump Rate (mL/hr): Dosing Weight: (in kg): Dose: (in ng/kg/min):
<input type="checkbox"/> Pharmacy Consult	
<input type="checkbox"/> Pharmacy consult to review pulmonary arterial hypertension (PAH) infusions	STAT, Until discontinued, Starting S
<input type="checkbox"/> Therapy Titration	
<input type="checkbox"/> Vital Signs	"And" Linked Panel
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Per unit protocol Upon initiation of therapy or dose increase, every 15 minutes x 4, then every 30 minutes x2, then every 4 hours.

☐ Infusion Options (Single Response)

- | | |
|---|--|
| <p>() Intravenous Infusion - CADD-Legacy Pump (mg/100mL)</p> | <p>intravenous, every 48 hours at 1400
Do NOT STOP treprostinil (Remodulin) for ANY REASON without first contacting MD who ordered treprostinil (Remodulin)
For CENTAL LINE administration.</p> <p>Titration Instructions:
Initiate treprostinil (Remodulin) as ordered.
{Increase/Decrease:34622} by {.:34620} mL/24hours every {.:34621} hours
{Maximum/Minimum:34623} infusion rate of ***mL/24hours
Infusion Pump Rate (mL per 24 hours):
Dosing Weight: (in kg):
Dose: (in ng/kg/min):</p> |
| <p>() SQ/IV Infusion - Crono 5 Syringe (mg/20mL)</p> | <p>every 48 hours at 1400
Do NOT STOP treprostinil (Remodulin) for ANY REASON without first contacting MD who ordered treprostinil (Remodulin)
For CENTAL LINE administration.</p> <p>Titration Instructions:
Initiate treprostinil (Remodulin) as ordered.
{Increase/Decrease:34622} by (***)mL/hr every (**) hours
{Maximum/Minimum:34623} infusion rate of *** mL/hr
Infusion Pump Rate (mcl per hour):
Dosing Weight: (in kg):
Dose: (in ng/kg/min):</p> |
| <p>() SQ Infusion - CADD MS3 pump (3mL)</p> | <p>3 mL, subcutaneous, every 72 hours at 1400
Do NOT STOP treprostinil (Remodulin) for ANY REASON without first contacting MD who ordered treprostinil (Remodulin) For CENTAL LINE subcutaneous administration</p> <p>Titration Instructions:
Initiate treprostinil (Remodulin) as ordered.
{Increase/Decrease:34622} by (***)mL/hr every (**) hours
{Maximum/Minimum:34623} infusion rate of *** mL/hr
Infusion Pump Rate (mL/hr):
Dosing Weight: (in kg):
Dose: (in ng/kg/min):</p> |

☐ Pharmacy Consult

- | | |
|--|---|
| <p><input type="checkbox"/> Pharmacy consult to review pulmonary arterial hypertension (PAH) infusions</p> | <p>STAT, Until discontinued, Starting S</p> |
|--|---|