

Amphotericin B LIPOSOMAL (AMBISOME) Infusion (RESTRICTED) [2389]

!!! LIPOSOMAL formulation !!!

Amphotericin B Liposomal (AMBISOME) infusion is RESTRICTED to Infectious Diseases (ID), Solid Organ Transplant (SOT), Bone Marrow Transplant (BMT), and Hematology/Oncology (Heme/Onc) specialists.

Medications

amphotericin B (AMBISOME) with fluids +/- pre-medications (Single Response) (Selection Required)

() amphotericin with pre-medications & fluids	"And" Linked Panel
<input type="checkbox"/> sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours, Starting H+2 Hours Give 1 hour BEFORE amphotericin
<input type="checkbox"/> acetaminophen (TYLENOL) tablet	650 mg, oral, every 24 hours, Starting H+150 Minutes Give 30 minute BEFORE amphotericin
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection	25 mg, intravenous, every 24 hours, Starting H+150 Minutes Give 30 minute BEFORE amphotericin
<input type="checkbox"/> hydrocortisone sodium succinate (Solu-CORTEF) injection	25 mg, intravenous, every 24 hours, Starting H+150 Minutes Give 30 minute BEFORE amphotericin
<input type="checkbox"/> dextrose 5% flush syringe	10 mL, intravenous, every 24 hours, Starting H+3 Hours Flush line immediately BEFORE amphotericin.
<input type="checkbox"/> amphotericin B (liposomal) (AmBisome) infusion (RESTRICTED)	intravenous, for 2 Hours, every 24 hours, Starting H+3 Hours RESTRICTED to Infectious Diseases (ID), Solid Organ Transplant (SOT), Bone Marrow Transplant (BMT), and Hematology/Oncology (Heme/Onc) specialists. Are you an ID, SOT, BMT, or Heme/Onc specialist or ordering on behalf of one? Reason for Therapy:
<input type="checkbox"/> dextrose 5% flush syringe	10 mL, intravenous, every 24 hours, Starting H+5 Hours Flush line immediately AFTER amphotericin.
<input type="checkbox"/> sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours, Starting H+5 Hours Give AFTER amphotericin and dextrose 5% flush.
() amphotericin with fluids	"And" Linked Panel
<input type="checkbox"/> sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours, Starting H+2 Hours Give 1 hour BEFORE amphotericin
<input type="checkbox"/> dextrose 5% flush syringe	10 mL, intravenous, every 24 hours, Starting H+3 Hours Flush line immediately BEFORE amphotericin.
<input type="checkbox"/> amphotericin B (liposomal) (AmBisome) infusion (RESTRICTED)	intravenous, for 2 Hours, every 24 hours, Starting H+3 Hours RESTRICTED to Infectious Diseases (ID), Solid Organ Transplant (SOT), Bone Marrow Transplant (BMT), and Hematology/Oncology (Heme/Onc) specialists. Are you an ID, SOT, BMT, or Heme/Onc specialist or ordering on behalf of one? Reason for Therapy:
<input type="checkbox"/> dextrose 5% flush syringe	10 mL, intravenous, every 24 hours, Starting H+5 Hours Flush line immediately AFTER amphotericin.
<input type="checkbox"/> sodium chloride 0.9 % bolus	500 mL, intravenous, for 60 Minutes, every 24 hours, Starting H+5 Hours Give AFTER amphotericin and dextrose 5% flush.

Labs

Laboratory

<input type="checkbox"/> Basic metabolic panel - STAT	STAT For 1 Occurrences
<input type="checkbox"/> Basic metabolic panel	Once
<input type="checkbox"/> Magnesium level	Once
<input type="checkbox"/> Phosphorus level	Once