

Bivalirudin Infusion for Heparin Induced Thrombocytopenia [3284]

General

Nursing

Nursing

[X] Weigh patient	STAT, Once For 1 Occurrences if baseline measured weight not in medical record. Notify pharmacist if patient cannot be weighed.
[X] Notify Pharmacist	Routine, Until discontinued, Starting S For Until specified, NOTIFY pharmacist if patient CANNOT be weighed.
[X] Bivalirudin instructions	Routine, Until discontinued, Starting S Do not interrupt bivalirudin infusion unless ordered. Contact pharmacist regarding compatibility with other IV drugs if access is a concern. Contact pharmacist immediately if bivalirudin infusion is stopped for any reason.
[X] Notify pharmacist	Routine, Until discontinued, Starting S, If a physician writes bivalirudin orders. Bivalirudin dose or infusion changes to be made by pharmacist only. Immediately if patient transfers or has an off-unit procedure. If compatibility with other IV drugs or if IV access is a concern IMMEDIATELY if bivalirudin infusion is stopped for any reason.
[X] Notify Physician and Pharmacist	Routine, Until discontinued, Starting S, If hematoma, bleeding or suspected bleeding occurs.
[X] Draw blood for PTT from arm that doesn't have bivalirudin infusion.	Routine, Until discontinued, Starting S If there is no other access, other than the bivalirudin line, then stop the bivalirudin, flush the line, aspirate 10 ml of blood to waste, and then reflush the line prior to drawing a specimen.
[X] Do not call PTT results to physician unless subsequently ordered.	Routine, Until discontinued, Starting S For Until specified Do not call PTT results to physician unless subsequently ordered.

IV Fluids

Medications

bivalirudin (ANGIOMAX) infusion

[X] bivalirudin (ANGIOMAX) 1 mg/mL in sodium chloride 0.9 % 250 mL infusion	0.1 mg/kg/hr, intravenous, continuous Indication: HIT Therapeutic Monitoring Target: PTT – 50 to 90 secs
-----------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------

VTE

Labs

Lab - Initiation

[X] Anti Xa, unfractionated	STAT For 1 Occurrences
[X] Prothrombin time with INR	STAT For 1 Occurrences
[X] Partial thromboplastin time	STAT For 1 Occurrences
[] CBC with platelet and differential	STAT For 1 Occurrences
[] Basic metabolic panel	STAT For 1 Occurrences

Lab - Continuing

[] CBC with platelet and differential	AM draw For 1 Occurrences while on bivalirudin therapy
----------------------------------------	-----------------------------------------------------------

<input type="checkbox"/> Partial thromboplastin time	Once For Until specified Three (3) hours after initiation of bivalirudin infusion. Do not draw blood from the arm that has bivalirudin infusion. If there is no other access, other than the bivalirudin line for a Partial thromboplastin time (PTT) specimen draw, then stop the bivalirudin, flush the line, and aspirate 10 ml of blood to waste prior to drawing a specimen.
<input type="checkbox"/> Occult blood, stool	Conditional Frequency, Stool guiac stools when there is a hemoglobin decrease of more than 2 gm/dL in a 24 hour period

Cardiology

Imaging

Other Studies

Respiratory

Rehab

Consults

For Physician Consult orders use sidebar

Additional Orders