

Midline Catheter Maintenance [3400]

For Midline

Nursing Care (NOT HMH HMSNF HMWB HMSJ HMW)

[X] Change dressing: Midline	Routine, Once Change every 7 days and as needed; If sponge/gauze applied under dressing, then change dressing every 48 hrs. Only reapply gauze dressing if site is actively draining.
[X] Needleless Connector	Routine, Weekly Replace needleless connector with each IV blood draw, IV tubing change, and as needed
[X] Replace Stabilization Device every 7 days	Routine, Until discontinued, Starting S
[] Measure and Record - Arm Circumference Every 8 hours	Routine, Now then every 8 hours
[X] Notify Physician - (Midline)	Routine, Until discontinued, Starting S, If arm swelling, drainage, increase in arm circumference, or other complications arise concerning the Midline
[X] Notify Vascular Access Team - (Midline)	Routine, Once If arm circumference increases
[] Ok to use (Midline)	Routine, Until discontinued, Starting S Device: Midline

Nursing Care (HMH HMWB HMSJ & HMW ONLY)

[X] Change dressing: Midline	Routine, Once Change every 7 days and as needed; If sponge/gauze applied under dressing, then change dressing every 48 hrs. Only reapply gauze dressing if site is actively draining.
[] Midline has CT Scan capability- Contrast media via lumens designated for power contrast only.	Routine, Until discontinued, Starting S
[X] Needleless Connector	Routine, Weekly Replace needleless connector with each IV blood draw, IV tubing change, and as needed
[X] Replace Stabilization Device every 7 days	Routine, Until discontinued, Starting S
[] Measure and Record - Arm Circumference Every 8 hours	Routine, Now then every 8 hours
[X] Notify Physician - (Midline)	Routine, Until discontinued, Starting S, If arm swelling, drainage, increase in arm circumference, or other complications arise concerning the Midline
[X] Notify Vascular Access Team - (Midline)	Routine, Once If arm circumference increases
[] Ok to use (Midline)	Routine, Until discontinued, Starting S Device: Midline

Notify MD

Notify MD

[X] Notify Physician if arm swelling, drainage, or other complications arise concerning the midline	Routine, Until discontinued, Starting S
---	---

Medications

IV Flush

[X] sodium chloride 0.9 % flush	10 mL, intravenous, every 8 hours Aspirate and flush every 8 hours for saline locked IV. ONLY USE 10 mL SYRINGE OR LARGER TO FLUSH LINE.
[X] sodium chloride 0.9 % flush: 10 mL	10 mL, intravenous, PRN, line care Aspirate and flush 10 mL IV push before and after medications and as needed using turbulent method. ONLY USE 10 mL SYRINGE OR LARGER TO FLUSH LINE.

[X] sodium chloride 0.9 % flush: 20 mL

20 mL, intravenous, PRN, line care
Flush with 20 mLs IV push after blood withdrawals. ONLY
USE 10 mL SYRINGE OR LARGER TO FLUSH LINE.

Medications - NOT HMSTC, HMH

[] lidocaine (XYLOCAINE) 1% injection

5 mL, intradermal, once PRN, Midline insertion

Medications - HMH Only

[] lidocaine PF (XYLOCAINE) 1 % injection

50 mg, intradermal, once PRN, Midline insertion

Medications - HMSTC Only

[] lidocaine (XYLOCAINE) 1% injection

5 mL, intradermal, once PRN, Midline insertion