

**General**

**Nursing**

**Consent**

<input type="checkbox"/> Complete consent for	Routine, Once Procedure: Diagnosis/Condition: Physician: Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate? Pre-Procedure
---	--

**Vital Signs**

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Once For 1 Occurrences Upon arrival to Interventional Radiology unit., Pre-Procedure
<input type="checkbox"/> Pulse oximetry	Routine, Once For 1 Occurrences Current FIO2 or Room Air: Upon arrival to Interventional Radiology unit., Pre-Procedure

**Nursing**

<input type="checkbox"/> Hold AM meds	Routine, Once For 1 Occurrences, Pre-Procedure
<input type="checkbox"/> Insert peripheral IV	Routine, Once Use 20-22 gauge Jelco catheter, Pre-Procedure
<input type="checkbox"/> Insert and maintain Foley	
<input type="checkbox"/> Insert Foley catheter	Routine, Once Type: Size: Urinometer needed: Pre-Procedure
<input type="checkbox"/> Foley Catheter Care	Routine, Until discontinued, Starting S Orders: Maintain Pre-Procedure
<input type="checkbox"/> Bedside glucose	Routine, Once If diabetic, upon arrival to Interventional Radiology/AOD unit. Notify Radiologist if glucose level less than 60 or greater than 200., Pre-Procedure

**Diet**

<input type="checkbox"/> NPO after midnight	Diet effective midnight, Starting S+1 at 12:01 AM NPO: Except meds Pre-Operative fasting options: Pre-Procedure
---	--

**Medications**

**Buffered Lidocaine 1% Injection**

<input type="checkbox"/> lidocaine 1% buffered with 8.4% sodium bicarbonate injection	10 mL, injection, once, For 1 Doses, Pre-Procedure
---	--

**Prophylaxis Recommendation for Patients with an Allergy to Iodine Contrast Media (Single Response)**

Medications may be administered by oral or IV route. The equivalent of these medications may be administered as determined by the attending or referring physician.

In a true medical emergency situation the exam should proceed with an accelerated premedication schedule at the direction of the attending physician. (attachment to Radiology Policy R 711)

<input type="checkbox"/> Oral methylPREDNISolone (MEDROL) tablet	32 mg, oral, once, For 1 Doses, Pre-Procedure For iodine allergy prophylaxis. Give 6-12 hours prior to the examination.
<input type="checkbox"/> IV methylPREDNISolone succinate (Solu-MEDROL) 40 mg/mL injection	40 mg, intravenous, for 5 Minutes, once, For 1 Doses, Pre-Procedure For iodine allergy prophylaxis. Give 2 hours prior to the examination.

#### For patients with allergic reaction

<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection	50 mg, intravenous, once, For 1 Doses, Pre-Procedure Given in Interventional Radiology unit
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection	125 mg, intravenous, once, For 1 Doses, Pre-Procedure Given in Interventional Radiology unit

## VTE

## Labs

### COVID-19 Qualitative PCR

<input type="checkbox"/> COVID-19 qualitative RT-PCR - Nasal Swab	STAT For 1 Occurrences Specimen Source: Nasal Swab Is this for pre-procedure or non-PUI assessment? Yes Pre-Procedure
---	--

### Labs

<input type="checkbox"/> CBC and differential	Once, Starting S, Pre-Procedure
<input type="checkbox"/> Comprehensive metabolic panel	Once, Pre-Procedure
<input type="checkbox"/> Basic metabolic panel	Once, Starting S, Pre-Procedure
<input type="checkbox"/> Protime-INR	Once, Starting S, Pre-Procedure
<input type="checkbox"/> Platelet count	Once, Pre-Procedure
<input type="checkbox"/> Partial thromboplastin time	Once, Starting S, Pre-Procedure
<input type="checkbox"/> hCG qualitative, serum	Once, Pre-Procedure

## Cardiology

## Imaging

## Other Studies

## Respiratory

## Rehab

## Consults

For Physician Consult orders use sidebar

## Additional Orders