

# CVIR Post Port/Central Line/Central Venous Catheter Placement [1408]

## General

## Nursing

### Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 15 min Every 15 min times 2, PACU & Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 15 min Every 15 min times 4, PACU & Post-op
<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Every 15 min Every 15 min times 4, then every 30 min times 4, then every 1 hour times 2, then every 4 hours., PACU & Post-op

### Activity

<input type="checkbox"/> Bed rest	Routine, Until discontinued, Starting S Bathroom Privileges: for 30 min, PACU & Post-op
<input type="checkbox"/> Bed rest	Routine, Until discontinued, Starting S For 1 Hours Bathroom Privileges: PACU & Post-op
<input type="checkbox"/> Bed rest	Routine, Until discontinued, Starting S For 2 Hours Bathroom Privileges: PACU & Post-op

### Nursing

<input type="checkbox"/> Assess operative site	Routine, With vitals Assess puncture/surgical site with vital signs, PACU & Post-op
<input type="checkbox"/> Change dressing	Routine, Every 72 hours For central venous catheter. Change dressing and needleless connector every Tuesday and Friday or as needed., PACU & Post-op
<input type="checkbox"/> Head of bed 45 degrees	Routine, Until discontinued, Starting S Head of bed: 45 degrees PACU & Post-op
<input type="checkbox"/> Ok to use - portacath	Routine, Until discontinued, Starting S Device: Portacath Port has been flushed with Heparin: aspirate before use, PACU & Post-op
<input type="checkbox"/> Ok to use - central venous catheter	Routine, Until discontinued, Starting S Device: Other Other: Central venous catheter PACU & Post-op
<input type="checkbox"/> Ok to use - dialysis access	Routine, Until discontinued, Starting S Device: Dialysis Access PACU & Post-op
<input type="checkbox"/> Ok to use - tunneled central venous cath	Routine, Until discontinued, Starting S Device: Other Other: tunneled central venous cath PACU & Post-op
<input type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, every 8 hours, PACU & Post-op Flush each lumen with 10 milliliters normal saline after each use or every 8 hours.
<input type="checkbox"/> sodium chloride 0.9 % flush	10 mL, intravenous, PRN, line care, PACU & Post-op Flush after every use, flush each lumen with 10 mL normal saline

<input type="checkbox"/> Nursing communication	Routine, Until discontinued, Starting S Instructions: ***, PACU & Post-op
<input type="checkbox"/> Nursing communication	Routine, Until discontinued, Starting S Instructions: ***, PACU & Post-op

### Diet

<input type="checkbox"/> NPO except meds	Diet effective now, Starting S For 1 Hours NPO: Except meds Pre-Operative fasting options: PACU & Post-op
<input type="checkbox"/> Diet	Diet effective now, Starting S Diet(s): Other Options: Advance Diet as Tolerated? IDDSI Liquid Consistency: Fluid Restriction: Foods to Avoid: Foods to Avoid: PACU & Post-op
<input type="checkbox"/> Tube feeding	Diet effective now, Starting S Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Formula: Tube Feeding Schedule: Tube Feeding Schedule: Dietitian to manage Tube Feed? PACU & Post-op

### Notify

<input type="checkbox"/> Notify Interventional Radiology of worsening pain or bleeding from cath site.	Routine, Until discontinued, Starting S, Please call IR at your campus., PACU & Post-op
--	---

## IV Fluids

## Return to Floor

### Return to Floor (Single Response)

<input type="checkbox"/> OK to return to previous IP bed	Routine, Until discontinued, Starting S, PACU
<input type="checkbox"/> Discharge to home	Routine, Once Discharge Criteria: Clearing specialty: PACU

## Medications

### Post Procedure Medications

<input type="checkbox"/> fentaNYL (SUBLIMAZE) injection	25 mcg, intravenous, every 15 min PRN, severe pain (score 7-10), persistent pain, For 2 Doses, PACU & Post-op Hold for respirations LESS than 12 breath/minute. For persistent pain, call interventional MD.
<input type="checkbox"/> ondansetron (ZOFTRAN) IV	4 mg, intravenous, once PRN, nausea, vomiting, PACU & Post-op

## VTE

Labs

Cardiology

Diagnostic Imaging

Other Diagnostic Studies

Respiratory

Rehab

Consults

Additional Orders