

Nursing

IV Fluids

IV Fluids

<input type="checkbox"/> heparin 1000 unit/500 mL 0.9% sodium chloride	100 mL, intravenous, once in imaging, Flush
<input type="checkbox"/> HEParin 2000 Units in 1000 mL 0.9% NaCl infusion	100 mL, once in imaging, Flush
<input type="checkbox"/> sodium chloride 0.9 % bolus	50 mL, intravenous, for 1 Minutes, once in imaging, Flush
<input type="checkbox"/> sodium chloride 0.9% infusion	250 mL, intravenous, once in imaging

Medications

IR Contrasts

<input type="checkbox"/> iohexol (OMNIPAQUE) 180 mg iodine/mL injection	intrathecal, once in imaging, contrast
<input type="checkbox"/> iohexol (OMNIPAQUE) 240 mg iodine/mL injection	intravenous, once in imaging, contrast
<input type="checkbox"/> iohexol (OMNIPAQUE) 300 mg iodine/mL injection	intravenous, once in imaging, contrast
<input type="checkbox"/> iohexol (OMNIPAQUE) 350 mg iodine/mL injection	intravenous, once in imaging, contrast
<input type="checkbox"/> iodixanol (VISIPAQUE) 270 mg iodine/mL injection	50 mL, intravenous, once in imaging, contrast
<input type="checkbox"/> iodixanol (VISIPAQUE) 320 mg iodine/mL injection	intravenous, once in imaging, contrast
<input type="checkbox"/> carbon dioxide (CO2) gas	intra-arterial, once in imaging, contrast

XR Arthrograms - HMM, HMSJ, HMSTC Only

<input type="checkbox"/> gadobutrol (GADAVIST) 7.5 mL vial solution	7.5 mL, intravenous, once in imaging, contrast, xr arthrogram
<input type="checkbox"/> gadopentetate dimeglumine (MAGNEVIST) 5 mL vial solution	5 mL, intra-articular, once in imaging, contrast, xr arthrogram
<input type="checkbox"/> iohexol (OMNIPAQUE 180)	20 mL, intra-arterial, once in imaging, contrast Intra-articular route
<input type="checkbox"/> iohexol (OMNIPAQUE 300)	30 mL, Other, once in imaging, contrast, xr arthrogram Intra-articular route
<input type="checkbox"/> lidocaine 1% buffered with 8.4% sodium bicarbonate injection	5 mL, Other, once in imaging, xr arthrogram Intra-articular route

XR Arthrograms - HMW Only

<input type="checkbox"/> gadopentetate dimeglumine (MAGNEVIST) 5 mL vial solution	5 mL, intra-articular, once in imaging, contrast, xr arthrogram
<input type="checkbox"/> gadobutrol (GADAVIST) 7.5 mL vial solution	7.5 mL, intravenous, once in imaging, contrast
<input type="checkbox"/> iohexol (OMNIPAQUE 180)	20 mL, intra-arterial, once in imaging, contrast Intra-articular route
<input type="checkbox"/> iohexol (OMNIPAQUE 300)	30 mL, Other, once in imaging, contrast, xr arthrogram Intra-articular route
<input type="checkbox"/> lidocaine 1% buffered with 8.4% sodium bicarbonate injection	5 mL, Other, once in imaging, xr arthrogram Intra-articular route

XR Arthrograms - HMSL Only

<input type="checkbox"/> gadopentetate dimeglumine (MAGNEVIST) 5 mL vial solution	5 mL, intra-articular, once in imaging, contrast, xr arthrogram
<input type="checkbox"/> iohexol (OMNIPAQUE 180)	20 mL, intra-arterial, once in imaging, contrast Intra-articular route
<input type="checkbox"/> iohexol (OMNIPAQUE 300)	30 mL, Other, once in imaging, contrast, xr arthrogram Intra-articular route
<input type="checkbox"/> lidocaine 1% buffered with 8.4% sodium bicarbonate injection	5 mL, Other, once in imaging, xr arthrogram Intra-articular route
<input type="checkbox"/> lidocaine PF (XYLOCAINE) 10 mg/mL (1 %) injection	5 mL, Other, once in imaging, xr arthrogram Intra-articular route
<input type="checkbox"/> gadobutrol (GADAVIST) vial solution	7.5 mL, intravenous, once in imaging, contrast, xr arthrogram

<input type="checkbox"/> methylPREDNISolone acetate (DEPO-MEDROL) injection	40 mg, intra-articular, once in imaging, xr arthrogram
<input type="checkbox"/> bupivacaine (PF) (MARCAINE) 0.25 % (2.5 mg/mL) solution	injection, once in imaging

XR Arthrograms - HMWB Only

<input type="checkbox"/> gadopentetate dimeglumine (MAGNEVIST) 5 mL vial solution	5 mL, intra-articular, once in imaging, contrast, xr arthrogram
<input type="checkbox"/> iohexol (OMNIPAQUE 180)	20 mL, intra-arterial, once in imaging, contrast Intra-articular route
<input type="checkbox"/> iohexol (OMNIPAQUE 300)	30 mL, Other, once in imaging, contrast, xr arthrogram Intra-articular route
<input type="checkbox"/> lidocaine 2% buffered with 8.4% sodium bicarbonate injection	5 mL, Other, once in imaging, xr arthrogram Intra-articular route

XR Arthrograms - HMSTJ Only

<input type="checkbox"/> gadopentetate dimeglumine (MAGNEVIST) 5 mL vial solution	5 mL, intra-articular, once in imaging, contrast, xr arthrogram
<input type="checkbox"/> gadobutrol (GADAVIST) 10 mL vial solution	10 mL, intra-articular, once in imaging, contrast, xr arthrogram
<input type="checkbox"/> iohexol (OMNIPAQUE 180)	20 mL, intra-arterial, once in imaging, contrast Intra-articular route
<input type="checkbox"/> iohexol (OMNIPAQUE 300)	30 mL, Other, once in imaging, contrast, xr arthrogram Intra-articular route
<input type="checkbox"/> lidocaine 1% buffered with 8.4% sodium bicarbonate injection	5 mL, Other, once in imaging, xr arthrogram

Additional Orders