

General

Nursing

Consent

<input type="checkbox"/> Complete consent for	Routine, Once Procedure: Diagnosis/Condition: Physician: Risks, benefits, and alternatives (as outlined by the Texas Medical Disclosure Panel, as appears on Houston Methodist Medical/Surgical Consent forms) were discussed with patient/surrogate? Pre-Procedure
---	--

Vital Signs

<input type="checkbox"/> Vital signs - T/P/R/BP	Routine, Once For 1 Occurrences Upon arrival to Interventional Radiology unit., Pre-Procedure
<input type="checkbox"/> Pulse oximetry	Routine, Once For 1 Occurrences Current FIO2 or Room Air: Upon arrival to Interventional Radiology unit., Pre-Procedure

Nursing

<input type="checkbox"/> Hold AM meds	Routine, Once For 1 Occurrences, Pre-Procedure
<input type="checkbox"/> Insert peripheral IV	Routine, Once Use 20-22 gauge Jelco catheter, Pre-Procedure
<input type="checkbox"/> Insert and maintain Foley	
<input type="checkbox"/> Insert Foley catheter	Routine, Once Type: Size: Urinometer needed: Pre-Procedure
<input type="checkbox"/> Foley Catheter Care	Routine, Until discontinued, Starting S Orders: Maintain Pre-Procedure
<input type="checkbox"/> Bedside glucose	Routine, Once If diabetic, upon arrival to Interventional Radiology/AOD unit. Notify Radiologist if glucose level less than 60 or greater than 200., Pre-Procedure

Diet

<input type="checkbox"/> NPO after midnight	Diet effective midnight, Starting S+1 at 12:01 AM NPO: Except meds Pre-Operative fasting options: Pre-Procedure
---	--

Medications

Antibiotics (Single Response) (Selection Required)

- For Tunneled HD Catheter; Gastrostomy Tube Placement; Gastrojejunostomy Tube Placement; Port Placement (Single Response)
- Antibiotics: For Patients GREATER than 120 kg (Single Response)

<input type="checkbox"/> ceFAZolin (ANCEF) IV - For Patients GREATER than 120 kg	3 g, intravenous, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:
<input type="checkbox"/> vancomycin (VANCOCIN) IV - For PCN allergic	15 mg/kg, intravenous, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:
<input type="checkbox"/> Antibiotics: For Patients LESS than or EQUAL to 120 kg (Single Response)	
<input type="checkbox"/> ceFAZolin (ANCEF) IV - For Patients LESS than or EQUAL to 120 kg	2 g, intravenous, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:
<input type="checkbox"/> vancomycin (VANCOCIN) IV - For PCN allergic	15 mg/kg, intravenous, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:
<input type="checkbox"/> For Uterine Artery Embolization	
<input type="checkbox"/> Antibiotics: For Patients GREATER than 120 kg (Single Response)	
<input type="checkbox"/> ceFAZolin (ANCEF) IV	3 g, intravenous, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> clindamycin (CLEOCIN) IV AND gentamicin (GARAMYCIN) IV - For PCN allergic For Penicillin Allergic Patients	"And" Linked Panel
Recommended ONLY for patients with high risk for penicillin anaphylaxis that are culture isolate sensitive to Clindamycin	
<input type="checkbox"/> clindamycin (CLEOCIN) IV	900 mg, intravenous, for 30 Minutes, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> gentamicin (GARAMYCIN) IVPB	1.5 mg/kg, intravenous, for 30 Minutes, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> Antibiotics: For Patients LESS than or EQUAL to 120 kg (Single Response)	
<input type="checkbox"/> ceFAZolin (ANCEF) IV	2 g, intravenous, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> clindamycin (CLEOCIN) IV AND gentamicin (GARAMYCIN) IV - For PCN allergic For Penicillin Allergic Patients	"And" Linked Panel
Recommended ONLY for patients with high risk for penicillin anaphylaxis that are culture isolate sensitive to Clindamycin	
<input type="checkbox"/> clindamycin (CLEOCIN) IV	900 mg, intravenous, for 30 Minutes, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis

<input type="checkbox"/> gentamicin (GARAMYCIN) IVPB	1.5 mg/kg, intravenous, for 30 Minutes, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis
() For Biliary Drainage Catheter; Nephrostomy Tube Insertion; TIPS (Single Response)	
<input type="checkbox"/> cefTRIAxone (ROCEPHIN) IV	1 g, intravenous, for 30 Minutes, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis Surgical Prophylaxis:
<input type="checkbox"/> Aztreonam + Vancomycin - For PCN allergic	"And" Linked Panel
<input type="checkbox"/> aztreonam (AZACTAM) IV	2 g, intravenous, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis
<input type="checkbox"/> vancomycin (VANCOGIN) IV	15 mg/kg, intravenous, once, For 1 Doses, Pre-Procedure On call to the operating room. Administer one hour prior to the opening incision. Reason for Therapy: Surgical Prophylaxis

lidocaine (XYLOCAINE) 1% injection

<input type="checkbox"/> lidocaine (XYLOCAINE) 10 mg/mL (1 %) injection	injection, once, For 1 Doses, Pre-Procedure
---	---

Buffered Lidocaine 1% Injection

<input type="checkbox"/> lidocaine 1% buffered with 8.4% sodium bicarbonate injection	10 mL, injection, once, For 1 Doses, Pre-Procedure
---	--

Prophylaxis Recommendation for Patients with an Allergy to Iodine Contrast Media (Single Response)

Medications may be administered by oral or IV route. The equivalent of these medications may be administered as determined by the attending or referring physician.

In a true medical emergency situation the exam should proceed with an accelerated premedication schedule at the direction of the attending physician. (attachment to Radiology Policy R 711)

<input type="checkbox"/> Oral methylPREDNISolone (MEDROL) tablet	32 mg, oral, once, For 1 Doses, Pre-Procedure For iodine allergy prophylaxis. Give 6-12 hours prior to the examination.
<input type="checkbox"/> IV methylPREDNISolone succinate (Solu-MEDROL) 40 mg/mL injection	40 mg, intravenous, for 5 Minutes, once, For 1 Doses, Pre-Procedure For iodine allergy prophylaxis. Give 2 hours prior to the examination.

For patients with allergic reaction

<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection	50 mg, intravenous, once, For 1 Doses, Pre-Procedure Given in Interventional Radiology unit
<input type="checkbox"/> methylPREDNISolone sodium succinate (Solu-MEDROL) injection	125 mg, intravenous, once, For 1 Doses, Pre-Procedure Given in Interventional Radiology unit

VTE

Labs

COVID-19 Qualitative PCR

<input type="checkbox"/> COVID-19 qualitative RT-PCR - Nasal Swab	STAT For 1 Occurrences Specimen Source: Nasal Swab Is this for pre-procedure or non-PUI assessment? Yes Pre-Procedure
---	--

Labs

<input type="checkbox"/> CBC and differential	Once, Starting S, Pre-Procedure
<input type="checkbox"/> Comprehensive metabolic panel	Once, Pre-Procedure
<input type="checkbox"/> Basic metabolic panel	Once, Starting S, Pre-Procedure
<input type="checkbox"/> Protime-INR	Once, Starting S, Pre-Procedure
<input type="checkbox"/> Platelet count	Once, Pre-Procedure
<input type="checkbox"/> Partial thromboplastin time	Once, Starting S, Pre-Procedure
<input type="checkbox"/> hCG qualitative, serum	Once, Pre-Procedure

Cardiology**Imaging****Other Studies****Respiratory****Rehab****Consults**

For Physician Consult orders use sidebar

Additional Orders