

# Retinopathy of Prematurity Exam [714]

## General

## Nursing

### Nursing

- |  |   |
|--|---|
| <input type="checkbox"/> Provide equipment / supplies at bedside- Retinopathy      | Routine, Once For 1 Occurrences<br>Supplies: Other (specify)<br>Other: Have the sterile lid speculum and lens loop at the infant's bedside. |
| <input type="checkbox"/> Apply phototherapy mask to eyes for 4 hours post ROP exam | Routine, Continuous<br>Phototherapy: Double<br>Apply phototherapy mask to eyes 4 hours post dilation.                                       |

## IV Fluids

## Medications

### Medications

- |  |   |
|--|---|
| <input checked="" type="checkbox"/> cyclopentolate-phenylephrine (CYCLOMYDRYL) 0.2-1 % ophthalmic solution | 1 drop, Both Eyes, PRN, instill every 5 minutes for 3 doses<br>To be administered 45 minutes PRIOR to eye exam. |
| <input checked="" type="checkbox"/> sucrose 24 % (TOOT SWEET) oral solution                                | 0.1 mL, oral, once PRN, mild pain (score 1-3), pain   |
| <input checked="" type="checkbox"/> proparacaine (ALCAINE) 0.5 % ophthalmic solution                       | 1 drop, Both Eyes, PRN, for eye exam<br>To be administered 30 seconds PRIOR to eye exam.                        |
| <input type="checkbox"/> artificial tears(hypromellose) (SYSTANE/GENTEAL) 0.3 % ophthalmic gel             | 1 application, Both Eyes, PRN, dry eyes, for eye exam<br>To be administered for each eye exam.                  |

## VTE

## Labs

## Cardiology

## Diagnostic Imaging

## Other Diagnostic Studies

## Respiratory

## Rehab

## Consults

## Additional Orders