Adjust doses based on therapeutic response and digoxin level monitoring.

For elderly patients (GREATER THAN 65 years), maximum maintenance dose of 125 mcg daily is recommended.

Medications

Digoxin (LANOXIN) (Selection Required)

() Maintenance (Selection Required)

[X] Digoxin Maintenance Only (Selection Required)

(X) digOXIN (LANOXIN) tablet

oral, Daily, Indication: Arrhythmia, HR HOLD parameters for this order: HR Hold Parameters requested, HR HOLD for: Heart Rate LESS than 60 bpm

() digOXIN (LANOXIN) oral solution

oral, Daily, Indication: Arrhythmia, HR HOLD parameters for this order: HR Hold Parameters requested, HR HOLD for: Heart Rate LESS than 60 bpm

() digOXIN (LANOXIN) injection

intravenous, Daily, Indication: Arrhythmia, HR HOLD parameters for this order: HR Hold Parameters requested, HR HOLD for: Heart Rate LESS than 60 bpm

[X] Digoxin level

AM Draw, For 1 Occurrences

() Load & Maintenance

() Digoxin Load & Maintenance - NO active Amiodarone Order (Selection Required)

[X] Digoxin level

AM Draw, Starting S+5, For 1 Occurrences

[X] Digoxin load and maintenance based on CrCl (Selection Required)

() CrCl GREATER than or EQUAL to 30 mL/min (Selection Required)

[X] Digoxin Load and Maintenance (Selection Required)

digOXIN (LANOXIN) injection

5 mcg/kg, intravenous, Q6H, Starting Today, For 1 Doses, Indication: Arrhythmia

Followed by

digOXIN (LANOXIN) injection

2.5 mcg/kg, intravenous, Q6H, For 2 Doses, Indication: Arrhythmia

Followed by

digOXIN (LANOXIN) tablet

125 mcg, oral, Daily, Starting Tomorrow, Indication: Arrhythmia, HR HOLD parameters for this order: HR Hold Parameters requested, HR HOLD for: Heart Rate LESS than 60 bpm

[X] Digoxin level

Once, Starting 18 Hours after signing, For 1 Occurrences

() CrCl LESS than 30 mL/min (Selection Required)

[X] HM IP DIGOXIN 5MCG/KG X 1 X 62.5 MG TAB (Selection Required)

digOXIN (LANOXIN) injection

5 mcg/kg, intravenous, Once, For 1 Doses, Indication: Arrhythmia

Followed by

digOXIN (LANOXIN) tablet

62.5 mcg, oral, Daily, Starting Tomorrow

[X] Digoxin level

Once, Starting 12 Hours after signing, For 1 Occurrences

() Digoxin Load & Maintenance WITH active Amiodarone Order (Selection Required)

[X] Digoxin level

AM Draw, Starting S+5, For 1 Occurrences

[X] Digoxin load and maintenance based on CrCl (Selection Required) () CrCl greater than or equal to 30 mL/min (Selection Required) [X] Digoxin Load and Maintenance (Selection Required) digOXIN (LANOXIN) injection 5 mcg/kg, intravenous, Once, For 1 Doses, Indication: Arrhythmia Followed by digOXIN (LANOXIN) tablet 125 mcg, oral, Daily, Starting Tomorrow [X] Digoxin level Once, Starting 12 Hours after signing, For 1 Occurrences () CrCl LESS than 30 mL/min (Selection Required) [X] HM IP DIGOXIN 2.5 MCG/KG X 1 X 62.5 TAB (Selection Required) digOXIN (LANOXIN) injection 2.5 mcg/kg, intravenous, Once, For 1 Doses, Indication: Arrhythmia Followed by digOXIN (LANOXIN) tablet 62.5 mcg, oral, Daily, Starting Tomorrow [X] Digoxin level Once, Starting 12 Hours after signing, For 1 Occurrences () Load (Selection Required) () Load Only - WITHOUT active amiodarone orders (Selection Required) () CrCl Greater than or Equal to 30 mL/min (Selection Required) [X] HM IP DIGOXIN IV NO ACTIVE AMIODARONE LOAD ONLY (Selection Required) digOXIN (LANOXIN) injection 5 mcg/kg, intravenous, Once, For 1 Doses Followed by digOXIN (LANOXIN) injection 2.5 mcg/kg, intravenous, Q6H, For 2 Doses, Indication: Arrhythmia [X] Digoxin level Once, Starting 18 Hours after signing, For 1 Occurrences () CrCl LESS THAN 30 mL/min (Selection Required) [X] digOXIN (LANOXIN) injection 5 mcg/kg, intravenous, Once, For 1 Doses, Indication: Arrhythmia Once, Starting 12 Hours after signing, For 1 Occurrences () Load Only -WITH concurrent active amiodarone use (Selection Required) () CrCl GREATER THAN or EQUAL 30 mL/min (Selection Required) [X] digOXIN (LANOXIN) injection 5 mcg/kg, intravenous, Once, For 1 Doses, Indication: Arrhythmia [X] Digoxin level Once, Starting 12 Hours after signing, For 1 Occurrences () CrCl LESS THAN 30 mL/min (Selection Required) [X] digOXIN (LANOXIN) injection 2.5 mcg/kg, intravenous, Once, For 1 Doses, Indication: Arrhythmia [X] Digoxin level Once, Starting 12 Hours after signing, For 1 Occurrences () One Time Dose Only (Selection Required) () digOXIN (LANOXIN) tablet

oral, Once, For 1 Doses

() digOXIN (LANOXIN) oral solution

oral, Once, For 1 Doses

() digOXIN (LANOXIN) injection

intravenous, Once, For 1 Doses

Magnesium Baseline Level

[X] Magnesium level

AM Draw, For 1 Occurrences

Digoxin Follow-up Level

[] Digoxin level

AM Draw, Starting S+5, At: 0400, For 1 Occurrences