

Location: _____

General

Vital Signs

☐ **Vital signs - T/P/R/BP** Per unit protocol, STAT

Labs

Exposure Labs

☒ **HIV 1/2 antigen/antibody, fourth generation, with reflexes** Once, 1, Occurrences, STAT, Blood, 3
Release to patient (Note: If manual release option is selected, result will auto release 5 days from finalization.):☒ **Hepatitis B surface antibody** Once, 1, Occurrences, STAT, Blood, 3☒ **Hepatitis B surface antigen** Once, 1, Occurrences, STAT, Blood, 3☒ **Hepatitis C Virus (HCV) Antibody With Reflex to PCR** Once, 1, Occurrences, STAT, 3☒ **Comprehensive metabolic panel** Once, 1, Occurrences, STAT, Blood, 3☒ **CBC with platelet and differential** Once, 1, Occurrences, STAT, Blood, 3

Medications

HIV Prophylaxis Medications

☐ **Monday - Thursday (HIV Prophylaxis Medications)**☒ **emtricitabine-tenofovir DF (TRUVADA) + dolutegravir (TIVICAY)**☒ **emtricitabine-tenofovir DF (TRUVADA) 200-300 mg per tablet** 1 tablet, oral, daily

Indication: ○ Medical Prophylaxis

☒ **dolutegravir (TIVICAY) tablet** 50 mg, oral, daily

Indication: ○ Medical Prophylaxis

Administer 2 hours before any milk product, antacid or iron.

☒ **Discharge medication prescriptions - emtricitabine-tenofovir DF (TRUVADA) and dolutegravir (TIVICAY) tablets**☒ **emtricitabine-tenofovir DF (Truvada) 200-300 mg per tablet** 200-300 , daily

Indication:

☒ **dolutegravir (Tivicay) 50 mg tablet** 50 , daily

Indication:

Administer 2 hours before any milk product, antacid or iron.

☐ **Friday - Sunday (HIV Prophylaxis Medications)**☒ **First dose and Take Home Med - Truvada and Tivicay tablets**☒ **First Dose - emtricitabine-tenofovir DF (TRUVADA) tablet 200-300 mg** 1 tablet, oral, once, 1, Occurrences

Indication: ○ Medical Prophylaxis

Administer first dose today.

☒ **First Dose - dolutegravir (TIVICAY) tablet 50 mg** 50 mg, oral, once, 1, Occurrences

Indication: Medical Prophylaxis

Administer first dose today.

Administer 2 hours before any milk product, antacid or iron.

☒ **Take Home Med - emtricitabine-tenofovir DF (TRUVADA) tablet 200-300 mg** 1 tablet, oral, daily, 3, Days

Indication: ○ Medical Prophylaxis

RN: do NOT administer. Patient to take 1 tablet by mouth daily starting tomorrow. Pharmacy to dispense 3 tablets to patient to take at home.

☒ **Take Home Med - dolutegravir (TIVICAY) tablet 50 mg** 50 mg, oral, daily, 3, Days

Indication: Medical Prophylaxis

RN: do NOT administer. Patient to take 1 tablet by mouth daily starting tomorrow. Pharmacy to dispense 3 tablets to patient to take at home.

Administer 2 hours before any milk product, antacid or iron.

☒ **Discharge medication prescriptions - emtricitabine-tenofovir DF (TRUVADA) and dolutegravir (TIVICAY) tablets**☒ **emtricitabine-tenofovir DF (Truvada) 200-300 mg per tablet** 200-300 , daily

Indication:

☒ **dolutegravir (Tivicay) 50 mg tablet** 50 , daily

Indication:

Administer 2 hours before any milk product, antacid or iron.

Sign: _____ Printed Name: _____ Date/Time: _____

Meningitis

☐ **riFAMPin (RIFADIN) capsule** 600 mg, oral, once, 1, Occurrences, STAT

Indication: ○ CNS

Give for 2 days with q12h dosing regimen.

☐ **ciprofloxacin HCl (CIPRO) tablet** 500 mg, oral, once, 1, Occurrences, STAT

Indication: ○ CNS

Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.:

May cause QTc prolongation. Tablets may be crushed if needed to be administered via a feeding tube route.

If administering with tube feeds, mix with water to avoid interaction with tube feed.

Sign: _____ Printed Name: _____ Date/Time: _____