

Location: _____

General

Common Present on Admission Diagnosis

- ☐ **Acidosis** Frequency: Once Priority: Routine
- ☐ **Acute Post-Hemorrhagic Anemia** Frequency: Once Priority: Routine
- ☐ **Acute Renal Failure** Frequency: Once Priority: Routine
- ☐ **Acute Respiratory Failure** Frequency: Once Priority: Routine
- ☐ **Acute Thromboembolism of Deep Veins of Lower Extremities** Frequency: Once Priority: Routine
- ☐ **Anemia** Frequency: Once Priority: Routine
- ☐ **Bacteremia** Frequency: Once Priority: Routine
- ☐ **Bipolar disorder, unspecified** Frequency: Once Priority: Routine
- ☐ **Cardiac Arrest** Frequency: Once Priority: Routine
- ☐ **Cardiac Dysrhythmia** Frequency: Once Priority: Routine
- ☐ **Cardiogenic Shock** Frequency: Once Priority: Routine
- ☐ **Decubitus Ulcer** Frequency: Once Priority: Routine
- ☐ **Dementia in Conditions Classified Elsewhere** Frequency: Once Priority: Routine
- ☐ **Disorder of Liver** Frequency: Once Priority: Routine
- ☐ **Electrolyte and Fluid Disorder** Frequency: Once Priority: Routine
- ☐ **Intestinal Infection due to Clostridium Difficile** Frequency: Once Priority: Routine
- ☐ **Methicillin Resistant Staphylococcus Aureus Infection** Frequency: Once Priority: Routine
- ☐ **Obstructive Chronic Bronchitis with Exacerbation** Frequency: Once Priority: Routine
- ☐ **Other Alteration of Consciousness** Frequency: Once Priority: Routine
- ☐ **Other and Unspecified Coagulation Defects** Frequency: Once Priority: Routine
- ☐ **Other Pulmonary Embolism and Infarction** Frequency: Once Priority: Routine
- ☐ **Phlebitis and Thrombophlebitis** Frequency: Once Priority: Routine
- ☐ **Protein-calorie Malnutrition** Frequency: Once Priority: Routine
- ☐ **Psychosis, unspecified psychosis type** Frequency: Once Priority: Routine
- ☐ **Schizophrenia Disorder** Frequency: Once Priority: Routine
- ☐ **Sepsis** Frequency: Once Priority: Routine
- ☐ **Septic Shock** Frequency: Once Priority: Routine
- ☐ **Septicemia** Frequency: Once Priority: Routine
- ☐ **Type II or Unspecified Type Diabetes Mellitus with Mention of Complication, Not Stated as Uncontrolled** Frequency: Once Priority: Routine
- ☐ **Urinary Tract Infection, Site Not Specified** Frequency: Once Priority: Routine

Elective Outpatient, Observation, or Admission

- ☐ **Elective outpatient procedure: Discharge following routine recovery** Frequency: Continuous Phase of Care: PACU & Post-op Priority: Routine
- ☐ **Outpatient observation services under general supervision** Frequency: Once Phase of Care: PACU & Post-op Priority: Routine

Question(s):

Admitting Physician:

Attending Provider:

Patient Condition:

Bed request comments:

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Outpatient in a bed - extended recovery** Frequency: Once Phase of Care: PACU & Post-op Priority: Routine

Question(s):

Admitting Physician:

Bed request comments:

☐ **Admit to Inpatient** Frequency: Once Ordering Quantity: 1 Phase of Care: PACU & Post-op Priority: Routine

Question(s):

Admitting Physician:

Level of Care:

Patient Condition:

Bed request comments:

Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.

Admission or Observation**Patient has active status order on file**

☐ **Admit to Inpatient** Frequency: Once Ordering Quantity: 1 Priority: Routine

Question(s):

Admitting Physician:

Level of Care:

Patient Condition:

Bed request comments:

Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.

☐ **Outpatient observation services under general supervision** Frequency: Once Priority: Routine

Question(s):

Admitting Physician:

Attending Provider:

Patient Condition:

Bed request comments:

☐ **Outpatient in a bed - extended recovery** Frequency: Once Priority: Routine

Question(s):

Admitting Physician:

Bed request comments:

Admission or Observation**Patient has active status order on file**

☐ **Admit to Inpatient** Frequency: Once Ordering Quantity: 1 Priority: Routine

Question(s):

Admitting Physician:

Level of Care:

Patient Condition:

Bed request comments:

Certification: I certify that based on my best clinical judgment and the patient's condition as documented in the HP and progress notes, I expect that the patient will need hospital services for two or more midnights.

☐ **Outpatient observation services under general supervision** Frequency: Once Priority: Routine

Question(s):

Admitting Physician:

Attending Provider:

Patient Condition:

Bed request comments:

☐ **Outpatient in a bed - extended recovery** Frequency: Once Priority: Routine

Question(s):

Admitting Physician:

Bed request comments:

Code Status

@CERMSGREFRESHOPT(674511:21703,,1)@

☒ **Code Status**

DNR and Modified Code orders should be placed by the responsible physician.

☐ **Full code** Frequency: Continuous Priority: Routine

Question(s):

Code Status decision reached by:

☐ **DNR (Do Not Resuscitate)** (Required)

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **DNR (Do Not Resuscitate) Frequency:** Continuous **Priority:** Routine**Question(s):**

Did the patient/surrogate require the use of an interpreter?

Did the patient/surrogate require the use of an interpreter?

Does patient have decision-making capacity?

I acknowledge that I have communicated with the patient/surrogate/representative that the Code Status order is NOT Active until Signed by the Responsible Attending Physician.:

Code Status decision reached by:

☐ **Consult to Palliative Care Service**☒ **Consult to Palliative Care Service Frequency:** Once **Priority:** Routine**Question(s):**

Priority:

Reason for Consult?

Order?

Name of referring provider:

Enter call back number:

Reason for Consult?

Process Instructions:

Note: Please call Palliative care office 832-522-8391. Due to current resource constraints, consultation orders received after 2:00 pm M-F will be seen the following business day. Consults placed over weekend will be seen on Monday.

☐ **Consult to Social Work Frequency:** Once **Priority:** Routine**Question(s):**

Reason for Consult:

Reason for Consult?

☐ **Modified Code Frequency:** Continuous **Priority:** Routine**Question(s):**

Did the patient/surrogate require the use of an interpreter?

Did the patient/surrogate require the use of an interpreter?

Does patient have decision-making capacity?

Modified Code restrictions:

I acknowledge that I have communicated with the patient/surrogate/representative that the Code Status order is NOT Active until Signed by the Responsible Attending Physician.:

Code Status decision reached by:

☐ **Treatment Restrictions ((For use when a patient is NOT in a cardiopulmonary arrest)) Frequency:** Continuous -
Treatment Restrictions **Priority:** Routine**Question(s):**

I understand that if the patient is NOT in a cardiopulmonary arrest, the selected treatments will NOT be provided. I understand that all other unselected medically indicated treatments will be provided.:

Treatment Restriction decision reached by:

Specify Treatment Restrictions:

Code Status decision reached by:

Process Instructions:

Treatment Restrictions is NOT a Code Status order. It is NOT a Modified Code order. It is strictly intended for Non Cardiopulmonary situations.

The Code Status and Treatment Restrictions are two SEPARATE sets of physician's orders. For further guidance, please click on the link below: [Guidance for Code Status & Treatment Restrictions](#)

Examples of Code Status are Full Code, DNR, or Modified Code. An example of a Treatment Restriction is avoidance of blood transfusion in a Jehovah's Witness patient.

If the Legal Surrogate is the Primary Physician, consider ordering a Biomedical Ethics Consult PRIOR to placing this order. A Concurring Physician is required to second sign the order when the Legal Surrogate is the Primary Physician.

Isolation☐ **Airborne isolation status**☒ **Airborne isolation status Frequency:** Continuous **Priority:** Routine☐ **Mycobacterium tuberculosis by PCR - If you suspect Tuberculosis, please order this test for rapid diagnostics.****Frequency:** Once **Priority:** Routine

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Contact isolation status** Frequency: Continuous Priority: Routine

☐ **Droplet isolation status** Frequency: Continuous Priority: Routine

☐ **Enteric isolation status** Frequency: Continuous Priority: Routine

Precautions

☐ **Aspiration precautions** Frequency: Continuous Priority: Routine

☐ **Fall precautions** Frequency: Continuous Priority: Routine

Question(s):

Increased observation level needed:

☐ **Latex precautions** Frequency: Continuous Priority: Routine

☐ **Seizure precautions** Frequency: Continuous Priority: Routine

Question(s):

Increased observation level needed:

Nursing

Vital Signs

☒ **Vital signs - T/P/R/BP** Frequency: Per unit protocol Priority: Routine

Activity/Patient Position

☐ **Strict bed rest** Frequency: Until discontinued Priority: Routine Comments: Up in chair in AM

☐ **Up in chair** Frequency: Until discontinued Priority: Routine

Question(s):

Specify: ☐ Up in chair

☐ **Ambulate with assistance (specify frequency)** Frequency: 3 times daily Priority: Routine

Question(s):

Specify:

☐ **Ambulate with assistance every 4 hours** Frequency: Now then every 4 hours Priority: Routine

Question(s):

Specify: ☐ with assistance

☐ **Ambulate with assistance every 8 hours** Frequency: Every 8 hours Priority: Routine

Question(s):

Specify: ☐ with assistance

☐ **Activity as tolerated** Frequency: Until discontinued Priority: Routine

Question(s):

Specify: ☐ Activity as tolerated

☐ **Avoid pressure to** Frequency: Once Priority: Routine

Question(s):

Orientation:

Location:

☐ **Head of bed** Frequency: Until discontinued Priority: Routine

Question(s):

Head of bed:

☐ **Patient position: Elevate foot of bed** Frequency: Until discontinued Priority: Routine

Question(s):

Additional instructions: ☐ elevate foot of bed

Position:

☐ **Patient position: Semi-Fowler's** Frequency: Until discontinued Priority: Routine Comments: With bed flexed in semi-fowler's (lawn chair) position.

Question(s):

Position: ☐ semi-Fowler's

Additional instructions:

☐ **Patient position: Do not reposition patient** Frequency: Until discontinued Priority: Routine

Question(s):

Additional instructions: ☐ do not reposition

Position:

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Shower patient** Frequency: Daily Priority: Routine

Question(s):

Specify:

Additional modifier:

Nursing Care

☐ **Neurological assessment** Frequency: Once Priority: Routine

Question(s):

Assessment to Perform:

☐ **Peripheral vascular assessment** Frequency: Once Priority: Routine

☐ **Assess head** Frequency: Every 4 hours Priority: Routine

Question(s):

Assess: o Head (Facial, eyelids) for color, refill, hematoma. Notify Resident or staff for any changes.

☐ **Assess breast** Frequency: Every 4 hours Priority: Routine

Question(s):

Assess: o Breast - assess nipple for color, refill, and hematoma. Notify Resident or staff for any changes.

☐ **Assess abdomen** Frequency: Every 4 hours Priority: Routine

Question(s):

Assess: o Abdomen - assess for color, refill, and hematoma. Notify Resident or staff for any changes.

☐ **Assess On-Q Pump** Frequency: Every 4 hours Priority: Routine

Question(s):

Assess: o Assess On-Q Pump every 4 hours

☐ **Intake and output** Frequency: Per unit protocol Priority: Routine Comments: Include amount from surgical drain in intake and output

☐ **No ice pack** Frequency: Until discontinued Priority: Routine Comments: Unless ordered otherwise

☐ **Limb precautions** Frequency: Continuous Priority: Routine

Question(s):

Location:

Precaution:

☐ **May use either arm for blood pressure or needle sticks** Frequency: Until discontinued Priority: Routine

☐ **Supportive bra** Frequency: Until discontinued Priority: Routine Comments: Do not remove post-operative bra.

☐ **Abdominal binder** Frequency: Once Priority: Routine Comments: Keep abdominal binder open and loose while in bed.

When patient gets up in chair, place binder on. Open when back in bed.

Question(s):

Waking hours only?

Nurse to schedule?

Special Instructions:

☐ **Compression garment** Frequency: Until discontinued Priority: Routine

Question(s):

Intervention: o Remove every 3 hours for 1 hour

Flap Assessment

☐ **Flap assessment** Frequency: Every hour Priority: Routine Comments: Notify Resident or staff for any changes.

Question(s):

Side:

Location:

Tubes and Drain Care

☐ **Drain care- Compression Suction; Attach bulbs to gown with safety pins. Do NOT tape drains to patient.; Strip tubing and record output every 4 hours** Frequency: Every 4 hours Priority: Routine

Question(s):

All Drains: o Jackson Pratt

Care Details: Attach bulbs to gown with safety pins. Do NOT tape drains to patient.

Drainage/Suction: o To Compression (Bulb) Suction o Strip tubing o Other (specify)

Specify: Empty drain and record output every 4 hours.

Drain 1:

Drain 2:

Drain 3:

Drain 4:

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Drain care- Clean site daily with normal saline. Apply ointment and cover with gauze. Frequency: Daily Priority:**

Routine

Question(s):

All Drains: ☐ Jackson Pratt

Care Details: Clean site daily with normal saline. Apply ointment and cover with gauze.

Drainage/Suction: To Compression (Bulb) Suction

Drain 1:

Drain 2:

Drain 3:

Drain 4:

☐ **Drain care- Clean site daily with peroxide. Apply bacitracin ointment and cover with gauze. Frequency: Daily Priority:**

Routine

Question(s):

All Drains: ☐ Jackson Pratt

Care Details: Clean site daily with peroxide. Apply bacitracin ointment and cover with gauze.

Drainage/Suction: To Compression (Bulb) Suction

Drain 1:

Drain 2:

Drain 3:

Drain 4:

☐ **Do not remove Foley Frequency: Until discontinued Priority: Routine**

Question(s):

Rationale:

☐ **Foley catheter care Frequency: Until discontinued Priority: Routine**

Question(s):

Orders: ☐ Maintain ☐ to gravity

☐ **Remove Foley catheter Frequency: Once Priority: Routine**

Wound/Incision Care

☐ **Surgical/incision site care- Wet to dry, Normal Saline Frequency: Every 8 hours Priority: Routine**

Question(s):

Dressing Type: ☐ Moist to Dry, Normal Saline

Location:

Site:

Apply:

Open to air?

☐ **Surgical/incision site care- Wet to dry, Dakins Frequency: Every 8 hours Priority: Routine**

Question(s):

Dressing Type: ☐ Other

Specify: Wet to dry, Dakins

Location:

Site:

Apply:

Open to air?

☐ **Surgical/incision site care- Do not remove dressing Frequency: Once Priority: Routine Comments: Do not remove or change surgical dressing**

Question(s):

Location:

Site:

Apply:

Dressing Type:

Open to air?

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Wound care orders** Frequency: Every 12 hours Priority: Routine

Question(s):

Location:

Site:

Irrigate wound?

Apply:

Dressing Type:

Process Instructions:

This Nursing Order is NOT for a CONSULT for PT Wound Care or WOC nurse. The order is not transmitted to any department.

Do NOT use this order to request :

Bedside debridement, Ultrasound Therapy, Pulsed Lavage, Negative Pressure Vacuum Therapy, Compression therapy, WOC ongoing wound /ostomy management and teaching.

☐ **Provide equipment / supplies at bedside** Frequency: Once Priority: Routine

Question(s):

Supplies:

☐ **Provide equipment / supplies at bedside: Extra Bra to bedside** Frequency: Once Priority: Routine Comments: Size ***

Question(s):Supplies: ☐ Other (specify)

Other: Extra Bra to bedside.

☐ **Negative pressure wound therapy (Not a consult order)** Frequency: Every Mon, Wed, Fri Priority: Routine

Question(s):

Existing wound vac?

Type of Wound:

Wound Location:

Pressure (mmHg): 125

Therapy Settings:

Intensity:

Foam Type:

☐ **Consult to Wound Ostomy Care Nurse** Frequency: Once Priority: Routine

Question(s):

Reason for consult:

Reason for consult:

Reason for consult:

Reason for consult:

Consult for NPWT:

Reason for consult:

Reason for consult:

Reason for Consult?

Process Instructions:

This is NOT for PT Wound Care Consult order.

Skin Graft Donor Site Care

☐ **Heat lamp** Frequency: 4 times daily Priority: Routine

Question(s):Duration of treatment (minutes): ☐ 15Distance from site: ☐ 2-3 feet

☐ **Skin graft donor site care** Frequency: 4 times daily Priority: Routine

Question(s):

Instructions: ☐ Leave donor site intact for 48 hours, clean any excessive fluid leakage as needed. After 48 hours, remove clear Tegaderm but DO NOT remove Zeroform gauze. Wipe off any excess fluid gently PRN. Use heat lamp to treat donor site 4 x/day when patient is aw

☐ **Negative pressure wound therapy (Not a consult order)** Frequency: Every Mon, Wed, Fri Priority: Routine Comments:

DO NOT change negative pressure wound therapy dressing**

Question(s):

NPWT to be applied by: Physician

Pressure (mmHg): 125

Existing wound vac?

Type of Wound:

Wound Location:

Therapy Settings:

Intensity:

Foam Type:

Sign: _____ Printed Name: _____ Date/Time: _____

Notify

- ☐ **Notify Physician- Notify Plastic Surgery resident on-call or Plastics Attending Surgeon for ANY questions regarding the flap or change in flap assessment** Frequency: Until discontinued Priority: Routine Comments: Notify Plastic Surgery resident on-call or Plastics Attending Surgeon for ANY questions regarding the flap or change in flap assessment
- ☐ **Notify Physician- or Resident of any acute changes in patient status** Frequency: Until discontinued Priority: Routine Comments: or Resident of any acute changes in patient status

Diet

- ☐ **NPO** Frequency: Diet effective now Priority: Routine

Question(s):

NPO:

Pre-Operative fasting options:

Process Instructions:

An NPO order without explicit exceptions means nothing can be given orally to the patient.

- ☐ **NPO after midnight** Frequency: Diet effective midnight Priority: Routine

Question(s):

NPO:

Pre-Operative fasting options:

Process Instructions:

An NPO order without explicit exceptions means nothing can be given orally to the patient.

- ☐ **Diet- Clear Liquids** Frequency: Diet effective now Priority: Routine

Question(s):

Diet(s): o Clear Liquids

Cultural/Special:

Cultural/Special:

Other Options:

Other Options:

Advance Diet as Tolerated?

IDDSI Liquid Consistency:

Fluid Restriction:

Foods to Avoid:

Foods to Avoid:

- ☐ **Diet - Easy to digest (GERD)** Frequency: Diet effective now Priority: Routine

Question(s):

Diet(s): o Easy to digest (GERD)

Cultural/Special:

Cultural/Special:

Other Options:

Other Options:

Advance Diet as Tolerated?

IDDSI Liquid Consistency:

Fluid Restriction:

Foods to Avoid:

Foods to Avoid:

- ☐ **Diet - 1800 Kcal / 202 gm Carbohydrate** Frequency: Diet effective now Priority: Routine

Question(s):

Diet(s): o Consistent Carbohydrate

Consistent Carbohydrate: 1800 Kcal/202 gm Carbohydrate

Cultural/Special:

Cultural/Special:

Other Options:

Other Options:

Advance Diet as Tolerated?

IDDSI Liquid Consistency:

Fluid Restriction:

Foods to Avoid:

Foods to Avoid:

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Diet- Regular Frequency:** Diet effective now **Priority:** Routine

Question(s):Diet(s): ☐ Regular

Cultural/Special:

Cultural/Special:

Other Options:

Other Options:

Advance Diet as Tolerated?

IDDSI Liquid Consistency:

Fluid Restriction:

Foods to Avoid:

Foods to Avoid:

Education

☐ **Patient education- Drain care Frequency:** Prior to discharge **Priority:** Routine

Question(s):Education for: ☐ Drain care

Patient/Family:

☐ **Patient education- Dressing change Frequency:** Once **Priority:** Routine

Question(s):Education for: ☐ Other (specify)

Specify: Dressing change

Patient/Family:

☐ **Patient education- Lovenox teaching Frequency:** Prior to discharge **Priority:** Routine

Question(s):Education for: ☐ Self admin of medication ☐ Other (specify)

Specify: Lovenox teaching for home administration.

Patient/Family:

☐ **Patient education- Pain pump Frequency:** Prior to discharge **Priority:** Routine

Question(s):Patient/Family: ☐ PatientEducation for: ☐ Other (specify)

Specify: Pain pump

☐ **Patient education- Post-op urine color Frequency:** Once **Priority:** Routine

Question(s):Patient/Family: ☐ BothEducation for: ☐ Other (specify)

Specify: Blue/green urine post op is normal

☐ **Patient education- Scopolamine patch Frequency:** Once **Priority:** Routine

Question(s):Education for: ☐ Other (specify)

Specify: Scopolamine patch side effects education

Patient/Family:

☐ **Patient education- Surgeons post op instructions Frequency:** Prior to discharge **Priority:** Routine

Question(s):Education for: ☐ Other (specify)

Specify: Dispense surgeon's post op instructions prior to discharge.

Patient/Family:

IV Fluids**IV Fluids**

☐ **dextrose 5 % and sodium chloride 0.45 % with potassium chloride 20 mEq/L infusion Dose:** 125 mL/hr **Route:** intravenous **Frequency:** continuous **Phase of Care:** Post-op

☐ **lactated Ringer's infusion Dose:** 125 mL/hr **Route:** intravenous **Frequency:** continuous **Phase of Care:** Post-op

☐ **sodium chloride 0.9 % infusion Dose:** 125 mL/hr **Route:** intravenous **Frequency:** continuous **Phase of Care:** Post-op

☐ **sodium chloride 0.9 % with potassium chloride 20 mEq/L infusion Dose:** 125 mL/hr **Route:** intravenous **Frequency:** continuous **Phase of Care:** Post-op

☐ **sodium chloride 0.45 % with potassium chloride 20 mEq/L infusion Dose:** 125 mL/hr **Route:** intravenous **Frequency:** continuous **Phase of Care:** Post-op

Sign: _____ Printed Name: _____ Date/Time: _____

Medications

Reminder: If you need to place orders for PCA Analgesia, after using this order set go to the [Order Set Activity](#) and access the **General Patient Controlled Analgesia (PCA) Therapy for Opioid Naive Patients** (or Tolerant Patients if appropriate).

Pharmacy Consult

☐ **Pharmacy consult to manage dosing of medication** Frequency: Until discontinued Priority: STAT

Question(s):

Which drug do you need help dosing?

Contact Number:

IV Antibiotics: For Patients LESS than or EQUAL to 120 kg

☐ **ampicillin-sulbactam (UNASYN) IV** Dose: 3 g Route: intravenous Frequency: every 6 hours Priority: STAT

Question(s):

Indication: ○ Surgical Prophylaxis

Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.:

☐ **cefazolin (ANCEF) IV - For Patients LESS than or EQUAL to 120 kg** Dose: 2 g Route: intravenous Frequency: every 8 hours Priority: STAT

Question(s):

Indication: ○ Surgical Prophylaxis

Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.:

☐ **cefepime (MAXIPIME) IV - For antipseudomonal coverage** Dose: 1 g Route: intravenous Frequency: every 8 hours Priority: STAT

Question(s):

Indication: ○ Surgical Prophylaxis

Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.:

Admin Instructions:

STANDARD Infusion

☐ **clindamycin (CLEOCIN) IV** Dose: 900 mg Route: intravenous Frequency: every 8 hours Priority: STAT

Question(s):

Indication: Surgical Prophylaxis

Indication:

Admin Instructions:

Use if patient penicillin allergic.

☐ **vancomycin IV plus Optional Pharmacy Consult to Dose Vancomycin**

☒ **IV Vancomycin Loading Dose + Pharmacy Consult**

☒ **vancomycin (VANCOCIN) IV** Route: intravenous Frequency: once Frequency Limit: 1 Occurrences Priority: STAT

Question(s):

Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.:

Indication:

Admin Instructions:

Loading Dose

☒ **Pharmacy consult to manage vancomycin** Frequency: Until discontinued Priority: Routine

Question(s):

Indication:

Anticipated Duration of Vancomycin Therapy (Days):

Process Instructions:

All eligible patients to receive Vancomycin at AUC 400-600 and Trough 10-20.

IV Antibiotics: For Patients GREATER than 120 kg

☐ **ampicillin-sulbactam (UNASYN) IV** Dose: 3 g Route: intravenous Frequency: every 6 hours Priority: STAT

Question(s):

Indication: ○ Surgical Prophylaxis

Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.:

☐ **cefazolin (ANCEF) IV - For Patients GREATER than 120 kg** Dose: 3 g Route: intravenous Frequency: every 8 hours Priority: STAT

Question(s):

Indication: ○ Surgical Prophylaxis

Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.:

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **cefepime (MAXIPIME) IV - For antipseudomonal coverage** Dose: 2 g Route: intravenous Frequency: every 8 hours

Priority: STAT

Question(s):

Indication: o Surgical Prophylaxis

Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.:

Admin Instructions:

STANDARD Infusion

☐ **clindamycin (CLEOCIN) IV** Dose: 900 mg Route: intravenous Frequency: every 8 hours Priority: STAT

Question(s):

Indication: Surgical Prophylaxis

Indication:

Admin Instructions:

Use if patient penicillin allergic.

☐ **IV Vancomycin Loading Dose + Pharmacy Consult**

☒ **vancomycin (VANCOCIN) IV** Route: intravenous Frequency: once Frequency Limit: 1 Occurrences Priority: STAT

Question(s):

Per Med Staff Policy, R.Ph. will automatically renally dose this medication based on current SCr and CrCl values.:

Indication:

Admin Instructions:

Loading Dose

☒ **Pharmacy consult to manage vancomycin** Frequency: Until discontinued Priority: Routine

Question(s):

Indication:

Anticipated Duration of Vancomycin Therapy (Days):

Process Instructions:

All eligible patients to receive Vancomycin at AUC 400-600 and Trough 10-20.

Topical Antibiotics

☐ **bacitracin ointment** Dose: 500 Route: Topical Frequency: 3 times daily

Admin Instructions:

Apply to drain site.

☐ **bacitracin-polymyxin B (POLYSPORIN) ointment** Dose: 500-10000 Route: Topical Frequency: 3 times daily

Admin Instructions:

Apply to drain site.

☐ **neomycin-bacitracin-polymyxinB (NEOSPORIN) ointment** Dose: 3.5mg-400 unit- Route: Topical Frequency: 3 times daily

Admin Instructions:

Apply to drain site.

☐ **mupirocin (BACTROBAN) 2 % ointment** Dose: 2 Route: Topical Frequency: 3 times daily

Admin Instructions:

Apply to drain site.

☐ **povidone-iodine (BETADINE) ointment** Dose: 10 Route: Topical Frequency: 3 times daily

Admin Instructions:

Apply to drain site.

Ophthalmic Antibiotic Ointments

☐ **tobramycin-dexamethasone (TOBRADEX) ophthalmic ointment** Dose: 0.3-0.1 Frequency: 3 times daily

Product Admin Instructions:

For Ophthalmic use only

Facial Operations

☐ **chlorhexidine (PERIDEX) 0.12 % solution** Dose: 15 mL Route: Mouth/Throat Frequency: 2 times daily

Admin Instructions:

Swish and Spit

☐ **artificial tears ophthalmic solution** Dose: 2 drop Route: Both Eyes Frequency: every 4 hours PRN PRN Reasons: dry eyes

Product Admin Instructions:

For Ophthalmic use only

☐ **artificial tears ointment** Route: Both Eyes Frequency: nightly PRN PRN Reasons: dry eyes

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **clonIDINE HCl (CATAPRES) tablet** Dose: .1 Route: oral Frequency: 2 times daily PRN PRN Reasons: high blood pressure

Question(s):

BP & HR HOLD parameters for this order:

Contact Physician if:

Bowel Care - NOT HMSJ

☐ **docusate sodium (COLACE) capsule** Dose: 100 mg Route: oral Frequency: 2 times daily PRN PRN Reasons: constipation

Admin Instructions:

Use docusate for stool softener as needed.

☐ **simethicone (MYLICON) chewable tablet** Dose: 160 mg Route: oral Frequency: 4 times daily PRN PRN Reasons: flatulence

☐ **bisacodyl (DULCOLAX) suppository** Dose: 10 mg Route: rectal Frequency: daily PRN PRN Reasons: constipation

Admin Instructions:

Suppository can be used if oral therapy is not tolerated or ineffective.

☐ **senna (SENOKOT) tablet** Dose: 1 tablet Route: oral Frequency: 2 times daily PRN PRN Reasons: constipation

☐ **diphenoxylate-atropine (LOMOTIL) 2.5-0.025 mg per tablet** Dose: 1 tablet Route: oral Frequency: 4 times daily PRN PRN Reasons: diarrhea

Anxiolytics: For Patients LESS than 65 years old

☐ **LORAZepam (ATIVAN) tablet** Dose: 1 mg Route: oral Frequency: every 6 hours PRN PRN Reasons: anxiety

Question(s):Indication(s): ☐ Anxiety**Anxiolytics: For Patients GREATER than or EQUAL to 65 years old**

☐ **LORAZepam (ATIVAN) tablet** Dose: 0.5 mg Route: oral Frequency: every 6 hours PRN PRN Reasons: anxiety

Question(s):Indication(s): ☐ Anxiety**Muscle Spasms****Caution: muscle relaxants should be minimized in patients over 65 years of age.**

☐ **cyclobenzaprine (FLEXERIL) tablet** Dose: 5 mg Route: oral Frequency: every 8 hours PRN PRN Reasons: muscle spasms

☐ **methocarbamol (ROBAXIN) tablet** Dose: 500 mg Route: oral Frequency: 3 times daily PRN PRN Reasons: muscle spasms

Muscle Pain

☐ **diazepam (VALIUM) tablet** Dose: 5 mg Route: oral Frequency: every 6 hours PRN PRN Comment: muscle pain

Question(s):Indication(s): ☐ Other

Specify: Muscle Pain

On-Q Pump

☐ **ropivacaine 0.2% (PF) (NAROPIN) solution for On-Q Pump** Route: infiltration Frequency: continuous

Question(s):

Regional Block:

Location:

Catheter:

Continuous Rate:

Bolus Dose (Optional):

Product Admin Instructions:

This is a disposable, single-use product for Regional Block use. Do NOT refill

☐ **ropivacaine 0.5% (PF) (NAROPIN) solution for On-Q Pump** Route: infiltration Frequency: continuous

Question(s):

Regional Block:

Location:

Catheter:

Continuous Rate:

Bolus Dose (Optional):

Product Admin Instructions:

This is a disposable, single-use product for Regional Block use. Do NOT refill

PCA Medications

☐ **morPHINE PCA 30 mg/30 mL**

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **morPHINE PCA 30 mg/30 mL Dose:** 3 **Route:** intravenous **Frequency:** continuous

Admin Instructions:

Management of breakthrough pain. Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber. For breakthrough pain in patients ages 19-59 years old with normal renal function, may bolus {Bolus Dose:26657::"2"} mg every {Bolus Frequency:26659::"3"} hours as needed. If pain persists, may increase PCA demand dose by {PCA Dose:26660::"0.5"} mg ONCE. Adjust doses for age, renal function or other factors.

☒ **Vital signs - T/P/R/BP Frequency:** Per unit protocol **Priority:** Routine **Comments:** - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change

☒ **Richmond agitation sedation scale Frequency:** Once **Priority:** Routine **Comments:** 60 minutes after administration of pain medication AND every 4 hours. Assess and document side effects of at least every 4 hours for duration of therapy and when patient complains of pain and/or side effects.

Question(s):

Hold infusion daily at:

Target RASS:

BIS Monitoring (Target BIS: 40-60):

☒ **Notify Physician (Specify) Frequency:** Until discontinued **Priority:** Routine **Comments:** - PCA pump infusion discontinued for any reason - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy

☒ **Stop the PCA pump and call ordering physician and/or CERT team for any of the following: Frequency:** Until discontinued **Priority:** Routine **Comments:** - Respiratory rate 10 per minute or less - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention

☒ **naloxone (NARCAN) 0.4 mg/mL injection 0.2 mg Dose:** 0.2 mg **Frequency:** once PRN **Priority:** Routine **PRN Comment:** as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3).

Admin Instructions:

Repeat Naloxone 0.2 mg once in 2 minutes if necessary (MAXIMUM 0.4 mg). If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times.

☐ **hydromorPHONE PCA (DILAUDID) 15 mg/30 mL**

☒ **hydromorPHONE (DILAUDID) 15 mg/30 mL PCA Dose:** .5 **Route:** intravenous **Frequency:** continuous

Admin Instructions:

Management of breakthrough pain. Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber. For breakthrough pain in patients ages 19-59 years old with normal renal function, may bolus {Bolus Dose:26662::"0.2"} mg every {Bolus Frequency:26663::"3"} hours as needed. If pain persists, may increase PCA demand dose by {PCA Dose:26664::"0.1"} mg ONCE. Adjust doses for age, renal function or other factors.

☒ **Vital signs - T/P/R/BP Frequency:** Per unit protocol **Priority:** Routine **Comments:** - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change

☒ **Richmond agitation sedation scale Frequency:** Once **Priority:** Routine **Comments:** 60 minutes after administration of pain medication AND every 4 hours. Assess and document side effects of at least every 4 hours for duration of therapy and when patient complains of pain and/or side effects.

Question(s):

Hold infusion daily at:

Target RASS:

BIS Monitoring (Target BIS: 40-60):

☒ **Notify Physician (Specify) Frequency:** Until discontinued **Priority:** Routine **Comments:** - PCA pump infusion discontinued for any reason - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy

☒ **Stop the PCA pump and call ordering physician and/or CERT team for any of the following: Frequency:** Until discontinued **Priority:** Routine **Comments:** - Respiratory rate 10 per minute or less - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **naloxone (NARCAN) 0.4 mg/mL injection 0.2 mg** Dose: 0.2 mg Frequency: once PRN Priority: Routine PRN

Comment: as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3).

Admin Instructions:

Repeat Naloxone 0.2 mg once in 2 minutes if necessary (MAXIMUM 0.4 mg). If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times.

☐ **fentaNYL PCA (SUBLIMAZE) 1500 mcg/30 mL**

☒ **fentaNYL (SUBLIMAZE) 1500 mcg/30 mL PCA** Dose: 50 Route: intravenous Frequency: continuous

Admin Instructions:

Management of breakthrough pain. Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber. For breakthrough pain in patient 19-59 years old, may bolus {Bolus Dose:26656::"25"} mcg every {Bolus Frequency:26655::"2"} hours as needed. If pain persists, may increase PCA demand dose by {PCA Dose:26654::"10"} mcg ONCE. Adjust doses for age, renal function or other factors.

☒ **Vital signs - T/P/R/BP** Frequency: Per unit protocol Priority: Routine Comments: - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change

☒ **Richmond agitation sedation scale** Frequency: Once Priority: Routine Comments: 60 minutes after administration of pain medication AND every 4 hours. Assess and document side effects of at least every 4 hours for duration of therapy and when patient complains of pain and/or side effects.

Question(s):

Hold infusion daily at:

Target RASS:

BIS Monitoring (Target BIS: 40-60):

☒ **Notify Physician (Specify)** Frequency: Until discontinued Priority: Routine Comments: - PCA pump infusion discontinued for any reason - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy

☒ **Stop the PCA pump and call ordering physician and/or CERT team for any of the following:** Frequency: Until discontinued Priority: Routine Comments: - Respiratory rate 10 per minute or less - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention

☒ **naloxone (NARCAN) 0.4 mg/mL injection 0.2 mg** Dose: 0.2 mg Frequency: once PRN Priority: Routine PRN

Comment: as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3).

Admin Instructions:

Repeat Naloxone 0.2 mg once in 2 minutes if necessary (MAXIMUM 0.4 mg). If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times.

PCA Medications

☐ **morPHINE PCA 30 mg/30 mL**

☒ **morPHINE PCA 30 mg/30 mL** Dose: 3 Route: intravenous Frequency: continuous

Admin Instructions:

Management of breakthrough pain. Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber. For breakthrough pain in patients ages 19-59 years old with normal renal function, may bolus {Bolus Dose:26657::"2"} mg every {Bolus Frequency:26659::"3"} hours as needed. If pain persists, may increase PCA demand dose by {PCA Dose:26660::"0.5"} mg ONCE. Adjust doses for age, renal function or other factors.

☒ **Vital signs - T/P/R/BP** Frequency: Per unit protocol Priority: Routine Comments: - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change

☒ **Richmond agitation sedation scale** Frequency: Once Priority: Routine Comments: 60 minutes after administration of pain medication AND every 4 hours. Assess and document side effects of at least every 4 hours for duration of therapy and when patient complains of pain and/or side effects.

Question(s):

Hold infusion daily at:

Target RASS:

BIS Monitoring (Target BIS: 40-60):

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **Notify Physician (Specify) Frequency:** Until discontinued **Priority:** Routine **Comments:** - PCA pump infusion discontinued for any reason - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy

☒ **Stop the PCA pump and call ordering physician and/or CERT team for any of the following: Frequency:** Until discontinued **Priority:** Routine **Comments:** - Respiratory rate 10 per minute or less - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention

☒ **naloxone (NARCAN) 0.4 mg/mL injection 0.2 mg Dose:** 0.2 mg **Frequency:** once PRN **Priority:** Routine **PRN Comment:** as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3).

Admin Instructions:

Repeat Naloxone 0.2 mg once in 2 minutes if necessary (MAXIMUM 0.4 mg). If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times.

☐ **hydromorPHONE PCA (DILAUDID) 15 mg/30 mL**

☒ **hydromorPHONE (DILAUDID) 15 mg/30 mL PCA Dose:** .5 **Route:** intravenous **Frequency:** continuous

Admin Instructions:

Management of breakthrough pain. Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber. For breakthrough pain in patients ages 19-59 years old with normal renal function, may bolus {Bolus Dose:26662::"0.2"} mg every {Bolus Frequency:26663::"3"} hours as needed. If pain persists, may increase PCA demand dose by {PCA Dose:26664::"0.1"} mg ONCE. Adjust doses for age, renal function or other factors.

☒ **Vital signs - T/P/R/BP Frequency:** Per unit protocol **Priority:** Routine **Comments:** - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change

☒ **Richmond agitation sedation scale Frequency:** Once **Priority:** Routine **Comments:** 60 minutes after administration of pain medication AND every 4 hours. Assess and document side effects of at least every 4 hours for duration of therapy and when patient complains of pain and/or side effects.

Question(s):

Hold infusion daily at:

Target RASS:

BIS Monitoring (Target BIS: 40-60):

☒ **Notify Physician (Specify) Frequency:** Until discontinued **Priority:** Routine **Comments:** - PCA pump infusion discontinued for any reason - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy

☒ **Stop the PCA pump and call ordering physician and/or CERT team for any of the following: Frequency:** Until discontinued **Priority:** Routine **Comments:** - Respiratory rate 10 per minute or less - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention

☒ **naloxone (NARCAN) 0.4 mg/mL injection 0.2 mg Dose:** 0.2 mg **Frequency:** once PRN **Priority:** Routine **PRN Comment:** as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3).

Admin Instructions:

Repeat Naloxone 0.2 mg once in 2 minutes if necessary (MAXIMUM 0.4 mg). If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times.

☐ **fentaNYL PCA (SUBLIMAZE) 600 mcg/30 mL**

☒ **fentaNYL (SUBLIMAZE) 600 mcg/30 mL PCA Dose:** 20 **Route:** intravenous **Frequency:** continuous

Admin Instructions:

Management of breakthrough pain. Administer only if respiratory rate 12 per minute or more and POSS level of 2 or less. If more than 2 bolus doses in 12 hours or if pain persists after increase in demand dose, call ordering prescriber. For breakthrough pain in patient 19-59 years old, may bolus {Bolus Dose:26656::"25"} mcg every {Bolus Frequency:26655::"2"} hours as needed. If pain persists, may increase PCA demand dose by {PCA Dose:26654::"10"} mcg ONCE. Adjust doses for age, renal function or other factors.

☒ **Vital signs - T/P/R/BP Frequency:** Per unit protocol **Priority:** Routine **Comments:** - Initially and every 30 minutes for 1 hour after PCA started, bolus administration or dose change; then - Every hour x 2 starting second hour after PCA started, bolus administered or dose change; then - Every 4 hours until PCA therapy is discontinued. - Immediately following PCA administration tubing change

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **Richmond agitation sedation scale** **Frequency:** Once **Priority:** Routine **Comments:** 60 minutes after administration of pain medication AND every 4 hours. Assess and document side effects of at least every 4 hours for duration of therapy and when patient complains of pain and/or side effects.

Question(s):

Hold infusion daily at:

Target RASS:

BIS Monitoring (Target BIS: 40-60):

☒ **Notify Physician (Specify)** **Frequency:** Until discontinued **Priority:** Routine **Comments:** - PCA pump infusion discontinued for any reason - Inadequate analgesia - Prior to administration of any other narcotics, antiemetics, or sedatives other than those ordered by the prescriber responsible for IV PCA therapy - PCA pump discontinued by any service other than the prescriber responsible for IV PCA therapy

☒ **Stop the PCA pump and call ordering physician and/or CERT team for any of the following:** **Frequency:** Until discontinued **Priority:** Routine **Comments:** - Respiratory rate 10 per minute or less - Severe and/or recent confusion or disorientation - POSS sedation level 4: Somnolent and difficult to arouse - Sustained hypotension (SBP less than 90) - Excessive nausea or vomiting - Urinary retention

☒ **naloxone (NARCAN) 0.4 mg/mL injection 0.2 mg** **Dose:** 0.2 mg **Frequency:** once PRN **Priority:** Routine **PRN Comment:** as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3).

Admin Instructions:

Repeat Naloxone 0.2 mg once in 2 minutes if necessary (MAXIMUM 0.4 mg). If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times.

Mild Pain (Pain Score 1-3) or Fever

☐ **acetaminophen (TYLENOL) tablet** **Dose:** 650 mg **Route:** oral **Frequency:** every 6 hours PRN **PRN Reasons:** mild pain (score 1-3)
fever

Question(s):

Allowance for Patient Preference:

Admin Instructions:

Contact physician for fever GREATER than 101 F

Product Admin Instructions:

Maximum of 3 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources).

Oral for Moderate Pain (Pain Score 4-6)

☐ **HYDROcodone-acetaminophen (NORCO) 5-325 mg per tablet** **Dose:** 1 tablet **Route:** oral **Frequency:** every 4 hours PRN **PRN Reasons:** moderate pain (score 4-6)

Question(s):

Allowance for Patient Preference:

Product Admin Instructions:

Give if patient can receive oral tablet/capsule.

☐ **HYDROcodone-acetaminophen (NORCO) 7.5-325 mg per tablet** **Dose:** 1 tablet **Route:** oral **Frequency:** every 4 hours PRN **PRN Reasons:** moderate pain (score 4-6)

Question(s):

Allowance for Patient Preference:

Product Admin Instructions:

Give if patient can receive oral tablet/capsule.

☐ **traMADol (ULTRAM) tablet** **Dose:** 50 mg **Route:** oral **Frequency:** every 6 hours PRN **PRN Reasons:** moderate pain (score 4-6)

Question(s):

Allowance for Patient Preference:

Product Admin Instructions:

Give if patient can receive oral tablet/capsule.

☐ **oxyCODONE-acetaminophen (PERCOCET) 5-325 mg per tablet** **Dose:** 1 tablet **Route:** oral **Frequency:** every 4 hours PRN **PRN Reasons:** moderate pain (score 4-6)

Question(s):

Allowance for Patient Preference:

Product Admin Instructions:

Give if patient can receive oral tablet/capsule. Maximum of 4 grams of acetaminophen per day from all sources. (Cirrhosis patients maximum: 2 grams per day from all sources).

IV for Moderate Pain (Pain Score 4-6)

If you select a PCA option you will not be allowed to also order IV PRN pain medications from this section.

Sign: _____ Printed Name: _____ Date/Time: _____

- ☐ **morPHINE injection** Dose: 1 mg Route: intravenous Frequency: every 4 hours PRN PRN Reasons: moderate pain (score 4-6)

Admin Instructions:

Give if patient cannot tolerate oral medications or a faster onset of action is required.

Oral for Severe Pain (Pain Score 7-10)

- ☐ **HYDROcodone-acetaminophen (NORCO 10-325) 10-325 mg per tablet** Dose: 1 tablet Route: oral Frequency: every 4 hours PRN PRN Reasons: severe pain (score 7-10)

Question(s):

Allowance for Patient Preference:

- ☐ **traMADol (ULTRAM) tablet** Dose: 100 mg Route: oral Frequency: every 6 hours PRN PRN Reasons: severe pain (score 7-10)

Question(s):

Allowance for Patient Preference:

Product Admin Instructions:

Give if patient can receive oral tablet/capsule.

- ☐ **oxyCODone-acetaminophen (PERCOCET) 10-325 mg per tablet** Dose: 1 tablet Route: oral Frequency: every 4 hours PRN PRN Reasons: severe pain (score 7-10)

Question(s):

Allowance for Patient Preference:

Product Admin Instructions:

Give if patient can receive oral tablet/capsule.

IV for Severe Pain (Pain Score 7-10)

If you select a PCA option you will not be allowed to also order IV PRN pain medications from this section.

- ☐ **morPHINE injection** Dose: 2 mg Route: intravenous Frequency: every 4 hours PRN PRN Reasons: severe pain (score 7-10)

Admin Instructions:

Give if patient cannot tolerate oral medications or a faster onset of action is required.

Respiratory

- ☒ **naloxone (NARCAN) injection** Dose: 0.2 mg Route: intravenous Frequency: once PRN Priority: Routine PRN Comment: as needed for respiratory rate 8 per minute or less OR patient somnolent and difficult to arouse (POSS GREATER than 3). PRN Reasons: respiratory depression

Admin Instructions:

Repeat Naloxone 0.2 mg once in 2 minutes if necessary (MAXIMUM 0.4 mg).

If naloxone is needed, please call the ordering physician and/or CERT team. Monitor vital signs (pulse oximetry, P/R/BP) every 15 minutes for 3 times.

Bowel Care - NOT HMSJ

- ☐ **docusate sodium (COLACE) capsule** Dose: 100 mg Route: oral Frequency: 2 times daily PRN PRN Reasons: constipation

Admin Instructions:

Use docusate for stool softener as needed.

- ☐ **simethicone (MYLICON) chewable tablet** Dose: 160 mg Route: oral Frequency: 4 times daily PRN PRN Reasons: flatulence

- ☐ **bisacodyl (DULCOLAX) suppository** Dose: 10 mg Route: rectal Frequency: daily PRN PRN Reasons: constipation

Admin Instructions:

Suppository can be used if oral therapy is not tolerated or ineffective.

- ☐ **senna (SENOKOT) tablet** Dose: 1 tablet Route: oral Frequency: 2 times daily PRN PRN Reasons: constipation

- ☐ **diphenoxylate-atropine (LOMOTIL) 2.5-0.025 mg per tablet** Dose: 1 tablet Route: oral Frequency: 4 times daily PRN PRN Reasons: diarrhea

Laxatives - HMSJ Only

- ☐ **docusate sodium (COLACE) capsule** Dose: 100 mg Route: oral Frequency: 2 times daily PRN PRN Reasons: constipation

Admin Instructions:

Use docusate for stool softener as needed.

- ☐ **simethicone (MYLICON) chewable tablet** Dose: 160 mg Route: oral Frequency: 4 times daily PRN PRN Reasons: flatulence

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **bisacodyl (DULCOLAX) suppository** Dose: 10 mg Route: rectal Frequency: daily PRN PRN Reasons: constipation

Admin Instructions:

Suppository can be used if oral therapy is not tolerated or ineffective.

☐ **sennosides-docusate sodium (SENOKOT-S) 8.6-50 mg per tablet** Dose: 1 tablet Route: oral Frequency: 2 times daily PRN PRN Reasons: constipation

☐ **diphenoxylate-atropine (LOMOTIL) 2.5-0.025 mg per tablet** Dose: 1 tablet Route: oral Frequency: 4 times daily PRN PRN Reasons: diarrhea

Antiemetics

☒ **ondansetron (ZOFTRAN) IV or Oral (Required)**

☒ **ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet** Dose: 4 mg Route: oral Frequency: every 8 hours PRN

PRN Reasons: nausea

vomiting

Admin Instructions:

Give if patient is able to tolerate oral medication.

Product Admin Instructions:

May cause QTc prolongation.

☒ **ondansetron (ZOFTRAN) 4 mg/2 mL injection** Dose: 4 mg Route: intravenous Frequency: every 8 hours PRN PRN

Reasons: nausea

vomiting

Admin Instructions:

Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.

Product Admin Instructions:

May cause QTc prolongation.

☐ **promethazine (PHENERGAN)**

☒ **promethazine (PHENERGAN) 12.5 mg IV** Dose: 12.5 mg Route: intravenous Frequency: every 6 hours PRN PRN

Reasons: nausea

vomiting

Admin Instructions:

Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.

☒ **promethazine (PHENERGAN) tablet** Dose: 12.5 mg Route: oral Frequency: every 6 hours PRN PRN Reasons:

nausea

vomiting

Admin Instructions:

Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

☒ **promethazine (PHENERGAN) suppository** Dose: 12.5 mg Route: rectal Frequency: every 6 hours PRN PRN

Reasons: nausea

vomiting

Admin Instructions:

Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

☒ **promethazine (PHENERGAN) intraMUSCULAR injection** Dose: 12.5 mg Route: intramuscular Frequency: every 6

hours PRN PRN Reasons: nausea

vomiting

Admin Instructions:

Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics

☒ **ondansetron (ZOFTRAN) IV or Oral (Required)**

☒ **ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet** Dose: 4 mg Route: oral Frequency: every 8 hours PRN

PRN Reasons: nausea

vomiting

Admin Instructions:

Give if patient is able to tolerate oral medication.

Product Admin Instructions:

May cause QTc prolongation.

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **ondansetron (ZOFTRAN) 4 mg/2 mL injection** Dose: 4 mg Route: intravenous Frequency: every 8 hours PRN PRN

Reasons: nausea
vomiting

Admin Instructions:

Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.

Product Admin Instructions:

May cause QTc prolongation.

☐ **promethazine (PHENERGAN) IV or Oral or Rectal**

☒ **promethazine (PHENERGAN) injection** Dose: 12.5 mg Route: intravenous Frequency: every 6 hours PRN PRN

Reasons: nausea
vomiting

Admin Instructions:

Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.

☒ **promethazine (PHENERGAN) tablet** Dose: 12.5 mg Route: oral Frequency: every 6 hours PRN PRN Reasons:

nausea
vomiting

Admin Instructions:

Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.

☒ **promethazine (PHENERGAN) suppository** Dose: 12.5 mg Route: rectal Frequency: every 6 hours PRN PRN

Reasons: nausea
vomiting

Admin Instructions:

Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Antiemetics

☒ **ondansetron (ZOFTRAN) IV or Oral (Required)**

☒ **ondansetron ODT (ZOFTRAN-ODT) disintegrating tablet** Dose: 4 mg Route: oral Frequency: every 8 hours PRN PRN

Reasons: nausea
vomiting

Admin Instructions:

Give if patient is able to tolerate oral medication.

Product Admin Instructions:

May cause QTc prolongation.

☒ **ondansetron (ZOFTRAN) 4 mg/2 mL injection** Dose: 4 mg Route: intravenous Frequency: every 8 hours PRN PRN

Reasons: nausea
vomiting

Admin Instructions:

Give if patient is UNable to tolerate oral medication OR if a faster onset of action is required.

Product Admin Instructions:

May cause QTc prolongation.

☐ **promethazine (PHENERGAN) IVPB or Oral or Rectal**

☒ **promethazine (PHENERGAN) 25 mg in sodium chloride 0.9 % 50 mL IVPB** Dose: 12.5 mg Route: intravenous

Frequency: every 6 hours PRN Minimum Infusion Duration: 30.000 Minutes PRN Reasons: nausea
vomiting

Admin Instructions:

Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral or rectal medication OR if a faster onset of action is required.

☒ **promethazine (PHENERGAN) tablet** Dose: 12.5 mg Route: oral Frequency: every 6 hours PRN PRN Reasons:

nausea
vomiting

Admin Instructions:

Give if ondansetron (ZOFTRAN) is ineffective and patient is able to tolerate oral medication.

☒ **promethazine (PHENERGAN) suppository** Dose: 12.5 mg Route: rectal Frequency: every 6 hours PRN PRN

Reasons: nausea
vomiting

Admin Instructions:

Give if ondansetron (ZOFTRAN) is ineffective and patient is UNable to tolerate oral medication.

Itching: For Patients GREATER than 70 years old

☒ **cetirizine (Zyrtec) tablet** Dose: 5 mg Route: oral Frequency: daily PRN PRN Reasons: itching

Sign: _____ Printed Name: _____ Date/Time: _____

HM IP MEDICATIONS - COMMON PRN - ITCHING FOR PTS 70-76 YO

- ☐ **cetirizine (Zyrtec) tablet** Dose: 5 mg Route: oral Frequency: daily PRN PRN Reasons: itching

Itching: For Patients LESS than 70 years old

- ☐ **diphenhydramine (BENADRYL) tablet** Dose: 25 mg Route: oral Frequency: every 6 hours PRN PRN Reasons: itching
- ☐ **hydroxyzine (ATARAX) tablet** Dose: 10 mg Route: oral Frequency: every 6 hours PRN PRN Reasons: itching
- ☒ **cetirizine (Zyrtec) tablet** Dose: 5 mg Route: oral Frequency: daily PRN PRN Reasons: itching
- ☐ **fexofenadine (ALLEGRA) tablet - For eGFR LESS than 80 mL/min, reduce frequency to once daily as needed** Dose: 60 mg Route: oral Frequency: 2 times daily PRN PRN Reasons: itching

Insomnia: For Patients GREATER than or EQUAL to 70 years old

- ☐ **ramelteon (ROZEREM) tablet** Dose: 8 mg Route: oral Frequency: nightly PRN PRN Reasons: sleep

Insomnia: For Patients LESS than 70 years old

- ☐ **zolpidem (AMBIEN) or ramelteon (ROZEREM) tablet** nightly PRN sleep
- ☐ **zolpidem (AMBIEN) tablet** Dose: 5 mg Route: oral Frequency: nightly PRN PRN Reasons: sleep
- ☐ **ramelteon (ROZEREM) tablet** Dose: 8 mg Route: oral Frequency: nightly PRN PRN Reasons: sleep

- ☐ **sodium chloride 0.9% bag** for line care

- ☒ **sodium chloride 0.9 % bag for line care** Dose: .9 Frequency: PRN PRN Reasons: line care

Admin Instructions:

For flushing of extension tubing sets after administration of intermittent infusions. Program sodium chloride bag to run at the same infusion rate as medication given for a total volume equal to contents of tubing sets used. Change bag every 24 hours.

VTE

DVT Risk and Prophylaxis Tool 1

1

VTE/DVT Risk Definitions (<http://epic-nas.et0922.epichosted.com/static/OrderSets/VTEDVTRISKDEFINITIONS.pdf>)

- ☐ Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Required)
- ☐ Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)
- ☒ **Moderate risk of VTE** Frequency: Once Priority: Routine
- ☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once Priority: Routine
- Question(s):**
- No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
- ☒ **Place sequential compression device**
- ☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine
- Question(s):**
- No mechanical VTE prophylaxis due to the following contraindication(s):
- ☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine
- Question(s):**
- Side: Bilateral
- Select Sleeve(s):
- ☐ Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)
- ☒ **Moderate risk of VTE** Frequency: Once Priority: Routine
- ☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once Priority: Routine
- Question(s):**
- No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:
- ☒ **Place sequential compression device**
- ☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine
- Question(s):**
- No mechanical VTE prophylaxis due to the following contraindication(s):

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** (Required)

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once

Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.

Therapy for the following:

☒ **Place sequential compression device**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** (Required)

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once

Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.

Therapy for the following:

☒ **Place sequential compression device**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **LOW Risk of VTE** (Required)

☒ **Low Risk** (Required)

☒ **Low risk of VTE** Frequency: Once Priority: Routine

Question(s):

Low risk: ☐ Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation ☐ Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation

☐ **MODERATE Risk of VTE - Surgical** (Required)

☒ **Moderate Risk** (Required)

☒ **Moderate risk of VTE** Frequency: Once Priority: Routine

☒ **Moderate Risk Pharmacological Prophylaxis - Surgical Patient** (Required)

☐ **Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device**

☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis**

☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation** (Required)

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily Start Date: S+1

Admin Instructions:

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

☐ **heparin**

Sign: _____ Printed Name: _____ Date/Time: _____

High Risk Bleeding Characteristics
Age $\geq$ 75
Weight < 50 kg
Unstable Hgb
Renal impairment
Plt count < 100 K/uL
Dual antiplatelet therapy
Active cancer
Cirrhosis/hepatic failure
Prior intra-cranial hemorrhage
Prior ischemic stroke
History of bleeding event requiring admission and/or transfusion
Chronic use of NSAIDs/steroids
Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours Dose:** 5000 Units **Frequency:** every 12 hours scheduled

☐ **HEParin (porcine) injection - Q8 Hours Dose:** 5000 Units **Frequency:** every 8 hours scheduled

☐ **Not high bleed risk**

☐ **Wt > 100 kg Dose:** 7500 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **Wt LESS than or equal to 100 kg Dose:** 5000 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **warfarin (COUMADIN)**

☐ **WITHOUT pharmacy consult Dose:** 1 **Route:** oral **Frequency:** daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

☒ **Pharmacy consult to manage warfarin (COUMADIN) Frequency:** Until discontinued **Priority:**

Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet Dose:** 1 **Route:** oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis (Required)**

☐ **Contraindications exist for mechanical prophylaxis Frequency:** Once **Priority:** Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous Frequency:** Continuous **Priority:** Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **MODERATE Risk of VTE - Non-Surgical (Required)**

☒ **Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Required)**

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **Moderate Risk (Required)**

☒ **Moderate risk of VTE Frequency:** Once **Priority:** Routine

☒ **Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Required)**

☐ **Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device**

☒ **Contraindications exist for pharmacologic prophylaxis Frequency:** Once **Priority:** Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous Frequency:** Continuous **Priority:** Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis**

☒ **Contraindications exist for pharmacologic prophylaxis Frequency:** Once **Priority:** Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Contraindications exist for mechanical prophylaxis Frequency:** Once **Priority:** Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin for Prophylactic Anticoagulation Nonsurgical (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection Dose:** 30 mg **Route:** subcutaneous **Frequency:** daily at 1700 **Start Date:** S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection Route:** subcutaneous **Start Date:** S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

Sign: _____ **Printed Name:** _____ **Date/Time:** _____

- ☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily

Admin Instructions:

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min

- ☐ **heparin**

High Risk Bleeding Characteristics
Age $\geq$ 75
Weight < 50 kg
Unstable Hgb
Renal impairment
Plt count < 100 K/uL
Dual antiplatelet therapy
Active cancer
Cirrhosis/hepatic failure
Prior intra-cranial hemorrhage
Prior ischemic stroke
History of bleeding event requiring admission and/or transfusion
Chronic use of NSAIDs/steroids
Active GI ulcer

- ☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

- ☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units Frequency: every 12 hours scheduled

- ☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units Frequency: every 8 hours scheduled

- ☐ **Not high bleed risk**

- ☐ **Wt > 100 kg** Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled

- ☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled

- ☐ **warfarin (COUMADIN)**

- ☐ **WITHOUT pharmacy consult** Dose: 1 Route: oral Frequency: daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

- ☐ **Medications**

- ☒ **Pharmacy consult to manage warfarin (COUMADIN)** Frequency: Until discontinued

Priority: Routine

Question(s):

Indication:

- ☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

- ☐ **Mechanical Prophylaxis** (Required)

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **HIGH Risk of VTE - Surgical** (Required)

☒ **High Risk** (Required)

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **High Risk Pharmacological Prophylaxis - Surgical Patient** (Required)

☐ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation** (Required)

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily Start Date: S+1

Admin Instructions:

If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

☐ **heparin**

Sign: _____ Printed Name: _____ Date/Time: _____

High Risk Bleeding Characteristics
Age $\geq$ 75
Weight < 50 kg
Unstable Hgb
Renal impairment
Plt count < 100 K/uL
Dual antiplatelet therapy
Active cancer
Cirrhosis/hepatic failure
Prior intra-cranial hemorrhage
Prior ischemic stroke
History of bleeding event requiring admission and/or transfusion
Chronic use of NSAIDs/steroids
Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours Dose:** 5000 Units **Frequency:** every 12 hours scheduled

☐ **HEParin (porcine) injection - Q8 Hours Dose:** 5000 Units **Frequency:** every 8 hours scheduled

☐ **Not high bleed risk**

☐ **Wt > 100 kg Dose:** 7500 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **Wt LESS than or equal to 100 kg Dose:** 5000 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **warfarin (COUMADIN)**

☐ **WITHOUT pharmacy consult Dose:** 1 **Route:** oral **Frequency:** daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

☒ **Pharmacy consult to manage warfarin (COUMADIN) Frequency:** Until discontinued **Priority:**

Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet Dose:** 1 **Route:** oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis (Required)**

☐ **Contraindications exist for mechanical prophylaxis Frequency:** Once **Priority:** Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous Frequency:** Continuous **Priority:** Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **HIGH Risk of VTE - Non-Surgical (Required)**

☒ **High Risk (Required)**

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **High Risk Pharmacological Prophylaxis - Non-Surgical Patient** (Required)

☐ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin for Prophylactic Anticoagulation Nonsurgical** (Required)

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily

Admin Instructions:

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

☐ **heparin**

Sign: _____ Printed Name: _____ Date/Time: _____

High Risk Bleeding Characteristics
Age $\geq$ 75
Weight < 50 kg
Unstable Hgb
Renal impairment
Plt count < 100 K/uL
Dual antiplatelet therapy
Active cancer
Cirrhosis/hepatic failure
Prior intra-cranial hemorrhage
Prior ischemic stroke
History of bleeding event requiring admission and/or transfusion
Chronic use of NSAIDs/steroids
Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours Dose:** 5000 Units **Frequency:** every 12 hours scheduled

☐ **HEParin (porcine) injection - Q8 Hours Dose:** 5000 Units **Frequency:** every 8 hours scheduled

☐ **Not high bleed risk**

☐ **Wt > 100 kg Dose:** 7500 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **Wt LESS than or equal to 100 kg Dose:** 5000 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **warfarin (COUMADIN)**

☐ **WITHOUT pharmacy consult Dose:** 1 **Route:** oral **Frequency:** daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

☒ **Pharmacy consult to manage warfarin (COUMADIN) Frequency:** Until discontinued **Priority:**

Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet Dose:** 1 **Route:** oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis (Required)**

☐ **Contraindications exist for mechanical prophylaxis Frequency:** Once **Priority:** Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous Frequency:** Continuous **Priority:** Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **HIGH Risk of VTE - Surgical (Hip/Knee) (Required)**

☒ **High Risk (Required)**

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **High risk of VTE Frequency:** Once **Priority:** Routine

☒ **High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Required)**

☐ **Contraindications exist for pharmacologic prophylaxis Frequency:** Once **Priority:** Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☐ **aspirin chewable tablet Dose:** 162 mg **Frequency:** daily **Start Date:** S+1

☐ **aspirin (ECOTRIN) enteric coated tablet Dose:** 162 mg **Frequency:** daily **Start Date:** S+1

☐ **Apixaban and Pharmacy Consult (Required)**

☒ **apixaban (ELIQUIS) tablet Dose:** 2.5 mg **Frequency:** 2 times daily **Start Date:** S+1

Question(s):

Indications: ☐ VTE prophylaxis

☒ **Pharmacy consult to monitor apixaban (ELIQUIS) therapy Frequency:** Until discontinued **Priority:**

STAT

Question(s):

Indications: VTE prophylaxis

☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection Dose:** 30 mg **Route:** subcutaneous **Frequency:** daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection Route:** subcutaneous **Start Date:** S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection Dose:** 2.5 mg **Route:** subcutaneous **Frequency:** daily **Start Date:** S+1

Admin Instructions:

If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min

☐ **heparin**

Sign: _____ **Printed Name:** _____ **Date/Time:** _____

High Risk Bleeding Characteristics
Age $\geq$ 75
Weight < 50 kg
Unstable Hgb
Renal impairment
Plt count < 100 K/uL
Dual antiplatelet therapy
Active cancer
Cirrhosis/hepatic failure
Prior intra-cranial hemorrhage
Prior ischemic stroke
History of bleeding event requiring admission and/or transfusion
Chronic use of NSAIDs/steroids
Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours Dose:** 5000 Units **Frequency:** every 12 hours scheduled

☐ **HEParin (porcine) injection - Q8 Hours Dose:** 5000 Units **Frequency:** every 8 hours scheduled

☐ **Not high bleed risk**

☐ **Wt > 100 kg Dose:** 7500 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **Wt LESS than or equal to 100 kg Dose:** 5000 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **Rivaroxaban and Pharmacy Consult (Required)**

☒ **rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission Dose:** 10 mg **Frequency:** daily at 0600 (TIME CRITICAL)

Question(s):

Indications: ☐ VTE prophylaxis

Admin Instructions:

For Xarelto 15 mg and 20 mg, give with food or follow administration with enteral feeding to increase medication absorption. Do not administer via post-pyloric routes.

☒ **Pharmacy consult to monitor rivaroxaban (XARELTO) therapy Frequency:** Until discontinued **Priority:** STAT

Question(s):

Indications: VTE prophylaxis

☐ **warfarin (COUMADIN)**

☐ **WITHOUT pharmacy consult Dose:** 1 **Route:** oral **Frequency:** daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

☒ **Pharmacy consult to manage warfarin (COUMADIN) Frequency:** Until discontinued **Priority:** Routine

Routine

Question(s):

Indication:

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis** (Required)

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

VTE Risk and Prophylaxis Tool (Required)

Low Risk Definition	Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.	High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed.
Age less than 60 years and NO other VTE risk factors	One or more of the following medical conditions :	One or more of the following medical conditions :
Patient already adequately anticoagulated	CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome	Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)
	Age 60 and above	Severe fracture of hip, pelvis or leg
	Central line	Acute spinal cord injury with paresis
	History of DVT or family history of VTE	Multiple major traumas
	Anticipated length of stay GREATER than 48 hours	Abdominal or pelvic surgery for CANCER
	Less than fully and independently ambulatory	Acute ischemic stroke
	Estrogen therapy	History of PE
	Moderate or major surgery (not for cancer)	
	Major surgery within 3 months of admission	

Anticoagulation Guide for COVID patients ([https://formweb.com/files/houstonmethodist/documents/COVID-19 Anticoagulation Guideline - 8.20.2021v15.pdf](https://formweb.com/files/houstonmethodist/documents/COVID-19%20Anticoagulation%20Guideline%20-%208.20.2021v15.pdf))

☐ Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Required)

☐ Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)

Sign: _____ Printed Name: _____ Date/Time: _____
Page 33 of 72

☒ **Moderate risk of VTE** Frequency: Once Priority: Routine

☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once
Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:

☒ **Place sequential compression device**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)**

☒ **Moderate risk of VTE** Frequency: Once Priority: Routine

☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once
Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:

☒ **Place sequential compression device**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)**

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once
Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:

☒ **Place sequential compression device**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)**

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once
Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:

☒ **Place sequential compression device**

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **LOW Risk of VTE (Required)**

☒ **Low Risk (Required)**

☒ **Low risk of VTE** Frequency: Once Priority: Routine

Question(s):

Low risk: ☐ Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation ☐ Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation

☐ **MODERATE Risk of VTE - Surgical (Required)**

☒ **Moderate Risk (Required)**

☒ **Moderate risk of VTE** Frequency: Once Priority: Routine

☒ **Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Required)**

☐ **Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device**

☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis**

☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily Start Date: S+1

Admin Instructions:

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

☐ **heparin**

High Risk Bleeding Characteristics

Age $\geq$ 75

Weight < 50 kg

Unstable Hgb

Renal impairment

Plt count < 100 K/uL

Dual antiplatelet therapy

Active cancer

Cirrhosis/hepatic failure

Prior intra-cranial hemorrhage

Prior ischemic stroke

History of bleeding event requiring admission and/or transfusion

Chronic use of NSAIDs/steroids

Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units Frequency: every 12 hours scheduled

☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units Frequency: every 8 hours scheduled

☐ **Not high bleed risk**

☐ **Wt > 100 kg** Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ **warfarin (COUMADIN)**

☐ **WITHOUT pharmacy consult** Dose: 1 Route: oral Frequency: daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

Sign: _____ Printed Name: _____ Date/Time: _____

☐ Medications

☒ **Pharmacy consult to manage warfarin (COUMADIN)** Frequency: Until discontinued Priority:

Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

☐ Mechanical Prophylaxis (Required)

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ MODERATE Risk of VTE - Non-Surgical (Required)

☒ **Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient** (Required)

☒ **Moderate Risk** (Required)

☒ **Moderate risk of VTE** Frequency: Once Priority: Routine

☒ **Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient** (Required)

☐ **Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device**

☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority:

Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis**

☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin for Prophylactic Anticoagulation Nonsurgical** (Required)

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Sign: _____ Printed Name: _____ Date/Time: _____

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700 Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily

Admin Instructions:

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min

☐ **heparin**

High Risk Bleeding Characteristics
Age $\geq$ 75
Weight < 50 kg
Unstable Hgb
Renal impairment
Plt count < 100 K/uL
Dual antiplatelet therapy
Active cancer
Cirrhosis/hepatic failure
Prior intra-cranial hemorrhage
Prior ischemic stroke
History of bleeding event requiring admission and/or transfusion
Chronic use of NSAIDs/steroids
Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Sign: _____ Printed Name: _____ Date/Time: _____

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units Frequency: every 12 hours scheduled

☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units Frequency: every 8 hours scheduled

☐ **Not high bleed risk**

☐ **Wt > 100 kg** Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ **warfarin (COUMADIN)**

☐ **WITHOUT pharmacy consult** Dose: 1 Route: oral Frequency: daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

☒ **Pharmacy consult to manage warfarin (COUMADIN)** Frequency: Until discontinued

Priority: Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis (Required)**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **HIGH Risk of VTE - Surgical (Required)**

☒ **High Risk (Required)**

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **High Risk Pharmacological Prophylaxis - Surgical Patient (Required)**

☐ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Sign: _____ Printed Name: _____ Date/Time: _____

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily Start Date: S+1

Admin Instructions:

If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

☐ **heparin**

High Risk Bleeding Characteristics

Age $\geq$ 75

Weight < 50 kg

Unstable Hgb

Renal impairment

Plt count < 100 K/uL

Dual antiplatelet therapy

Active cancer

Cirrhosis/hepatic failure

Prior intra-cranial hemorrhage

Prior ischemic stroke

History of bleeding event requiring admission and/or transfusion

Chronic use of NSAIDs/steroids

Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

Sign: _____ Printed Name: _____ Date/Time: _____

- ☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units **Frequency:** every 12 hours scheduled
- ☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units **Frequency:** every 8 hours scheduled

☐ **Not high bleed risk**

- ☐ **Wt > 100 kg** Dose: 7500 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled
- ☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **warfarin (COUMADIN)**

- ☐ **WITHOUT pharmacy consult** Dose: 1 **Route:** oral **Frequency:** daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

- ☒ **Pharmacy consult to manage warfarin (COUMADIN)** **Frequency:** Until discontinued **Priority:**

Routine

Question(s):

Indication:

- ☐ **warfarin (COUMADIN) tablet** Dose: 1 **Route:** oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis (Required)**

- ☐ **Contraindications exist for mechanical prophylaxis** **Frequency:** Once **Priority:** Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

- ☒ **Place/Maintain sequential compression device continuous** **Frequency:** Continuous **Priority:** Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **HIGH Risk of VTE - Non-Surgical (Required)**

- ☒ **High Risk (Required)**

- ☒ **High risk of VTE** **Frequency:** Once **Priority:** Routine

- ☒ **High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Required)**

- ☐ **Contraindications exist for pharmacologic prophylaxis** **Frequency:** Once **Priority:** Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

- ☐ **Enoxaparin for Prophylactic Anticoagulation Nonsurgical (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

Sign: _____ Printed Name: _____ Date/Time: _____

☐ ENOXAPARIN 30 MG DAILY☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ ENOXAPARIN SQ DAILY☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily**Admin Instructions:**

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

☐ heparin**High Risk Bleeding Characteristics**Age $\geq$ 75

Weight < 50 kg

Unstable Hgb

Renal impairment

Plt count < 100 K/uL

Dual antiplatelet therapy

Active cancer

Cirrhosis/hepatic failure

Prior intra-cranial hemorrhage

Prior ischemic stroke

History of bleeding event requiring admission and/or transfusion

Chronic use of NSAIDs/steroids

Active GI ulcer

☐ High Bleed Risk

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units Frequency: every 12 hours scheduled☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units Frequency: every 8 hours scheduled☐ Not high bleed risk☐ **Wt > 100 kg** Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled☐ warfarin (COUMADIN)

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **WITHOUT pharmacy consult** Dose: 1 Route: oral Frequency: daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

☒ **Pharmacy consult to manage warfarin (COUMADIN)** Frequency: Until discontinued Priority:

Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis (Required)**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **HIGH Risk of VTE - Surgical (Hip/Knee) (Required)**

☒ **High Risk (Required)**

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Required)**

☐ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☐ **aspirin chewable tablet** Dose: 162 mg Frequency: daily Start Date: S+1

☐ **aspirin (ECOTRIN) enteric coated tablet** Dose: 162 mg Frequency: daily Start Date: S+1

☐ **Apixaban and Pharmacy Consult (Required)**

☒ **apixaban (ELIQUIS) tablet** Dose: 2.5 mg Frequency: 2 times daily Start Date: S+1

Question(s):

Indications: ☐ VTE prophylaxis

☒ **Pharmacy consult to monitor apixaban (ELIQUIS) therapy** Frequency: Until discontinued Priority:

STAT

Question(s):

Indications: VTE prophylaxis

☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Sign: _____ Printed Name: _____ Date/Time: _____

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily Start Date: S+1

Admin Instructions:

If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min

☐ **heparin**

High Risk Bleeding Characteristics

Age $\geq$ 75

Weight < 50 kg

Unstable Hgb

Renal impairment

Plt count < 100 K/uL

Dual antiplatelet therapy

Active cancer

Cirrhosis/hepatic failure

Prior intra-cranial hemorrhage

Prior ischemic stroke

History of bleeding event requiring admission and/or transfusion

Chronic use of NSAIDs/steroids

Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

Sign: _____ Printed Name: _____ Date/Time: _____

- ☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units Frequency: every 12 hours scheduled
- ☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units Frequency: every 8 hours scheduled
- ☐ **Not high bleed risk**
- ☐ **Wt > 100 kg** Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled
- ☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled
- ☐ **Rivaroxaban and Pharmacy Consult (Required)**
- ☒ **rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission** Dose: 10 mg Frequency: daily at 0600 (TIME CRITICAL)
- Question(s):**
Indications: ☐ VTE prophylaxis
- Admin Instructions:**
For Xarelto 15 mg and 20 mg, give with food or follow administration with enteral feeding to increase medication absorption. Do not administer via post-pyloric routes.
- ☒ **Pharmacy consult to monitor rivaroxaban (XARELTO) therapy** Frequency: Until discontinued Priority: STAT
- Question(s):**
Indications: VTE prophylaxis
- ☐ **warfarin (COUMADIN)**
- ☐ **WITHOUT pharmacy consult** Dose: 1 Route: oral Frequency: daily at 1700
- Question(s):**
Indication:
Dose Selection Guidance:
- ☐ **Medications**
- ☒ **Pharmacy consult to manage warfarin (COUMADIN)** Frequency: Until discontinued Priority: Routine
- Question(s):**
Indication:
- ☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral
- Question(s):**
Indication:
Dose Selection Guidance:
- ☐ **Mechanical Prophylaxis (Required)**
- ☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine
- Question(s):**
No mechanical VTE prophylaxis due to the following contraindication(s):
- ☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine
- Question(s):**
Side: Bilateral
Select Sleeve(s):

Sign: _____ Printed Name: _____ Date/Time: _____

VTE Risk and Prophylaxis Tool

Low Risk Definition	Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.	High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed.
Age less than 60 years and NO other VTE risk factors	One or more of the following medical conditions :	One or more of the following medical conditions :
Patient already adequately anticoagulated	CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome	Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)
	Age 60 and above	Severe fracture of hip, pelvis or leg
	Central line	Acute spinal cord injury with paresis
	History of DVT or family history of VTE	Multiple major traumas
	Anticipated length of stay GREATER than 48 hours	Abdominal or pelvic surgery for CANCER
	Less than fully and independently ambulatory	Acute ischemic stroke
	Estrogen therapy	History of PE
	Moderate or major surgery (not for cancer)	
	Major surgery within 3 months of admission	

Anticoagulation Guide for COVID patients ([https://formweb.com/files/houstonmethodist/documents/COVID-19 Anticoagulation Guideline - 8.20.2021v15.pdf](https://formweb.com/files/houstonmethodist/documents/COVID-19%20Anticoagulation%20Guideline%20-%208.20.2021v15.pdf))

☐ Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Required)

☐ Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **Moderate risk of VTE** Frequency: Once Priority: Routine

☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once
Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:

☒ **Place sequential compression device**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)**

☒ **Moderate risk of VTE** Frequency: Once Priority: Routine

☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once
Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:

☒ **Place sequential compression device**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)**

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once
Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:

☒ **Place sequential compression device**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)**

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once
Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:

☒ **Place sequential compression device**

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **LOW Risk of VTE (Required)**

☒ **Low Risk (Required)**

☒ **Low risk of VTE** Frequency: Once Priority: Routine

Question(s):

Low risk: ☐ Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation ☐ Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation

☐ **Moderate Risk of VTE - Surgical (Required)**

☒ **Moderate Risk (Required)**

☒ **Moderate risk of VTE** Frequency: Once Priority: Routine

☒ **Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Required)**

☐ **Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device**

☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis**

☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily Start Date: S+1

Admin Instructions:

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

☐ **heparin**

High Risk Bleeding Characteristics

Age $\geq$ 75

Weight < 50 kg

Unstable Hgb

Renal impairment

Plt count < 100 K/uL

Dual antiplatelet therapy

Active cancer

Cirrhosis/hepatic failure

Prior intra-cranial hemorrhage

Prior ischemic stroke

History of bleeding event requiring admission and/or transfusion

Chronic use of NSAIDs/steroids

Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units Frequency: every 12 hours scheduled

☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units Frequency: every 8 hours scheduled

☐ **Not high bleed risk**

☐ **Wt > 100 kg** Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ **warfarin (COUMADIN)**

☐ **WITHOUT pharmacy consult** Dose: 1 Route: oral Frequency: daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

Sign: _____ Printed Name: _____ Date/Time: _____

☐ Medications☒ **Pharmacy consult to manage warfarin (COUMADIN)** Frequency: Until discontinued Priority:

Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis (Required)**☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **Moderate Risk of VTE - Non-Surgical (Required)**☒ **Moderate Risk (Required)**☒ **Moderate risk of VTE** Frequency: Once Priority: Routine☒ **Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Required)**☐ **Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device**☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis**☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin for Prophylactic Anticoagulation Nonsurgical (Required)****Patient renal status: @CRCL@**

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

Sign: _____ Printed Name: _____ Date/Time: _____

☐ ENOXAPARIN 30 MG DAILY☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ ENOXAPARIN SQ DAILY☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily**Admin Instructions:**

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min

☐ heparin**High Risk Bleeding Characteristics**Age $\geq$ 75

Weight < 50 kg

Unstable Hgb

Renal impairment

Plt count < 100 K/uL

Dual antiplatelet therapy

Active cancer

Cirrhosis/hepatic failure

Prior intra-cranial hemorrhage

Prior ischemic stroke

History of bleeding event requiring admission and/or transfusion

Chronic use of NSAIDs/steroids

Active GI ulcer

☐ High Bleed Risk

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units Frequency: every 12 hours scheduled☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units Frequency: every 8 hours scheduled☐ Not high bleed risk☐ **Wt > 100 kg** Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled☐ warfarin (COUMADIN)

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **WITHOUT pharmacy consult** Dose: 1 Route: oral Frequency: daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

☒ **Pharmacy consult to manage warfarin (COUMADIN)** Frequency: Until discontinued Priority:

Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis** (Required)

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **High Risk of VTE - Surgical** (Required)

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

☒ **High Risk** (Required)

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **High Risk Pharmacological Prophylaxis - Surgical Patient** (Required)

☐ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation** (Required)

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily Start Date: S+1

Admin Instructions:

If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

☐ **heparin**

High Risk Bleeding Characteristics
Age $\geq$ 75
Weight < 50 kg
Unstable Hgb
Renal impairment
Plt count < 100 K/uL
Dual antiplatelet therapy
Active cancer
Cirrhosis/hepatic failure
Prior intra-cranial hemorrhage
Prior ischemic stroke
History of bleeding event requiring admission and/or transfusion
Chronic use of NSAIDs/steroids
Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units Frequency: every 12 hours scheduled

☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units Frequency: every 8 hours scheduled

☐ **Not high bleed risk**

☐ **Wt > 100 kg** Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ **warfarin (COUMADIN)**

☐ **WITHOUT pharmacy consult** Dose: 1 Route: oral Frequency: daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

Sign: _____ Printed Name: _____ Date/Time: _____

☐ Medications☒ **Pharmacy consult to manage warfarin (COUMADIN)** Frequency: Until discontinued Priority:

Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

☐ High Risk of VTE - Non-Surgical (Required)

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

☒ High Risk (Required)☒ **High risk of VTE** Frequency: Once Priority: Routine☒ **High Risk Pharmacological Prophylaxis - Non-Surgical Patient** (Required)☐ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin for Prophylactic Anticoagulation Nonsurgical** (Required)**Patient renal status:** @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ ENOXAPARIN 30 MG DAILY☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ ENOXAPARIN SQ DAILY☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily**Admin Instructions:**

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

Sign: _____ Printed Name: _____ Date/Time: _____

☐ heparin

High Risk Bleeding Characteristics
Age $\geq$ 75
Weight < 50 kg
Unstable Hgb
Renal impairment
Plt count < 100 K/uL
Dual antiplatelet therapy
Active cancer
Cirrhosis/hepatic failure
Prior intra-cranial hemorrhage
Prior ischemic stroke
History of bleeding event requiring admission and/or transfusion
Chronic use of NSAIDs/steroids
Active GI ulcer

☐ High Bleed Risk

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ HEParin (porcine) injection - Q12 Hours Dose: 5000 Units Frequency: every 12 hours scheduled

☐ HEParin (porcine) injection - Q8 Hours Dose: 5000 Units Frequency: every 8 hours scheduled

☐ Not high bleed risk

☐ Wt > 100 kg Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ Wt LESS than or equal to 100 kg Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ warfarin (COUMADIN)

☐ WITHOUT pharmacy consult Dose: 1 Route: oral Frequency: daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ Medications

☒ Pharmacy consult to manage warfarin (COUMADIN) Frequency: Until discontinued Priority:

Routine

Question(s):

Indication:

☐ warfarin (COUMADIN) tablet Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

☐ High Risk of VTE - Surgical (Hip/Knee) (Required)

Address both pharmacologic and mechanical prophylaxis by ordering from Pharmacological and Mechanical Prophylaxis.

☒ High Risk (Required)

☒ High risk of VTE Frequency: Once Priority: Routine

☒ High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Required)

☐ Contraindications exist for pharmacologic prophylaxis Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

Sign: _____ Printed Name: _____ Date/Time: _____

- ☐ **aspirin chewable tablet** Dose: 162 mg Frequency: daily Start Date: S+1
- ☐ **aspirin (ECOTRIN) enteric coated tablet** Dose: 162 mg Frequency: daily Start Date: S+1
- ☐ **Apixaban and Pharmacy Consult** (Required)
- ☒ **apixaban (ELIQUIS) tablet** Dose: 2.5 mg Frequency: 2 times daily Start Date: S+1
- Question(s):**
Indications: ○ VTE prophylaxis
- ☒ **Pharmacy consult to monitor apixaban (ELIQUIS) therapy** Frequency: Until discontinued Priority: STAT
- Question(s):**
Indications: VTE prophylaxis
- ☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation** (Required)
- Patient renal status:** @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

- ☐ **ENOXAPARIN 30 MG DAILY**
- ☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700
- Start Date:** S+1
- Question(s):**
Indication(s):
- Product Admin Instructions:**
Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.
- ☐ **ENOXAPARIN SQ DAILY**
- ☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1
- Question(s):**
Indication(s):
- Product Admin Instructions:**
Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.
- ☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily Start Date: S+1
- Admin Instructions:**
If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min
- ☐ **heparin**

Sign: _____ Printed Name: _____ Date/Time: _____

High Risk Bleeding Characteristics
Age $\geq$ 75
Weight < 50 kg
Unstable Hgb
Renal impairment
Plt count < 100 K/uL
Dual antiplatelet therapy
Active cancer
Cirrhosis/hepatic failure
Prior intra-cranial hemorrhage
Prior ischemic stroke
History of bleeding event requiring admission and/or transfusion
Chronic use of NSAIDs/steroids
Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours Dose:** 5000 Units **Frequency:** every 12 hours scheduled

☐ **HEParin (porcine) injection - Q8 Hours Dose:** 5000 Units **Frequency:** every 8 hours scheduled

☐ **Not high bleed risk**

☐ **Wt > 100 kg Dose:** 7500 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **Wt LESS than or equal to 100 kg Dose:** 5000 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **Rivaroxaban and Pharmacy Consult (Required)**

☒ **rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission Dose:** 10 mg **Frequency:** daily at 0600 (TIME CRITICAL)

Question(s):

Indications: ☐ VTE prophylaxis

Admin Instructions:

For Xarelto 15 mg and 20 mg, give with food or follow administration with enteral feeding to increase medication absorption. Do not administer via post-pyloric routes.

☒ **Pharmacy consult to monitor rivaroxaban (XARELTO) therapy Frequency:** Until discontinued **Priority:** STAT

Question(s):

Indications: VTE prophylaxis

☐ **warfarin (COUMADIN)**

☐ **WITHOUT pharmacy consult Dose:** 1 **Route:** oral **Frequency:** daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

☒ **Pharmacy consult to manage warfarin (COUMADIN) Frequency:** Until discontinued **Priority:** Routine

Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

VTE Risk and Prophylaxis Tool

Low Risk Definition	Moderate Risk Definition Pharmacologic prophylaxis must be addressed. Mechanical prophylaxis is optional unless pharmacologic is contraindicated.	High Risk Definition Both pharmacologic AND mechanical prophylaxis must be addressed.
Age less than 60 years and NO other VTE risk factors	One or more of the following medical conditions :	One or more of the following medical conditions :
Patient already adequately anticoagulated	CHF, MI, lung disease, pneumonia, active inflammation, dehydration, varicose veins, cancer, sepsis, obesity, previous stroke, rheumatologic disease, sickle cell disease, leg swelling, ulcers, venous stasis and nephrotic syndrome	Thrombophilia (Factor V Leiden, prothrombin variant mutations, anticardiolipin antibody syndrome; antithrombin, protein C or protein S deficiency; hyperhomocysteinemia; myeloproliferative disorders)
	Age 60 and above	Severe fracture of hip, pelvis or leg
	Central line	Acute spinal cord injury with paresis
	History of DVT or family history of VTE	Multiple major traumas
	Anticipated length of stay GREATER than 48 hours	Abdominal or pelvic surgery for CANCER
	Less than fully and independently ambulatory	Acute ischemic stroke
	Estrogen therapy	History of PE
	Moderate or major surgery (not for cancer)	
	Major surgery within 3 months of admission	

Sign: _____ Printed Name: _____

Date/Time: _____

Anticoagulation Guide for COVID patients ([https://formweb.com/files/houstonmethodist/documents/COVID-19 Anticoagulation Guideline - 8.20.2021v15.pdf](https://formweb.com/files/houstonmethodist/documents/COVID-19%20Anticoagulation%20Guideline%20-%208.20.2021v15.pdf))

☐ Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis with Risk Stratification (Required)

☐ Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)

☒ Moderate risk of VTE Frequency: Once Priority: Routine

☒ Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

☒ Place sequential compression device

☐ Contraindications exist for mechanical prophylaxis Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ Place/Maintain sequential compression device continuous Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ Moderate Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)

☒ Moderate risk of VTE Frequency: Once Priority: Routine

☒ Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

☒ Place sequential compression device

☐ Contraindications exist for mechanical prophylaxis Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ Place/Maintain sequential compression device continuous Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)

☒ High risk of VTE Frequency: Once Priority: Routine

☒ Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication. Therapy for the following:

☒ Place sequential compression device

☐ Contraindications exist for mechanical prophylaxis Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ Place/Maintain sequential compression device continuous Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ High Risk - Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis (Required)

☒ High risk of VTE Frequency: Once Priority: Routine

Sign: _____ Printed Name: _____ Date/Time: _____

☒ **Patient currently has an active order for therapeutic anticoagulant or VTE prophylaxis** Frequency: Once

Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis because: patient is already on therapeutic anticoagulation for other indication.
Therapy for the following:

☒ **Place sequential compression device**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **LOW Risk of VTE (Required)**

☒ **Low Risk (Required)**

☒ **Low risk of VTE** Frequency: Once Priority: Routine

Question(s):

Low risk: ☐ Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation ☐ Due to low risk, no VTE prophylaxis is needed. Will encourage early ambulation

☐ **MODERATE Risk of VTE - Surgical (Required)**

☒ **Moderate Risk (Required)**

☒ **Moderate risk of VTE** Frequency: Once Priority: Routine

☒ **Moderate Risk Pharmacological Prophylaxis - Surgical Patient (Required)**

☐ **Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device**

☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis**

☒ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☒ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Sign: _____ Printed Name: _____ Date/Time: _____

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily Start Date: S+1

Admin Instructions:

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

☐ **heparin**

High Risk Bleeding Characteristics

Age $\geq$ 75

Weight < 50 kg

Unstable Hgb

Renal impairment

Plt count < 100 K/uL

Dual antiplatelet therapy

Active cancer

Cirrhosis/hepatic failure

Prior intra-cranial hemorrhage

Prior ischemic stroke

History of bleeding event requiring admission and/or transfusion

Chronic use of NSAIDs/steroids

Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

Sign: _____ Printed Name: _____ Date/Time: _____

- ☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units **Frequency:** every 12 hours scheduled
- ☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units **Frequency:** every 8 hours scheduled
- ☐ **Not high bleed risk**
- ☐ **Wt > 100 kg** Dose: 7500 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled
- ☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **warfarin (COUMADIN)**

- ☐ **WITHOUT pharmacy consult** Dose: 1 **Route:** oral **Frequency:** daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

- ☐ **Medications**

- ☒ **Pharmacy consult to manage warfarin (COUMADIN)** **Frequency:** Until discontinued **Priority:**

Routine

Question(s):

Indication:

- ☐ **warfarin (COUMADIN) tablet** Dose: 1 **Route:** oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis (Required)**

- ☐ **Contraindications exist for mechanical prophylaxis** **Frequency:** Once **Priority:** Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

- ☒ **Place/Maintain sequential compression device continuous** **Frequency:** Continuous **Priority:** Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **MODERATE Risk of VTE - Non-Surgical (Required)**

- ☒ **Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Required)**

- ☒ **Moderate Risk (Required)**

- ☒ **Moderate risk of VTE** **Frequency:** Once **Priority:** Routine

- ☒ **Moderate Risk Pharmacological Prophylaxis - Non-Surgical Patient (Required)**

- ☐ **Contraindications exist for pharmacologic prophylaxis - Order Sequential compression device**

- ☒ **Contraindications exist for pharmacologic prophylaxis** **Frequency:** Once **Priority:** Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

- ☒ **Place/Maintain sequential compression device continuous** **Frequency:** Continuous **Priority:**

Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

- ☐ **Contraindications exist for pharmacologic prophylaxis AND mechanical prophylaxis**

- ☒ **Contraindications exist for pharmacologic prophylaxis** **Frequency:** Once **Priority:** Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

- ☒ **Contraindications exist for mechanical prophylaxis** **Frequency:** Once **Priority:** Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin for Prophylactic Anticoagulation Nonsurgical (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily
at 1700 Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily

Admin Instructions:

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT), do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min

☐ **heparin**

High Risk Bleeding Characteristics
Age $\geq$ 75
Weight < 50 kg
Unstable Hgb
Renal impairment
Plt count < 100 K/uL
Dual antiplatelet therapy
Active cancer
Cirrhosis/hepatic failure
Prior intra-cranial hemorrhage
Prior ischemic stroke
History of bleeding event requiring admission and/or transfusion
Chronic use of NSAIDs/steroids
Active GI ulcer

☐ **High Bleed Risk**

Sign: _____ Printed Name: _____ Date/Time: _____

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units Frequency: every 12 hours scheduled

☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units Frequency: every 8 hours scheduled

☐ **Not high bleed risk**

☐ **Wt > 100 kg** Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled

☐ **warfarin (COUMADIN)**

☐ **WITHOUT pharmacy consult** Dose: 1 Route: oral Frequency: daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

☒ **Pharmacy consult to manage warfarin (COUMADIN)** Frequency: Until discontinued

Priority: Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis (Required)**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **HIGH Risk of VTE - Surgical (Required)**

☒ **High Risk (Required)**

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **High Risk Pharmacological Prophylaxis - Surgical Patient (Required)**

☐ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Sign: _____ Printed Name: _____ Date/Time: _____

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily Start Date: S+1

Admin Instructions:

If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

☐ **heparin**

High Risk Bleeding Characteristics

Age $\geq$ 75

Weight < 50 kg

Unstable Hgb

Renal impairment

Plt count < 100 K/uL

Dual antiplatelet therapy

Active cancer

Cirrhosis/hepatic failure

Prior intra-cranial hemorrhage

Prior ischemic stroke

History of bleeding event requiring admission and/or transfusion

Chronic use of NSAIDs/steroids

Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

Sign: _____ Printed Name: _____ Date/Time: _____

- ☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units **Frequency:** every 12 hours scheduled
- ☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units **Frequency:** every 8 hours scheduled
- ☐ **Not high bleed risk**
- ☐ **Wt > 100 kg** Dose: 7500 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled
- ☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units **Route:** subcutaneous **Frequency:** every 8 hours scheduled

☐ **warfarin (COUMADIN)**

- ☐
- WITHOUT pharmacy consult**
- Dose: 1
- Route:**
- oral
- Frequency:**
- daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

- ☒
- Pharmacy consult to manage warfarin (COUMADIN)**
- Frequency:**
- Until discontinued
- Priority:**

Routine

Question(s):

Indication:

- ☐
- warfarin (COUMADIN) tablet**
- Dose: 1
- Route:**
- oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis (Required)**

- ☐
- Contraindications exist for mechanical prophylaxis**
- Frequency:**
- Once
- Priority:**
- Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

- ☒
- Place/Maintain sequential compression device continuous**
- Frequency:**
- Continuous
- Priority:**
- Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **HIGH Risk of VTE - Non-Surgical (Required)**

- ☒
- High Risk (Required)**

- ☒
- High risk of VTE**
- Frequency:**
- Once
- Priority:**
- Routine

- ☒
- High Risk Pharmacological Prophylaxis - Non-Surgical Patient (Required)**

- ☐
- Contraindications exist for pharmacologic prophylaxis**
- Frequency:**
- Once
- Priority:**
- Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

- ☐
- Enoxaparin for Prophylactic Anticoagulation Nonsurgical (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

Sign: _____ Printed Name: _____ Date/Time: _____

☐ ENOXAPARIN 30 MG DAILY☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ ENOXAPARIN SQ DAILY☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily**Admin Instructions:**

If the patient does not have a history of or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min.

☐ heparin**High Risk Bleeding Characteristics**Age $\geq$ 75

Weight < 50 kg

Unstable Hgb

Renal impairment

Plt count < 100 K/uL

Dual antiplatelet therapy

Active cancer

Cirrhosis/hepatic failure

Prior intra-cranial hemorrhage

Prior ischemic stroke

History of bleeding event requiring admission and/or transfusion

Chronic use of NSAIDs/steroids

Active GI ulcer

☐ High Bleed Risk

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units Frequency: every 12 hours scheduled☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units Frequency: every 8 hours scheduled☐ Not high bleed risk☐ **Wt > 100 kg** Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled☐ warfarin (COUMADIN)

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **WITHOUT pharmacy consult** Dose: 1 Route: oral Frequency: daily at 1700

Question(s):

Indication:

Dose Selection Guidance:

☐ **Medications**

☒ **Pharmacy consult to manage warfarin (COUMADIN)** Frequency: Until discontinued Priority:

Routine

Question(s):

Indication:

☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral

Question(s):

Indication:

Dose Selection Guidance:

☐ **Mechanical Prophylaxis (Required)**

☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine

Question(s):

No mechanical VTE prophylaxis due to the following contraindication(s):

☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine

Question(s):

Side: Bilateral

Select Sleeve(s):

☐ **HIGH Risk of VTE - Surgical (Hip/Knee) (Required)**

☒ **High Risk (Required)**

☒ **High risk of VTE** Frequency: Once Priority: Routine

☒ **High Risk Pharmacological Prophylaxis - Hip or Knee (Arthroplasty) Surgical Patient (Required)**

☐ **Contraindications exist for pharmacologic prophylaxis** Frequency: Once Priority: Routine

Question(s):

No pharmacologic VTE prophylaxis due to the following contraindication(s):

☐ **aspirin chewable tablet** Dose: 162 mg Frequency: daily Start Date: S+1

☐ **aspirin (ECOTRIN) enteric coated tablet** Dose: 162 mg Frequency: daily Start Date: S+1

☐ **Apixaban and Pharmacy Consult (Required)**

☒ **apixaban (ELIQUIS) tablet** Dose: 2.5 mg Frequency: 2 times daily Start Date: S+1

Question(s):

Indications: ☐ VTE prophylaxis

☒ **Pharmacy consult to monitor apixaban (ELIQUIS) therapy** Frequency: Until discontinued Priority:

STAT

Question(s):

Indications: VTE prophylaxis

☐ **Enoxaparin (LOVENOX) for Prophylactic Anticoagulation (Required)**

Patient renal status: @CRCL@

For patients with CrCl GREATER than or EQUAL to 30mL/min, enoxaparin orders will apply the following recommended doses by weight:

Sign: _____ Printed Name: _____ Date/Time: _____

Weight	Dose
LESS THAN 100kg	enoxaparin 40mg daily
100 to 139kg	enoxaparin 30mg every 12 hours
GREATER THAN or EQUAL to 140kg	enoxaparin 40mg every 12 hours

☐ **ENOXAPARIN 30 MG DAILY**

☒ **enoxaparin (LOVENOX) injection** Dose: 30 mg Route: subcutaneous Frequency: daily at 1700

Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **ENOXAPARIN SQ DAILY**

☒ **enoxaparin (LOVENOX) injection** Route: subcutaneous Start Date: S+1

Question(s):

Indication(s):

Product Admin Instructions:

Administer by deep subcutaneous injection into the left and right anterolateral or posterolateral abdominal wall. Alternate injection site with each administration.

☐ **fondaparinux (ARIXTRA) injection** Dose: 2.5 mg Route: subcutaneous Frequency: daily Start Date: S+1

Admin Instructions:

If the patient does not have a history or suspected case of Heparin-Induced Thrombocytopenia (HIT) do NOT order this medication. Contraindicated in patients LESS than 50kg, prior to surgery/invasive procedure, or CrCl LESS than 30 mL/min

☐ **heparin**

High Risk Bleeding Characteristics

Age $\geq$ 75

Weight < 50 kg

Unstable Hgb

Renal impairment

Plt count < 100 K/uL

Dual antiplatelet therapy

Active cancer

Cirrhosis/hepatic failure

Prior intra-cranial hemorrhage

Prior ischemic stroke

History of bleeding event requiring admission and/or transfusion

Chronic use of NSAIDs/steroids

Active GI ulcer

☐ **High Bleed Risk**

Every 12 hour frequency is appropriate for most high bleeding risk patients. However, some high bleeding risk patients also have high clotting risk in which every 8 hour frequency may be clinically appropriate.

Please weight the risks/benefits of bleeding and clotting when selecting the dosing frequency.

Sign: _____ Printed Name: _____ Date/Time: _____

- ☐ **HEParin (porcine) injection - Q12 Hours** Dose: 5000 Units Frequency: every 12 hours scheduled
- ☐ **HEParin (porcine) injection - Q8 Hours** Dose: 5000 Units Frequency: every 8 hours scheduled
- ☐ **Not high bleed risk**
- ☐ **Wt > 100 kg** Dose: 7500 Units Route: subcutaneous Frequency: every 8 hours scheduled
- ☐ **Wt LESS than or equal to 100 kg** Dose: 5000 Units Route: subcutaneous Frequency: every 8 hours scheduled
- ☐ **Rivaroxaban and Pharmacy Consult (Required)**
- ☒ **rivaroxaban (XARELTO) tablet for hip or knee arthroplasty planned during this admission** Dose: 10 mg Frequency: daily at 0600 (TIME CRITICAL)
- Question(s):**
Indications: ☐ VTE prophylaxis
- Admin Instructions:**
For Xarelto 15 mg and 20 mg, give with food or follow administration with enteral feeding to increase medication absorption. Do not administer via post-pyloric routes.
- ☒ **Pharmacy consult to monitor rivaroxaban (XARELTO) therapy** Frequency: Until discontinued Priority: STAT
- Question(s):**
Indications: VTE prophylaxis
- ☐ **warfarin (COUMADIN)**
- ☐ **WITHOUT pharmacy consult** Dose: 1 Route: oral Frequency: daily at 1700
- Question(s):**
Indication:
Dose Selection Guidance:
- ☐ **Medications**
- ☒ **Pharmacy consult to manage warfarin (COUMADIN)** Frequency: Until discontinued Priority: Routine
- Question(s):**
Indication:
- ☐ **warfarin (COUMADIN) tablet** Dose: 1 Route: oral
- Question(s):**
Indication:
Dose Selection Guidance:
- ☐ **Mechanical Prophylaxis (Required)**
- ☐ **Contraindications exist for mechanical prophylaxis** Frequency: Once Priority: Routine
- Question(s):**
No mechanical VTE prophylaxis due to the following contraindication(s):
- ☒ **Place/Maintain sequential compression device continuous** Frequency: Continuous Priority: Routine
- Question(s):**
Side: Bilateral
Select Sleeve(s):

Labs Today**Hematology/Coagulation**

- ☐ **Hemoglobin and hematocrit** Frequency: Once Priority: Routine Specimen Type: Blood Maximum Quantity: 3
- ☐ **CBC with platelet and differential** Frequency: Once Priority: Routine Specimen Type: Blood Maximum Quantity: 3
- ☐ **Prothrombin time with INR** Frequency: Once Priority: Routine Specimen Type: Blood Maximum Quantity: 3
- ☐ **Partial thromboplastin time** Frequency: Once Priority: Routine Specimen Type: Blood Maximum Quantity: 3

Primary Ordering Comments:

Do not draw blood from the arm that has heparin infusion. Do not draw from heparin flushed lines. If there is no other access other than the heparin line, then stop the heparin, flush the line, and aspirate 20 ml of blood to waste prior to drawing a specimen.

Chemistry

- ☐ **Basic metabolic panel** Frequency: Once Priority: Routine Specimen Type: Blood Maximum Quantity: 3

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Magnesium** Frequency: Once Priority: Routine Specimen Type: Blood Maximum Quantity: 3

☐ **Calcium** Frequency: Once Priority: Routine Specimen Type: Blood Maximum Quantity: 3

Labs Tomorrow

Hematology/Coagulation

☐ **Hemoglobin and hematocrit** Frequency: AM draw Frequency Limit: 1 Occurrences Priority: Routine Specimen Type: Blood Maximum Quantity: 3

☐ **CBC with platelet and differential** Frequency: AM draw Frequency Limit: 1 Occurrences Priority: Routine Specimen Type: Blood Maximum Quantity: 3

☐ **Prothrombin time with INR** Frequency: AM draw Frequency Limit: 1 Occurrences Priority: Routine Specimen Type: Blood Maximum Quantity: 3

☐ **Partial thromboplastin time** Frequency: AM draw Frequency Limit: 1 Occurrences Priority: Routine Specimen Type: Blood Maximum Quantity: 3

Primary Ordering Comments:

Do not draw blood from the arm that has heparin infusion. Do not draw from heparin flushed lines. If there is no other access other than the heparin line, then stop the heparin, flush the line, and aspirate 20 ml of blood to waste prior to drawing a specimen.

Chemistry

☐ **Basic metabolic panel** Frequency: AM draw Frequency Limit: 1 Occurrences Priority: Routine Specimen Type: Blood Maximum Quantity: 3

☐ **Magnesium** Frequency: AM draw Frequency Limit: 1 Occurrences Priority: Routine Specimen Type: Blood Maximum Quantity: 3

☐ **Calcium** Frequency: AM draw Frequency Limit: 1 Occurrences Priority: Routine Specimen Type: Blood Maximum Quantity: 3

Cardiology

Imaging

Other Studies

Respiratory

Respiratory

☐ **Incentive spirometry instructions** Frequency: Once Frequency Limit: 1 Occurrences Priority: Routine

Question(s):

Frequency of use: ○ Every hour, 10 times per hour

Rehab

Consults

For Physician Consult orders use sidebar

Ancillary Consults (For Physician Consults, use the Sidebar)

☐ **Consult to case management** Frequency: Once Priority: Routine

Question(s):

Consult Reason:

Reason for Consult?

Process Instructions:

If Ordering IV antimicrobial therapy, an additional consult to Case Management OPAT order is needed.

☐ **Consult to social work for discharge planning** Frequency: Once Priority: Routine

Question(s):

Reason for Consult: ○ Discharge Planning

Reason for Consult?

☐ **PT eval and treat** Frequency: Once Priority: Routine

Question(s):

Reasons for referral to Physical Therapy (mark all applicable):

Are there any restrictions for positioning or mobility?

Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal):

Weight Bearing Status:

Reason for PT?

Process Instructions:

If the patient is not medically/surgically stable for therapy, please obtain the necessary clearance prior to consulting physical therapy

If the patient currently has an order for bed rest, please consider revising the activity order to accommodate therapy

Sign: _____ Printed Name: _____ Date/Time: _____

☐ **Consult to PT Wound Care Eval and Treat** Frequency: Once Priority: Routine

Question(s):

Special Instructions:

Location of Wound?

Reason for PT?

☐ **OT eval and treat** Frequency: Once Priority: Routine

Question(s):

Reason for referral to Occupational Therapy (mark all that apply):

Are there any restrictions for positioning or mobility?

Please provide safe ranges for HR, BP, O2 saturation(if values are very abnormal):

Weight Bearing Status:

Reason for OT?

Process Instructions:

If the patient is not medically/surgically stable for therapy, please obtain the necessary clearance prior to consulting occupational therapy

If the patient currently has an order for bed rest, please consider revising the activity order to accommodate therapy.

☐ **Consult to Nutrition** Frequency: Once Priority: Routine

Question(s):

Reason For Consult?

Purpose/Topic:

Reason for Consult?

☐ **Consult to Respiratory Therapy** Frequency: Once Priority: Routine

Question(s):

Reason for Consult?

Reason for Consult?

☐ **Consult to Spiritual Care** Frequency: Once Priority: Routine

Question(s):

Reason for consult?

Reason for Consult?

Process Instructions:

For requests after hours, call the house operator.

☐ **Consult to Speech Language Pathology** Frequency: Once Priority: Routine

Question(s):

Reason for consult:

Reason for SLP?

☐ **Consult to Wound Ostomy Care Nurse** Frequency: Once Priority: Routine

Question(s):

Reason for consult:

Reason for consult:

Reason for consult:

Reason for consult:

Consult for NPWT:

Reason for consult:

Reason for consult:

Reason for Consult?

Process Instructions:

This is NOT for PT Wound Care Consult order.

Additional Orders